



Swansea University  
Prifysgol Abertawe



## Cronfa - Swansea University Open Access Repository

---

This is an author produced version of a paper published in :

*Cortex*

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa23883>

---

### **Paper:**

Ktori, M., Tree, J., Mousikou, P., Coltheart, M. & Rastle, K. (2016). Prefixes repel stress in reading aloud: Evidence from surface dyslexia. *Cortex*, 74, 191-205.

<http://dx.doi.org/10.1016/j.cortex.2015.10.009>

---

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

# Accepted Manuscript

Prefixes Repel Stress in Reading Aloud: Evidence from Surface Dyslexia

Maria Ktori, Jeremy J. Tree, Petroula Mousikou, Max Coltheart, Kathleen Rastle



PII: S0010-9452(15)00360-3

DOI: [10.1016/j.cortex.2015.10.009](https://doi.org/10.1016/j.cortex.2015.10.009)

Reference: CORTEX 1606

To appear in: *Cortex*

Received Date: 13 May 2015

Revised Date: 4 September 2015

Accepted Date: 20 October 2015

Please cite this article as: Ktori M, Tree JJ, Mousikou P, Coltheart M, Rastle K, Prefixes Repel Stress in Reading Aloud: Evidence from Surface Dyslexia, *CORTEX* (2015), doi: 10.1016/j.cortex.2015.10.009.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Maria Ktori<sup>1</sup>, Jeremy J. Tree<sup>2</sup>, Petroula Mousikou<sup>1</sup>, Max Coltheart<sup>3</sup>, & Kathleen Rastle<sup>1</sup>

1. Department of Psychology, Royal Holloway, University of London, UK
2. Department of Psychology, Swansea University, UK
3. ARC Centre of Excellence in Cognition and its Disorders, and Department of Cognitive Science, Macquarie University, Australia

#### Author Note

This research was supported by The Leverhulme Trust RPG 2013-024 awarded to Kathleen Rastle, Max Coltheart, Jeremy Tree, and Petroula Mousikou.

Correspondence concerning this article should be addressed to Maria Ktori, Department of Psychology, Royal Holloway, University of London, Egham, Surrey, TW20 0EX, U.K. E-mail: Maria.Ktori@rhul.ac.uk

during reading aloud English disyllabic words. In particular, we tested the hypothesis that prefixes repel stress (Rastle & Coltheart, 2000) by investigating the likelihood with which patients with surface dyslexia assign second-syllable stress to prefixed words. Five such patients were presented with three types of disyllabic words for reading aloud: ‘regular’ prefixed words with weak-strong stress pattern (e.g., remind); ‘irregular’ prefixed words with strong-weak stress pattern (e.g., reflex); and non-prefixed words with strong-weak stress pattern (e.g., scandal). Results showed that all five patients frequently regularized the strong-weak prefixed words by pronouncing them with second syllable stress. These regularization errors provide strong evidence for the functional role of prefixes in stress assignment during reading. Additional computational simulations using the rule-based algorithm for pronouncing disyllables developed by Rastle and Coltheart (2000) and the CDP++ model of reading aloud (Perry et al., 2010) allowed us to evaluate how these two opponent approaches to reading aloud fare in respect of the patient data.

*Keywords:* reading aloud, prefixes, surface dyslexia, computational modeling

Over the past couple of decades, research into the generation of sound from print has begun to move away from a focus on simple monosyllabic words, to consider the special problems posed by multisyllabic words (e.g., Arciuli, Monaghan, & Ševa, 2010; Rastle & Coltheart, 2000). Reading aloud a multisyllabic word requires more than the translation of an orthographic string to its phonological equivalent; it also requires the assignment of stress, which involves the phonetic accentuation of one of the syllables, along with the possible reduction of an unstressed vowel in the word. A clear illustration of these phonetic modulations can be seen in the case of noun/verb minimal pairs. For example, the disyllabic English word “suspect” is pronounced /ˈsʌspɛkt/ when used as a noun (e.g., the usual suspect) and /səˈspɛkt/ when used as a verb (e.g., to suspect foul play). While the pronunciation of the former is characterised by a first-syllable stress and two phonetically full vowels, the pronunciation of the latter is characterized by a second-syllable stress and the phonetic reduction (schwa) of the vowel in the first syllable.

Several recent studies have investigated the mental processes that underpin stress assignment during reading aloud. These studies have focused on languages characterised by a free-stress system such as English (e.g., Arciuli & Cupples, 2006; 2007; Guion, Clark, Harada, & Wayland, 2003), Italian (see Sulpizio, Burani, & Colombo, 2015 for a review), and Russian (Jouravlev & Lupker, 2014), where stress has neither a fixed position within the word nor is marked by the use of diacritics. These investigations have mainly sought to examine the extent to which stress is determined by word specific stored information (lexical) or statistical-distributional regularities of a given language (sublexical). In respect of this latter dimension, several factors have been identified as potential predictors of stress

orthographic sequences, in particular word beginnings and/or endings (e.g., Burani, Falzi, & Sulpizio, 2014; Cappa, Nespor, Ielasi, & Miozzo, 1997; Colombo, 1992; Ševa, Monaghan & Arciuli, 2009); syllabic weight both at the orthographic (Kelly, 2004; Kelly, Morris, & Verrekia, 1998) and phonological level (Guion et al., 2003); and vowel length (Baker & Smith, 1976; Guion et al., 2003). Of particular importance to the present study is the claim that the morphological structure of a word (i.e., the presence of affixes) also provides important information in determining stress assignment in reading aloud (Rastle & Coltheart, 2000).

Rastle and Coltheart (2000) were among the first researchers to explore the computational processes of stress assignment during the spelling-to-sound translation of a disyllabic stimulus, and to demonstrate how these mechanisms could be implemented within an existing theoretical framework of reading, namely the DRC model (Coltheart, Curtis, Atkins, & Haller, 1993; Coltheart & Rastle, 1994; Coltheart, Rastle, Perry, Langdon, & Ziegler, 2001; Rastle & Coltheart, 1999). The DRC model is a computational instantiation of the dual-route theory of reading, the central tenet of which is that the translation of spelling to sound involves two procedures, a lexical procedure whereby item-specific stored knowledge about the relationship between orthography and phonology is retrieved, and a sublexical procedure whereby phonological information is computed from an orthographic string by a set of rules (Coltheart, 1978; Forster & Chambers, 1973; Marshall and Newcombe, 1973). Rastle and Coltheart (2000) suggested that stress information could be stored in the lexical route of the model as a property of item-specific phonological representations, and thus retrieved during the reading aloud of known words. They concentrated instead on the more challenging task of implementing a stress assignment procedure along the sublexical route of

The rule-based process developed by Rastle and Coltheart (2000) was designed to execute both the mapping between sublexical orthographic and phonological representations (segmental information) and the assignment of stress along with the appropriate vowel reduction (suprasegmental information). Morphological structure plays an important role in the system of rules that Rastle and Coltheart (2000) implemented, particularly in relation to the assignment of stress (for an illustration of the stress rules refer to Figure 2, p. 349 in Rastle & Coltheart, 2000). Specifically, the identification of a prefix (e.g., pre-, de-, dis-, re-, mis-) results in the assignment of second-syllable stress, while the identification of a suffix results in the assignment of first-syllable stress (except in the case of a small group of stress-taking suffixes identified by Fudge (1984) such as -een, -ique, -oo). In the absence of an identifiable affix, first-syllable stress is assigned, which is the dominant stress pattern for disyllables in the English language. Rastle and Coltheart (2000) reported that the algorithm successfully predicted stress assignment on 89.7% of all disyllabic English words present in the CELEX database (Baayen, Piepenbrock, & van Rijn, 1993), and it also predicted the modal stress given to 84% of a large set of disyllabic nonwords read aloud by human subjects. This work thus provides evidence supporting the notion that prefixes can serve as important cues for stress assignment, and more generally, that sublexical cues for assigning stress to disyllables can be expressed within a system of rules relating spelling to sound.

The present study introduces a new approach to ascertaining the sublexical cues to stress assignment. Specifically, we denote prefixed words as ‘regular’ if they take second-syllable stress (e.g. remind) and ‘irregular’ if they take first-syllable stress (e.g. reflex). We then test whether patients with acquired surface dyslexia, an acquired disorder of reading in which the reading aloud of irregular words is impaired while the reading aloud of nonwords

pronunciation when reading aloud irregular monosyllabic words (e.g., reading *pint* as if it rhymed with *mint*). Thus, while these patients demonstrate an impairment in utilising lexical information during reading, their ability to translate orthography to phonology via sublexical operations appears to be intact. Accordingly, we hypothesized that these patients would commit stress regularisation errors when reading aloud irregularly-stressed disyllabic words (e.g., read 'reflex' with second-syllable stress).

While patients with surface dyslexia have typically been examined in respect of the segmental errors produced during reading aloud, it has long been known that they also produce errors with respect to suprasegmental information (Marshall & Newcombe, 1973). Stress regularization errors in acquired, as well as developmental, surface dyslexia have been observed in different languages, including English (Marshall & Newcombe, 1973), Italian (e.g., Galante, Trali, Zuffi, & Avanzi, 2000; Laganaro, Vacheresse, & Frauenfelder, 2002; Miceli & Caramazza, 1993; Paizi, Zoccolotti, & Burani, 2011; Trenta, Benassi, Di Filippo, Pontillo, Zoccolotti, 2013; Zoccolotti, De Luca, Di Pace, Judica, Orlandi, & Spinelli, 1999), German (Janssen, 2003), Filipino (Dulay & Hanley, 2015), and Hebrew (Lukov & Friedmann, 2008). In the majority of these studies, stress errors involved overgeneralizations of the most frequent stress pattern of the given language. Surprisingly, however, there has been hardly any work investigating the sublexical knowledge used in stress assignment by patients with surface dyslexia. The only study that examined this issue was carried out by Jansen (2003) in German. In particular, Jansen (2003) investigated stress error patterns in two patients with surface dyslexia in order to determine whether syllable structure is an important predictor of stress assignment in the German reading system. Inspection of the patients' regularization errors revealed that in German, an open final syllable (i.e., ending with a

Accordingly, in the present study, we sought to uncover the sublexical cues to stress assignment in the English language by investigating whether patients with acquired surface dyslexia are likely to assign second-syllable stress to prefixed words that are irregularly stressed (e.g. reflex). More specifically, we tested the hypothesis put forward by Rastle and Coltheart (2000) that prefixes repel stress, so that identification of a prefix in a two-syllable letter string leads readers to assign second syllable stress. Patients were given three types of disyllabic words for reading aloud. Two of these types were prefixed words that varied in regularity according to the prefixes-repel-stress rule. The regular prefixed words contained a weak-strong syllable stress pattern and required a second syllable stress (e.g., remind), whereas the irregular prefixed words contained a strong-weak syllable stress pattern and required a first-syllable stress (e.g., reflex). Stress assignment to regular and irregular prefixed words was examined against a third type of control word. These were non-prefixed words that contained a strong-weak stress pattern (e.g., scandal), which is the dominant stress pattern of English disyllabic words in the absence of prefixation. If prefixes repel stress within the sublexical procedure for reading aloud, then we would expect surface dyslexic patients to assign second syllable stress to prefixed words, thus yielding stress errors in the case of 'irregular' prefixed words with strong-weak stress.

In addition to reporting data from five surface dyslexic patients on the reading aloud of disyllabic words, we report simulations from the disyllabic algorithm developed by Rastle and Coltheart (2000) and from the CDP++ model developed by Perry, Ziegler, and Zorzi (2010), which are currently the only two publicly available computational implementations of reading aloud that provide both a stress marker and a phonological code for English

Coltheart (2000) reflects only the sublexical reading aloud process; the output of this process thus reflects reading in the absence of lexical information (i.e. pure surface dyslexia). Given the nature of the hard-coded rules in the implemented algorithm, hereafter referred to as RC00, our prediction was that the algorithm would assign second-syllable stress to prefixed words and thus, commit stress errors on 'irregular' prefixed words with strong-weak stress (e.g., saying reflex as /r@fIEks'/). We opted to contrast these simulations with those from the CDP++ model (Perry et al., 2010). CDP++ is a computational implementation of the dual-route theory much like the DRC model (Coltheart et al., 2001), comprising a lexical and a sublexical procedure. However, its sublexical procedure consists of a two-layer associative (TLA) network for mapping graphemes onto phonemes, as opposed to a set of rules. In this model, stress assignment to disyllabic words is coded in a stress buffer that is connected with both the lexical and the sublexical procedures. While the lexical procedure of the model directly activates the stress that is associated with a familiar word's spoken form, the sublexical procedure activates the stress that it learnt to associate with the graphemes of a word. These grapheme to stress associations are learnt during a training phase in the same way as grapheme to phoneme mappings are formed, that is, via a connectionist algorithm based on the statistical distributional regularities of the spelling-to-sound and spelling-to-stress mappings. Surface dyslexia can be simulated in the CDP++ model by lesioning connections within the lexical procedure to varying degrees. The CDP++ model provides an interesting contrast to the RC00 algorithm because prefixes are not explicitly represented in its sublexical pathway. Hence, this model is not expected to be sensitive to associations between prefixes and certain stress patterns. For this reason, we predicted that the CDP++

The comparison between the RC00 algorithm and the CDP++ model also bears additional theoretical value in respect of how sublexical orthographic knowledge is used to generate phonology at the suprasegmental level during reading in general. Indeed, the comparison of the stress error patterns produced by these models may provide a platform to evaluate the extent to which this generative knowledge is expressed as a set of explicit rules, or as an implicit system of statistical regularities that has been acquired through learning.

## **2. Case Reports**

We recruited five cases of acquired surface dyslexia, four female and one male. All were judged to be candidates for surface dyslexia on the basis of (a) impaired irregular word relative to regular word reading and (b) relatively preserved nonword reading. These criteria were assessed through administration of the PALPA 35 and 36 (Psycholinguistic Assessment of Language Processing in Aphasia; Kay, Lesser, & Coltheart, 1992) and the Coltheart and Leahy (1996) reading tests. These assessments were conducted as part of a comprehensive neuropsychological evaluation performed on each individual case. In particular, each case was screened for dementia with the Addenbrooke's Cognitive Examination Revised (ACE-R; Mioshi, Dawson, Arnold, & Hodges, 2006) test battery, while additional tests assessing the different memory and language components were also administered. The administered assessments and the neuropsychological profile of each case are summarized in Table 1. All cases had normal or corrected to normal vision. Below we provide a brief medical history report for each of the five cases, including a short discussion of the patients' cognitive impairments and reading aloud performance at the time in which the present study was conducted.

were subsequently resolved, and no current evidence of a general mood disorder. There was some family history of dementia, with her mother having had vascular dementia. Neurological investigation included a CT undertaken in April 2012, which reported *“asymmetry of the temporal horns of the lateral ventricles with some prominence of the sulci of the temporal lobe on the left. This is in keeping with a degree of left temporal lobe atrophy. No other significant abnormality.”* A subsequent MRI undertaken in August 2012 reported *“significant atrophy of the left temporal lobe and to a lesser extent the right. No mass lesion identified”*, consistent with primary progressive aphasia. The patient undertook a degree in natural sciences and after graduation worked as a teacher, married, and raised a family.

Upon presentation, general cognitive testing indicated an impaired profile (ACE-R score 50/100) somewhat complicated by clear expressive and receptive aphasia – although some episodic memory impairment was present. Expressive aphasic presentation was evident on the basis of verbal fluency impairment and severe naming problems, though word repetition testing and auditory verbal short-term memory were normal (see Table 1). Severe semantic memory impairment was also present as assessed by the Pyramids and Palm Trees test (PPT, Howard & Patterson, 1992), where she obtained a score of 36 out of 52. The presence of semantic errors in naming and impaired comprehension, along with anterior temporal lobe atrophy (accentuated in the left hemisphere) are considered consistent with semantic dementia (Gorno-Tempini et al., 2011). Critically, Patient’s 1 reading assessment (PALPA 35, 36, and Coltheart & Leahy tests combined) exhibited the distinctive pattern of surface dyslexia, characterized by impaired irregular word reading (40% correct) in the presence of relatively spared regular word and nonword reading (96.7% and 94.6% correct, respectively). Notably, the vast majority of reading errors on irregular words involved

**Patient 2.** This patient is a right-handed female, aged 63 at time of testing. She had a four-year history of memory and language problems, with some consequential anxiety but no other evidence of generalized mood disorder. Osteo-arthritis in the knees was her only other physical complaint. There was also a history of dementia in the family, with her mother having developed Alzheimer's Disease (AD). Neurological investigation included a CT undertaken in March 2012, which reported "*mild features of general atrophy with widening of the Sylvian fissure and associate bilateral temporal lobe convexity. No evidence of any ischaemic change or mass lesion identified*", with a diagnosis of fronto-temporal dementia with primary progressive aphasia of the non-fluent subtype. The patient had been a homemaker throughout her life, leaving school at 16 to work for a brief period in retail, and is bilingual in Welsh and English.

Upon presentation, general cognitive testing indicated a dementia cognitive profile (ACE-R score 25/100) with episodic memory impairment and constructional apraxia as specified by her performance in the Rey-Osterrieth Complex Figure test (ROCF; Osterrieth, 1944). Expressive and receptive aphasic presentation was evident. Speech production was non-fluent at presentation, with severe auditory verbal short memory impairment, and word repetition, fluency and word naming problems (see Table 1). Mild semantic memory impairment was also present (PPT score 46/52). Patient's 2 reading assessment (PALPA 35, 36, and Coltheart & Leahy tests combined) revealed a surface dyslexia profile. Specifically, Patient's 2 regular word and nonword reading was highly accurate (83.3% and 85.4% correct, respectively), whereas irregular word reading was considerably impaired (48.3% correct). Errors to irregular words included 64.3% of spelling-to-sound regularizations, indicative of a

**Patient 3.** This patient is a right-handed male, aged 61 at time of testing. He had a three-year history of memory problems with some history of anxiety (treated with medication), but no generalized mood disorder and no family history of dementia. Neurological investigation included a CT undertaken in March 2012, which reported “*mild widening of the subarachnoid spaces in relation to the temporal lobes with widening of the Sylvian fissures and sulci. No evidence of focal ischaemic change*”, with a diagnosis of Alzheimer’s Disease. The patient had left school at 18 and worked as an insulation engineer throughout his professional life, marrying, and having children.

Upon presentation, general cognitive testing indicated a dementia cognitive profile (ACE-R score 37/100) with episodic memory impairment and constructional apraxia as specified by the patient’s ROCF test performance. No expressive aphasic presentation was evident on the basis of only mild naming problems and normal word repetition testing, though verbal fluency was severely impaired, as was auditory verbal short-term memory (see Table 1); semantic memory impairment was also present (PPT score 37/52). Patient’s 3 reading assessment (PALPA 35, 36, and Coltheart & Leahy tests combined) revealed a surface dyslexia profile. Specifically, while Patient’s 3 regular word reading was virtually perfect (96.7% accurate) and his nonword reading highly accurate (83.3% correct), this patient’s irregular word reading was relatively impaired (73.3% correct). Patient 3 regularized the spelling-to-sound pronunciation of 47.3% of all irregular words, reflecting a reliance on the sublexical reading procedure. Examples of such regularizations included /tQm / for the word ‘tomb’, /kw1/ for the word ‘quay’, and /su/ for the word ‘sew’.

family history of dementia. Neurological investigation included a CT undertaken in June 2013, which reported “*a disproportionate level of atrophy affecting the frontal lobes bilaterally. Additionally marked prominence of widening of the Sylvian fissure associated with temporal lobe atrophy. No evidence of focal ischaemic change or mass lesion identified*”, with a diagnosis of dementia of the Alzheimer’s type. The patient left school at 16 to work in domestic employment at a children’s home and raise a family.

Upon presentation, general cognitive testing indicated a dementia cognitive profile (ACE-R score 32/100) with episodic memory impairment and constructional apraxia as specified by the patient’s ROCF test performance. No expressive aphasic presentation was evident on the basis of naming and word repetition testing, though verbal fluency was severely impaired, as was auditory verbal short-term memory (see Table 1); mild semantic memory impairment was also present (PPT score 44/52). Patient’s 4 reading assessment (PALPA 35, 36, and Coltheart & Leahy tests combined) revealed a surface dyslexia profile. Specifically, while Patient’s 4 regular word reading was virtually perfect (95% accurate) and nonword reading highly accurate (90.4% correct), this patient’s irregular word reading was relatively impaired (78.3% correct). Patient 4 regularized the spelling-to-sound pronunciation of 53.6% of all irregular words, reflecting a reliance on the sublexical reading procedure. Examples of such regularizations included /rut2n/ for the word ‘routine’, /kw1/ for the word ‘quay’, and /Izl}nd/ for the word ‘island’.

**Patient 5.** This patient is a right-handed female, aged 69 at time of testing. She had a three-year history of memory problems, but no evidence of any mood disorder and no family history of dementia. She also receives medication for hypothyroidism. Neurological investigation included a CT conducted in December 2013, which reported “*mild generalized*

patient left school at 16 to work as a shop assistant and subsequently as a warehouse manager, marrying, and having children.

Upon presentation, general cognitive testing indicated a dementia cognitive profile (ACE-R score 62/100) with episodic memory impairment. Some expressive aphasic presentation was evident on the basis of moderate naming problems and impaired verbal fluency - but normal word repetition testing and mild impairments of auditory verbal short-term memory (see Table 1); mild semantic memory impairment was also present (PPT score 46/52). Patient's 5 reading assessment (PALPA 35, 36, and Coltheart & Leahy tests combined) revealed a surface dyslexia profile. Specifically, while Patient's 4 regular word reading was virtually perfect (96.7% accurate) and nonword reading highly accurate (90% correct), this patient's irregular word reading was relatively impaired (76.7% correct). Notably, Patient's 5 errors on irregular words included a striking 90.4% of spelling-to-sound regularizations. The nature of these errors reflect the patient's reliance on the sublexical reading procedure. Examples of such regularizations included /g9g/ for the word 'gauge', /IndIkt/ for the word 'indict, and /'k5l@nEl/ for the word 'colonel'.

[Table 1 about here]

### **3. Method**

#### **3.1 Stimuli**

One hundred and fifty disyllabic English words were selected for inclusion in three conditions. The 'Prefix W-S' condition comprised 50 words that had a prefix and were

according to the 'prefixes-repel-stress' rule. The 'Prefix S-W' condition comprised 50 words that also had a prefix but were characterised by a strong-weak syllable stress pattern; that is, their correct pronunciation required first syllable stress (e.g., reflex, subway). These items were considered 'irregular' according to the 'prefixes-repel-stress' rule. Finally, the 'NoPrefix S-W' condition comprised 50 words that had no prefix and were characterised by a strong-weak syllable stress pattern, which is the dominant stress pattern of English disyllabic words in the absence of prefixation. The correct pronunciation of these control words required first syllable stress (e.g., scandal, volume). Note that prefixation was defined on a purely orthographic basis (see Rastle & Davis, 2008 for relevant discussion). A word was considered as prefixed if it began with a letter sequence that could form an identifiable English prefix, and was followed by either an existing stem (e.g., remove) or by a potential stem (e.g., discreet). The two prefixed conditions (Prefix W-S and Prefix S-W) consisted of the same set of prefixes, which appeared equally frequently in each of these conditions and were predominantly associated with second syllable stress in disyllabic English words. The presence of an embedded word was also controlled, so that half of the items in each of the three conditions contained an embedded 'stem' (e.g., *remove*, *input*, *climate*, respectively), whereas the other half did not (e.g., *discreet*, *prospect*, *minute*, respectively).

Words in the three conditions were group-wise matched as closely as possible on CELEX word frequency (Baayen, Piepenbrock, & van Rijn, 1995), word length in letters, orthographic neighbourhood size (Coltheart's N), bigram type frequency (values extracted from the N-Watch database, Davis, 2005) and age of acquisition (AoA, Kuperman, Stadthagen-Gonzalez, & Brysbaert, 2012). The mean values of these variables across the experimental conditions are reported in Table 2. Stimuli and patient responses are listed in

### 3.2 Procedure

All five patients were presented with the 150 stimuli in a random order and were asked to read them aloud within a single testing session.

## 4. Results

### 4.1 Patient Data

For each patient, the percentage of stimuli given second-syllable stress was calculated in each condition (see Table 3). This percentage was based on all intelligible disyllabic responses. Critically, responses that comprised second-syllable stress in the Prefix S-W and NoPrefix S-W conditions were stress errors (e.g., /Ins2t/ for insight; /bISQp/ for bishop, respectively). Further examination of these stress errors sought to assess whether they could be deemed *pure* stress errors. Pure stress errors were defined as responses that contained an erroneous stress assignment but the correct phonemic pronunciation of the word. Pronunciations that were slightly altered as a result of the erroneous stress assignment (e.g., reducing to schwa the initial vowel in ‘agate’), or that involved spelling-to-sound regularizations (e.g., /dIpQt/ for the word ‘depot’), were also classified as pure stress errors. Non-pure stress errors involved incorrect pronunciations in addition to stress displacement (e.g., /pVt5t/ for the word ‘input’). As shown in Table 3, the vast majority of incorrect

Patients' responses are reported in Appendix B. It is noteworthy that beyond stress assignment errors, all patients produced a number of spelling-to-sound regularization errors, which are characteristic of surface dyslexia. In order to quantify the incidence of spelling-to-sound regularisations, we identified words in our stimulus set that both RC00 and the sub-lexical pathway of the CDP++ model produced incorrectly. There were 20 such items (or, 100 opportunities to observe spelling-to-sound regularisations). Patients produced regularisations in 40% of these cases while an additional 6% of the responses were either missing or otherwise incorrect. The words that were systematically regularized by the majority of patients (3 and above) were the following: depot (/dIpQt'/), climate (/ˈklɪm1t/), diamond (/ˈdɪmQnd/), minute (/mɪnʉt'/), surface (/ˈsʒf1s/), message (/ˈmEs1\_/), fountain (/ˈfʉnt1n/), mountain (/ˈmʉnt1n/), and input (/ˈɪnpVt/). These regularisations in pronunciation provide further evidence of the patients' surface dyslexia.

[Table 3 about here]

Logistic regression analyses were carried out for each patient individually. These analyses investigated the probability of second-syllable stress occurring (a binary variable) as a function of condition (3 levels) and the presence of an embedded word (another binary variable). Although our stimuli were closely matched on the psycholinguistic characteristics presented in Table 2, we included these continuous variables as covariates in the analysis of each patient to ensure that any effects of our factors of interest were not driven by small differences in these variables across the three conditions. However, for ease of exposition, we

hypothesis, if prefixes repeat stress, we would expect patients to depart from the typical strong-weak stress pattern of English disyllables and assign second-syllable stress to prefixed words. Specifically, relative to the control NoPrefix S-W words, we hypothesized that patients would be more likely to assign second-syllable stress to both regular Prefix W-S words and irregular Prefix S-W words. Second-syllable stress assignment to the latter words would denote stress regularisation errors.

Results revealed a significant influence of condition on the production of second-syllable stress for all patients. The Wald test statistics for the main effect of condition and the follow-up comparisons for each patient are reported in Table 4. All patients were more likely to assign second-syllable stress to Prefix W-S words, relative to NoPrefix S-W words, with an odds ratio ranging from 139.5 (for Patient 5) to 774.0 (for Patient 4). More importantly, all patients were more likely to incorrectly assign second-syllable stress to Prefix S-W words compared with NoPrefix S-W words, with an odds ratio ranging from 16.8 (for Patient 3) to 70.2 (for Patient 2). None of the patients revealed an effect of embedded words on the probability of assigning second-syllable stress. The Wald test statistics for the main effect of embedded word for each patient are also reported in Table 4. The covariates did not reveal significant effects on second-syllable stress assignment patients 2, 4, and 5 (all  $p$  values  $>.05$ ). For Patient 1, the effect of orthographic length was significant ( $W(1) = 10.61$ ,  $p = .001$ ), indicating that longer words were 3.2 times more likely to receive second-syllable stress than shorter words.<sup>2</sup> For Patient 3, the effects of age of acquisition and orthographic length were significant ( $W(1) = 4.32$ ,  $p = .038$ , and  $W(1) = 4.32$ ,  $p = .038$ , respectively), indicating that words that were acquired later in life were 1.3 times more likely to receive

[Table 4 about here]

## 4.2 Simulations

Simulations were run using the RC00 algorithm (Rastle & Coltheart, 2000) and the executable version of the CDP++ model (Perry et al., 2010). None of the patients showed totally impaired irregular word reading in the presence of totally unimpaired nonword and regular word reading. However, our simulations represent the case of pure surface dyslexia. While we acknowledge that this is a simplification, the RC00 algorithm only expresses a set of hypotheses about the sublexical rules relating spelling to sound and spelling to stress for disyllabic letter strings. Hence, this model can only simulate pure surface dyslexia. In order to simulate pure surface dyslexia with the CDP++ model, the lexical route was deactivated. In what follows, we describe the sublexical rules used by the RC00 algorithm and the sublexical procedure of the CDP++ model for the pronunciation and stress marking of English disyllabic words.

**RC00 algorithm.** The RC00 algorithm (Rastle & Coltheart, 2000) is an implementation of a rule-based sublexical pathway that translates printed disyllables to sound and applies a stress marker. The RC00 algorithm calls on the grapheme-to-phoneme translation rules used by the DRC model and in addition, it identifies orthographic strings corresponding to prefixes and suffixes to determine stress placement. The algorithm begins by searching a letter string for the presence of an existing prefix in the English language (e.g., pre-, de-, dis-, re-, mis-). A prefix is identified on a purely orthographic basis and only if it is

annex store; the remaining part of the word is translated via the grapheme-to-phoneme rules used by the DRC model (Coltheart et al., 2001). The pronunciation of the word is then assembled and the prefix is given non-stress and reduced vowel schwa if appropriate. In the absence of a prefix, the algorithm uses a similar procedure to search for the presence of a suffix at the end of the string. The presence of certain suffixes (i.e. the suffix -y or a suffix that begins with -e) lengthens the first phonological vowel of the string, and the suffix is given non-stress unless it belongs to a group of stress-taking suffixes identified by Fudge (1984; e.g., -een, -ique, -oo), in which case it is given stress. All non-suffixed letter strings are translated into a string of phonemes using the DRC grapheme-to-phoneme rules and are given first-syllable stress.

**CDP++ model.** The CDP++ model is a dual-pathway model of reading aloud comprising lexical and sublexical processes for mapping print-to-sound. The CDP++ model is a full processing model that produces a pronunciation, stress marker, and reaction time. It is built on its direct precursor, the CDP+ model (Perry et al., 2007), a successful model of reading aloud monosyllabic words. The CDP++ model is very similar to the CDP+ model except that it includes more letter and phoneme slots to accommodate longer words, a different input coding scheme to accommodate disyllables, it introduces the schwa phoneme to deal with vowel reduction and stress nodes to represent the position of stress, and it uses a larger training corpus and lexicon. As in the CDP+ model, the core component of the sublexical procedure of the CDP++ model is the TLA network of phonological assembly. The TLA network contains grapheme nodes that are linked to phoneme nodes via weighted connections that represent the most reliable mappings between orthography and phonology. These mappings are learnt via a connectionist algorithm during training, where the model is

sublexical stress nodes are fully connected to the sublexical grapheme nodes. During training, the model learns grapheme to stress relationships directly, in the same way and under the same parameters as it learns grapheme to phoneme relationships. Activation from the sublexical stress nodes is sent to two stress output nodes that are placed at the level of the phonological output buffer via an excitation parameter. In order to avoid a first-syllable stress bias in the phoneme output buffer, sublexical stress activation begins to activate the stress output nodes only after the last letter of the word is processed by the model's graphemic parser. Finally, a lateral inhibition parameter at the stress output level, allows activation from one stress output node to laterally inhibit the other. The stress output nodes also pool information from the lexical procedure of the model, where lexically defined stress information is sent via excitation and inhibition parameters from the phonological lexicon to the phonological output buffer. In the present simulation of pure surface dyslexia, the lexical procedure of the CDP++ model was completely switched off, so that it would not contribute at all to word reading aloud.

### **4.3 Simulation Data**

Table 5 below reports the percentages of second-syllable stress assigned by the RC00 algorithm and the CDP++ model across all conditions. As with the patient data above, stress assignments were calculated only for disyllabic responses. The responses produced by the algorithm and the CDP++ model are reported in Appendix B, where the excluded responses are also indicated. Logistic regression analyses were carried out separately for the RC00 algorithm and the CDP++ model, in the same way they were performed for the patient data.

**RC00 algorithm.** The simulation results revealed a significant influence of condition on the assignment of second-syllable stress ( $W(2) = 43.43, p < .001$ ). The RC00 algorithm was 160.9 times more likely to assign second-syllable stress to regular Prefix W-S words than NoPrefix S-W words ( $W(1) = 35.14, p < .001$ ). More importantly, the RC00 was 153.8 times more likely to stress the second syllable of the irregular Prefix S-W words, compared with the NoPrefix S-W words ( $W(1) = 30.83, p < .001$ ). There was no effect of embedded words on the probability of assigning second syllable stress ( $W(1) = .30, p = .584$ ). None of the covariates revealed any significant effects on second-syllable stress assignment (all  $p$  values  $>.05$ ).

**CDP++ model.** The simulation results revealed a significant influence of condition on the assignment of second-syllable stress ( $W(2) = 28.19, p < .001$ ). The CDP++ model was 182.8 times more likely to assign second-syllable stress to regular Prefix W-S words than NoPrefix S-W words ( $W(1) = 22.26, p < .001$ ). Importantly, the CDP++ model was 27.6 times more likely to stress the second syllable of the irregular Prefix S-W words, compared with the NoPrefix S-W words ( $W(1) = 9.19, p = .002$ ). There was no effect of embedded words on the probability of assigning second syllable stress ( $W(1) = .68, p = .410$ ). The effect of age of acquisition was nearly significant ( $W(1) = 3.83, p = .05$ ), indicating that words with a higher age of acquisition were 1.3 times more likely to be assigned second-syllable stress. All other covariates revealed no significant effects on second-syllable stress assignment (all  $p$  values  $>.05$ ).

phonology has focused on the segmental level – that is, the relationship between graphemes and phonemes (e.g., Andrews & Scarratt, 1998; Pritchard, Coltheart, Palethorpe, & Castles, 2012). The present study is concerned with the relationship between orthography and phonology at the suprasegmental level – specifically, the assignment of appropriate stress patterns to letter strings with more than one syllable. Multiple sublexical cues to stress have been put forward (e.g., vowel length, orthographic weight, morphological structure); in this work, we focused on the role of morphological structure on stress assignment. In particular, we tested the hypothesis that prefixes repel stress (Rastle & Coltheart, 2000) by examining whether patients with acquired surface dyslexia are likely to assign second-syllable stress to prefixed words with an irregular stress pattern (e.g. reflex). We reasoned that if prefixes repel stress along the sublexical reading process, these patients should be more likely to incorrectly assign second-syllable stress to these items, leading them to commit stress regularisation errors.

Results confirmed our prediction. All patients assigned significantly more second syllable stress to prefixed words relative to non-prefixed words with a strong-weak stress pattern (e.g., scandal). All patients correctly assigned second-syllable stress to the vast majority of regular weak-strong prefixed words. Critically, all patients committed regularization errors by incorrectly assigning second-syllable stress to irregular prefixed words with a strong-weak stress pattern (e.g., reflex) compared with non-prefixed words that shared the same stress pattern (e.g. scandal). This result was observed consistently across all five patients and interestingly, the likelihood of producing stress regularization errors varied across patients, reflecting their varying levels of surface dyslexia. For example, Patients 1 and

Coltheart & Leamy, 1996; see Table 1), these patients seem to rely almost exclusively on their sublexical procedures for the orthography to phonology mapping.

It is important to note that besides the patients' regularization errors on irregular strong-weak prefixed words, we also observed some stress errors in the other two types of words. In particular, patients incorrectly assigned second-syllable stress to some non-prefixed strong-weak words (Patients 1 and 2 especially), and their performance to regular prefixed weak-strong words was not perfect, as some of these words were assigned first-syllable stress. Thus, it could be argued that the patients' stress errors may have resulted from a cognitive impairment other than or additional to surface dyslexia, such as a deficit in the phonological output buffer (Patient 2 exhibits very poor word and nonword repetition, for example). However, if that were the case we would not expect to observe the systematic pattern of stress regularizations for irregular strong-weak prefixed items that all patients clearly demonstrated. If anything, we would expect more stress errors for prefixed words with a weak-strong stress pattern, which is by far more infrequent in the English language than the strong-weak stress pattern in English disyllabic words. Instead, the vast majority of the prefixed words with a weak-strong stress pattern were correctly assigned second-syllable stress by all patients. The great preponderance of stress errors on the irregular prefixed words, systematically shown by all patients, provides support for the idea that the presence of a prefix prompted patients to assign second-syllable stress, regularizing thus, stress assignment to these items. Overall, these findings confirm the hypothesis that prefixes repel stress in reading aloud disyllabic words and, together with Rastle and Coltheart's (2000) work, establish the functional value of prefixes as sublexical cues to stress assignment in reading.

producing a considerable number of stress errors for irregular strong-weak prefixed words.

This pattern was entirely expected, of course, since the RC00 algorithm includes a rule that assigns second syllable stress to disyllables with prefixes. One limitation of this algorithm is that it is not a fully implemented model (i.e. it is not integrated with the lexical route of the DRC model; Coltheart et al., 2001). For this reason, it is only able to simulate pure surface dyslexia – the total impairment of the lexical procedure in the presence of full functioning of the sublexical procedure. Thus, while this simulation has been able to approximate the number of regularization errors made by the most impaired patients, Patients 1 and 2, the model did not capture accurately the performance in of patients with milder cases of surface dyslexia could also be captured. We are also mindful that the most impaired patients additionally made a substantial number of stress errors on non-prefixed words. This was not the case for the RC00 algorithm. These errors likely reflect additional sublexical cues to stress that are not represented in the RC00 algorithm.

Simulation results from the CDP++ model (Perry et al., 2010) were rather surprising. In a simulation of pure surface dyslexia (with the lexical route of the model completely switched off), this model also produced a large number of stress errors for the irregular strong-weak prefixed words (50%) compared with the non-prefixed strong-weak words. This performance approximated the performance of Patients 3 and 4. These findings are rather surprising because unlike the RC00 algorithm, prefixes are not explicitly represented in the CDP++ model. The sublexical procedure of the CDP++ model consists of a two-layer associative (TLA) network, which maps graphemes to phonemes and graphemes to stress nodes. In this model, grapheme-to-phoneme and grapheme-to-stress relations are implicitly learned through their statistical distribution in the lexicon, rather than being represented as a

second-syllable stress. The CDP++ model appeared to do more poorly on correctly assigning second-syllable stress to regular weak-strong prefixed words than did most of the patients, but again, approximated the results from Patient 3 on this dimension (81.2%).

While our simulations with the CDP++ model represented the case of pure surface dyslexia, it is important to note that this model is well suited to simulating varying levels of severity in surface dyslexia. This is achieved by gradually lesioning connections within the lexical procedure to varying degrees, as opposed to turning this procedure off entirely. Because our patients clearly exhibit surface dyslexia of varying levels of severity, our original intention was to conduct these more graded simulations. However, the results from the current simulation, which assumed no contribution from the lexical route, removed the rationale for conducting this additional modeling work. Even with the lexical route completely turned off, the model yielded a percentage of stress errors in the irregular S-W prefix condition that was lower than four of the five patients. It is difficult to conceive how more subtle lesions within the lexical pathway could increase the percentage of stress errors in this condition to simulate the more severe cases of surface dyslexia (e.g. Patients 1 and 2). The CDP++ model also displays some problems in the actual responses that it produces, which are unlike those of the patients in various ways. For example, in contrast to the patients, the CDP++ model produced monosyllabic responses for a number of items (e.g., injure /In\_'/; pigeon /'p2\_'/; borough /'b9r/). It also produced impossible responses in which the schwa vowel was stressed (e.g., return /rEt@n'/), or both vowels were reduced to schwa (e.g., conclude /k@@dd'/). The fact that the model produces responses that are atypical of human responses has already been highlighted in the monosyllabic domain (Pritchard et al.,

In conclusion, we have presented a new approach to investigating sublexical cues to stress assignment in reading aloud. By investigating stress regularizations produced by five patients with surface dyslexia we have shown that the morphological structure of a stimulus (in particular, prefixation) provides important information as to its likely stress pattern. Further, while the simulations do not conclusively favour one theoretical approach to reading aloud over the other, they do highlight challenges for each of the models as they are developed further.

- Hough dou peapel rede gnew wirds. *Journal of Experimental Psychology: Human Perception & Performance*, 24, 1052–1088. doi:10.1037//0096-1523.24.4.1052
- Ans, B., Carbonnel, S., & Valdois, S. (1998). A connectionist multiple-trace memory model for polysyllabic word reading. *Psychological Review*, 105, 678–723. doi:10.1037/0033-295X.105.4.678-723
- Arciuli, J., & Cupples, L. (2006). The processing of lexical stress during visual word recognition: Typicality effects and orthographic correlates. *Quarterly Journal of Experimental Psychology*, 59, 920–948. doi:10.1080/02724980443000782
- Arciuli, J., & Cupples, L. (2007). Would you rather ‘embert a cudsert’ or ‘cudsert an embert’? How spelling patterns at the beginning of English disyllables can cue grammatical category. In A. Schalley & D. Khlentzos (Eds.), *Mental states: Language and cognitive structure* (pp. 213–237). Amsterdam, the Netherlands: John Benjamins.
- Arciuli, J., Monaghan, P., & Ševa, N. (2010). Learning to assign lexical stress during reading aloud: Corpus, behavioural, and computational investigations. *Journal of Memory and Language*, 63, 180–196. doi: 10.1016/j.jml.2010.03.005
- Baayen, R. H., Piepenbrock, R., & Gulikers, L. (1995). *The CELEX lexical database. Release 2 (CD-ROM)*. Philadelphia: Linguistic Data Consortium, University of Pennsylvania.
- Baayen, R. H., Piepenbrock, R., & van Rijn, H. (1993). *The CELEX lexical database (CD-ROM)*. Linguistic Data Consortium, University of Pennsylvania, Philadelphia, PA.
- Baker, R., & Smith, P. (1976). A psycholinguistic study of English stress assignment rules. *Language and Speech*, 19, 9–27.

Burani, C., Paizi, D., & Sulpizio, S. (2014). Stress assignment in reading Italian: Friendship outweighs dominance. *Memory & Cognition*, 42, 662–675. doi:10.3758/s13421-013-0379-5

Cappa, S. F., Nespola, M., Ielasi, W., & Miozzo, A. (1997). The representation of stress: Evidence from an aphasic patient. *Cognition*, 65, 1–13. doi:10.1016/S0010-0277(97)00024-3

Colombo, L. (1992). Lexical stress effect and its interaction with frequency in word pronunciation. *Journal of Experimental Psychology: Human Perception and Performance*, 18, 987–1003. doi:10.1037/0096-1523.18.4.987

Coltheart, M. (1978). Lexical access in simple reading tasks. In G. Underwood (Ed.), *Strategies of information processing* (pp. 151–216). London: Academic Press.

Coltheart, M., Curtis, B., Atkins, P., & Haller, M. (1993). Models of reading aloud: Dual-route and parallel-distributed-processing approaches. *Psychological Review*, 100, 589–608. doi:10.1037/0033-295X.100.4.589

Coltheart, M., & Leahy, J. (1996). Assessment of lexical and nonlexical reading abilities in children: Some normative data. *Australian Journal of Psychology*, 48, 136–140. doi:10.1080/00049539608259520

Coltheart, M., & Rastle, K. (1994). Serial processing in reading aloud: Evidence for dual-route models of reading. *Journal of Experimental Psychology: Human Perception and Performance*, 20, 1197–1211. doi:10.1037/0096-1523.20.6.1197

- Davis, C. J. (2005). N-Watch: A program for deriving neighborhood size and other psycholinguistic statistics. *Behavior Research Methods, 37*, 65–70.  
doi:10.3758/bf03206399
- Dulay, K. M. & Hanley, J.R. (2015). Stress errors in a case of developmental surface dyslexia in Filipino. *Cognitive Neuropsychology, 32*, 29–37.  
doi:10.1080/02643294.2014.984602
- Forster, K. I. & Chambers, S. M. (1973). Lexical access and naming time. *Journal of Verbal Learning and Verbal Behavior, 12*, 627–635. doi:10.1016/S0022-5371(73)80042-8
- Friedmann, N., & Lukov, L. (2008). Developmental surface dyslexias. *Cortex, 44*, 1146-1160. doi: 10.1016/j.cortex.2007.09.005
- Fudge, E. C. (1984). *English word stress*. London: Allen & Unwin.
- Galante, E., Tralli, A., Zuffi, M., & Avanzi, S. (2000). Primary progressive aphasia: a patient with stress assignment impairment in reading aloud. *Neurological Sciences, 21*, 39–48. doi:10.1007/s100720070117
- Gorno-Tempini, M. L., Hillis A. E., Weintraub, S., Kertesz, A., Mendez, M., Cappa, S. F., Ogar, J. M., Rohrer, J. D., Black, S., Boeve, B. F., Manes, F., Dronkers, N., F., Vandenberghe, R., Rascovsky, K., Patterson, K., Miller, B. L., Knopman, D. S., Hodges, J. R., Mesulam, M. M., & Grossman, M. (2011). Classification of primary progressive aphasia and its variants. *Neurology, 76*, 1006–1014.  
doi:10.1212/WNL.0b013e31821103e6

- Howard, D., & Patterson, K. (1992). *Pyramids and Palm Trees: A test of semantic access from pictures and words*. Bury St Edmunds, UK: Thames Valley Test Company.
- Janssen, U. (2003). Stress assignment in German patients with surface dyslexia. *Brain and Language*, 87, 114–115. doi:10.1016/S0093-934X(03)00225-6
- Jouravlev, O., & Lupker, S. L. (2014). Stress consistency and stress regularity effects in Russian. *Language and Cognitive Processes*, 29, 605–619.  
doi:10.1080/01690965.2013.813562
- Kay, J., Lesser, R., & Coltheart, M. (1992). *Psycholinguistic assessments of language processing in aphasia (PALPA)*. Sussex, UK: Lawrence Erlbaum Ltd.
- Kelly, M. H. (2004). Word onset patterns and lexical stress in English. *Journal of Memory and Language*, 50, 231–244. doi:10.1016/j.jml.2003.12.002
- Kelly, M. H., & Bock, J., K. (1988). Stress in time. *Journal of Experimental Psychology: Human Perception and Performance*, 14, 389–403. doi:10.1037/0096-1523.14.3.389
- Kelly, M. H., Morris, J., & Verrechia, L. (1998). Orthographic cues to lexical stress: Effects on naming and lexical decision. *Memory & Cognition*, 26, 822–832.  
doi:10.3758/BF03211401
- Kuperman, V., Stadthagen-Gonzalez, H., & Brysbaert, M. (2012). Age-of-acquisition ratings for 30 thousand English words. *Behavior Research Methods*, 44, 978–990.  
doi:10.3758/s13428-012-0210-4

- Marshall, J. C., & Newcombe, F. (1973). Patterns of paralexia: A psycholinguistic approach. *Journal of Psycholinguistic Research*, 2, 175–199. doi:10.1007/BF01067101
- Miceli, G., & Caramazza, A. (1992). The assignment of word stress in oral reading: Evidence from a case of acquired dyslexia. *Cognitive Neuropsychology*, 10, 272–296. doi:10.1080/02643299308253465
- Mioshi, E., Dawson, K., Mitchell, J., Arnold, R., & Hodges, J. (2006). The Addenbrooke's cognitive examination revised (ACE-R): A brief cognitive test battery for dementia screening. *International Journal of Geriatric Psychiatry*, 21, 1078–1085. doi:10.1002/gps.1610
- Monsell, S., Doyle, M. C., & Haggard, P. N. (1989). Effects of frequency on visual word recognition tasks: Where are they? *Journal of Experimental Psychology: General*, 118, 43–71. doi: 10.1037/0096-3445.118.1.43
- Nelson, H. E., & Wilson, J. (1991) *National Adult Reading Test (NART)*, Windsor, UK: NFER-Nelson.
- Osterrieth, P.A. (1944). Filetest de copie d'une figure complex: Contribution a l'etude de la perception et de la memoire [The test of copying a complex figure: A contribution to the study of perception and memory]. *Archives de Psychologie*, 30, 286–356.
- Paizi, D., Zoccolotti, P., & Burani, C. (2011). Lexical stress assignment in Italian developmental dyslexia. *Reading and Writing*, 24, 443–461. doi:10.1007/s11145-010-9236-0

- Pritchard, S. C., Coltheart, M., Palethorpe, S., & Castles, A. (2012). Nonword reading: Comparing dual-route cascaded and connectionist dual-process models with human data. *Journal of Experimental Psychology: Human Perception and Performance*, 38, 1268–1288.
- Psychological Corporation (1997). *WAIS-III/WMS-III Technical Manual*. San Antonio: Harcourt Brace & Company.
- Rastle, K., & Coltheart, M. (1999). Serial and strategic processing in reading aloud. *Journal of Experimental Psychology: Human Perception and Performance*, 25, 482–503. doi:10.1037//0096-1523.25.2.482
- Rastle, K., & Coltheart, M. (2000). Lexical and nonlexical print-to-sound translation of disyllabic words and nonwords. *Journal of Memory and Language*, 42, 342–364. doi:10.1037//0096-1523.25.2.482
- Rastle, K., & Davis, M. H. (2008). Morphological decomposition based on the analysis of orthography. *Language and Cognitive Processes*, 23, 942–971. doi:10.1080/01690960802069730
- Rey, A. (1941) Psychological examination in cases of traumatic encephalopathy. *Archives de Psychologie*, 28, 286–340.
- Schock, J., Cortese, M. J., & Khanna, M. M. (2012). Imageability estimates for 3,000 disyllabic words. *Behavior Research Methods*, 44, 374–379. doi:10.3758/s13428-011-0162-0

Spreen, O., & Strauss, E. (1998). *A compendium of neuropsychological tests: Administration, norms and commentary*. New York: Oxford University Press

Sulpizio, S., Burani, C., & Colombo, L. (2015). The process of stress assignment in reading aloud: Critical issues from studies on Italian, *Scientific Studies of Reading*, *19*, 5-20, doi:10.1080/10888438.2014.976340

Trenta, M., Benassi, M., Di Filippo, G., Pontillo, M., and Zoccolotti, P. (2013). Developmental dyslexia in a regular orthography: can the reading profile be reduced to strategic control? *Cognitive Neuropsychology*, *30*, 147–171. doi: 10.1080/02643294.2013.814569

Warrington, E. K. (1984). *Recognition memory test*. Windsor, UK: NFER-Nelson.

Zoccolotti, P., De Luca, M., Di Pace, E., Judica, A., Orlandi, M., & Spinelli, D. (1999). Markers of developmental surface dyslexia in a language (Italian) with high grapheme–phoneme correspondence. *Applied Psycholinguistics*, *20*, 191–216.

using the phonemic vocabulary of the dual-route cascaded model. The glossary of the DRC phonemic vocabulary is provided in Appendix A.

<sup>2</sup> Because of Patient 1's severe semantic memory impairment (see Table 1) we also examined the potential influence of word imageability on this patient's stress error data. In a separate regression analysis we included word imageability ratings (Schock, Cortese, & Khanna, 2012) as an additional covariate and found no imageability effect on the probability of producing second-syllable stress.

	<b>Patient</b>				
	1	2	3	4	5
<b>GENERAL COGNITION</b>					
Total ACE-R score	50/100	25/100	37/100	32/100	62/100
<b>Diagnosis</b>	PPA - SD	PPA - NFPA	AD	AD	FTD
<b>MEMORY</b>					
<i>Verbal/Nonverbal STM</i>					
Digit Span Forward/Backward (WMS-III)	6 vs 4	2 vs 0	4 vs 1	5 vs 2	4 vs 3
Spatial span (WMS-III)	6 vs 5	2 vs 0	4 vs 2	4 vs 3	3 vs 3
<i>Verbal/Nonverbal LTM</i>					
HVLT-R	3-3-4	---	3-5-6	---	3-3-4
ROCF copy	36/36	apraxia	apraxia	apraxia	32/36
ROCF delay	15/36	---	---	---	2/36
<i>Verbal/Nonverbal Recognition</i>					
Words (Warrington)	27/50	28/50	28/50	40/50	26/50
Faces (Warrington)	38/50	33/50	31/50	36/50	28/50
<b>LANGUAGE</b>					
<i>Verbal Fluency</i>					
Semantic					
<i>Animals (ACE-R)</i>	14	2	3	5	7
Letter					
<i>FAS</i>	13	4	3	0	16
<i>Comprehension</i>					
PPT	36/52	46/52	37/52	44/52	46/52
<i>Naming</i>					
Picture (PALPA 40)	11/40	5/40	34/40	38/40	32/40
<i>Repetition</i>					
Repetition (PALPA 9)					
<i>Words</i>	40/40	8/20	40/40	40/40	39/40
<i>Nonwords</i>	40/40	3/20	39/40	40/40	37/40

<i>Regular</i>	29/30	25/30	29/30	28/30	29/30
<i>Irregular</i> ( <i>regularization errors</i> )	12/30 (83.3%)	19/30 (63.6%)	25/30 (40.0%)	24/30 (50.0%)	26/30 (100%)
Nonwords (PALPA 36)	23/24	21/24	20/24	21/24	21/24
Coltheart & Leahy					
<i>Regular</i>	29/30	25/30	29/30	29/30	29/30
<i>Irregular</i> ( <i>regularization errors</i> )	12/30 (94.4%)	10/30 (65.0%)	19/30 (54.6%)	23/30 (57.1%)	20/30 (80.0%)
<i>NWs</i>	28/30	25/30	26/30	28/30	28/30

Note. **Diagnoses:** PPA = Primary Progressive Aphasia; SD = Semantic Dementia; NFPA = Non-Fluent Progressive Aphasia; AD = Alzheimer's Disease; FTD = Fronto-Temporal Dementia; apraxia = constructional apraxia. **Assessments:** ACE-R = Addenbrooke's Cognitive Examination Revised (Mioshi, Dawson, Arnold, & Hodges, 2006); Coltheart & Leahy tests, (Coltheart & Leahy, 1996); FAS letter fluency test (Spreen & Strauss, 1998); HVLRT-R = Hopkins Verbal Learning Test - Revised (Benedict, Schretlen, Groninger, & Brandt, 1998); NART = National Adult Reading Test (Nelson & Wilson, 1991); PALPA = Psycholinguistic Assessment of Language Processing in Aphasia (Kay, Lesser, & Coltheart, 1992); PPT = Pyramids & Palm Trees Test (Howard & Patterson, 1992); ROCF = Rey-Osterrieth Complex Figure test (Osterrieth, 1944); Warrington Words and Faces tests (Warrington, 1984); WMS-III = Wechsler Memory Scale-III (Psychological Corporation, 1997)

	<b>Word frequency</b>	<b>Word length</b>	<b>Coltheart's N</b>	<b>AoA</b>	<b>Bigram frequency</b>
<b>Prefix W-S</b> (e.g., remind)	Mean: 25.77 SD: 28.72	Mean: 6.46 SD: 0.89	Mean: 0.54 SD: 0.71	Mean: 9.26 SD: 2.16	Mean: 43.89 SD: 26.97
<b>Prefix S-W</b> (e.g., reflex)	Mean: 23.79 SD: 42.95	Mean: 6.30 SD: 0.91	Mean: 0.76 SD: 1.32	Mean: 10.22 SD: 2.42	Mean: 42.64 SD: 25.29
<b>NoPrefix S-W</b> (e.g., scandal)	Mean: 25.10 SD: 28.99	Mean: 6.72 SD: 0.78	Mean: 0.28 SD: 0.50	Mean: 7.52 SD: 2.24	Mean: 48.93 SD: 19.87

% 2 <sup>nd</sup> -syllable Stress			
Patient	Prefix W-S	Prefix S-W	NoPrefix S-W
	'regular'	'irregular'	
1	98.0	88.0 (84.0)	46.0 (42.0)
2	98.0	91.5 (89.3)	22.9 (22.9)
3	87.8	42.9 (36.7)	2.0 (2.0)
4	97.8	62.5 (47.9)	4.3 (4.3)
5	91.5	69.4 (67.3)	6.8 (6.8)

Note. Percentage of second syllable stress assignments for Prefix S-W and NoPrefix S-W words denote stress errors (pure stress errors are shown in parentheses).

## PREFIXES REPEL STRESS

Table 4.

Wald test statistics for the main effect of condition and pairwise comparisons, and the main effect of embedded word ratio (OR) is reported where appropriate.

		Patient				
		1	2	3	4	5
<b>Condition</b>		W(2) = 23.25 $p < .001$	W(2) = 32.09 $p < .001$	W(2) = 36.06 $p < .001$	W(2) = 28.76 $p < .001$	W(2) = 35.14 $p < .001$
<b>Prefix W-S</b> <b>'regular'</b>	<b>NoPrefix</b>	W(1) = 16.95 $p < .001$	W(1) = 22.49 $p < .001$	W(1) = 26.31 $p < .001$	W(1) = 27.76 $p < .001$	W(1) = 34.97 $p < .001$
		OR: 146.9	OR: 356.1	OR: 330.2	OR: 774.0	OR: 139.5
<b>Prefix S-W</b> <b>'irregular'</b>	<b>S-W</b>	W(1) = 15.92 $p < .001$	W(1) = 22.21 $p < .001$	W(1) = 6.64 $p = .01$	W(1) = 13.53 $p < .001$	W(1) = 19.27 $p < .001$
		OR: 23.6	OR: 70.2	OR: 16.8	OR: 22.7	OR: 28.3
<b>Embedded Word</b>		W(1) < .01 $p = .971$	W(1) = .56 $p = .456$	W(1) < .01 $p = .944$	W(1) = .20 $p = .652$	W(1) = .07 $p = .793$

	% 2 <sup>nd</sup> -syllable Stress		
	Prefix W-S 'regular'	Prefix S-W 'irregular'	NoPrefix S-W
RC00 algorithm	94.0	93.9 (93.9)	8.0 (8.0)
CDP++ model	81.2	50.0 (48.9)	2.1 (2.1)

Note. Percentage of second syllable stress assignments for Prefix S-W and NoPrefix S-W words denote stress errors (pure stress errors are shown in parentheses).

## PREFIXES REPEL STRESS

Appendix A. The phonemic vocabulary used in the transcription of the patients' and models' responses. Each symbol marked in bold in examples of English words. Adapted from Rastle & Coltheart (1999).

Symbol	Example	Symbol	Example	Symbol	Example	Symbol
1	<b>b</b> ay	J	<b>c</b> heap	h	<b>h</b> ad	v
2	<b>b</b> uy	N	<b>b</b> ang	i	<b>b</b> ean	w
3	<b>b</b> urn	Q	<b>p</b> ot	j	<b>y</b> ank	z
4	<b>b</b> oy	S	<b>s</b> heep	k	<b>c</b> ad	#
5	<b>n</b> o	T	<b>t</b> hin	l	<b>l</b> ad	{
6	<b>b</b> row	U	<b>p</b> ut	m	<b>m</b> ad	_
7	<b>p</b> eer	V	<b>p</b> utt	n	<b>n</b> at	@
8	<b>p</b> air	Z	<b>m</b> easure	p	<b>p</b> at	(schwa)
9	<b>p</b> oor	b	<b>b</b> ad	r	<b>r</b> at	
D	<b>t</b> hen	d	<b>d</b> ad	s	<b>s</b> ap	
E	<b>p</b> et	f	<b>f</b> at	t	<b>t</b> ack	
I	<b>p</b> it	g	<b>g</b> ame	u	<b>u</b> boon	

## PREFIXES REPEL STRESS

Appendix B. Stimuli and patients' and models' reading aloud responses.

Stimulus	Condition	Patient					RCO
		1	2	3	4	5	
adjust	Prefix W-S	@d_Vst'	@d_Vst'	@d_Vst'	@d_Vst'	@d_Vst'	@d_Vst'
alive	Prefix W-S	@l2v'	@l2v'	@l2v'	@l2v'	NR *	@l2v'
amuse	Prefix W-S	@mjuz'	@mus'	{mjuz'	@mjuz'	@mjuz'	@mjuz'
arise	Prefix W-S	@r2z'	@r2s'	@r2z'	@r2s'	@r2s'	@r2z'
confirm	Prefix W-S	kQnf3m'	kQnf3m'	kQnf3m'	kQnf3m'	kQnf3m'	k@n'
contempt	Prefix W-S	kQntEmpt'	kQntEmpt'	'kQntEmpt	kQmEnt'	kQntEmpt'	k@n'
design	Prefix W-S	dIz2n'	diz2n'	dIz2n'	diz2n'	NR *	dIsIn'
disgrace	Prefix W-S	dIsgr1s'	dIsgr1s'	dIsgr1s'	dIsgr1s'	dIsgr1s'	dIsgr1s'
enforce	Prefix W-S	Enf9s'	Inf9s'	Inf9s'	Inf9s'	Inf9s'	Inf9z'
explain	Prefix W-S	Ekspl1n'	Ekspl1n'	Ekspl1n'	Eks1n'	NR *	Ekspl1n'
improve	Prefix W-S	Impruv'	Impruv'	Impr5v'	Impruv'	Impruv'	Impruv'
inform	Prefix W-S	Inf9m'	Inf9m'	Inf9m'	Inf9m'	Inf9m'	Inf9m'
inspire	Prefix W-S	Insp2r'	Insp'2@ *	Insp2r'	Insp'2@ *	Insp2r'	'Insp2r'
intact	Prefix W-S	Int{kt'	Int{kt'	Int{kt'	Int{kt'	Int{kt'	Int{k'
intense	Prefix W-S	IntEns'	IntEns'	IntEns'	IntEst'	IntEns'	IntEn'
preside	Prefix W-S	prIz2d'	prIs2d'	prIs2d'	prIs2d'	prIs2d'	prIs2d'



## PREFIXES REPEL STRESS

## Appendix B. (continued)

Stimulus	Condition	Patient					RCO
		1	2	3	4	5	
endure	Prefix W-S	Endj9'	Indj9'	Endju'	Indj9'	Indj9'	Ind9
exploit	Prefix W-S	Eksp14t'	Eksp14t'	'Eksp14t	Eksp15t'	Eksp14t'	Eksp
immune	Prefix W-S	Imun'	Imun'	Imjun'	NR *	Imjun'	Imm
include	Prefix W-S	Inkljud'	Inkljud'	'Inklud	Inkljud'	Inkljud'	Inclu
insist	Prefix W-S	InsIst'	InsIst'	InsIst'	InsIst'	InsIst'	InsIs
invade	Prefix W-S	Inv1d'	Inv1d'	Inv1d'	Inv1d'	Inv1d'	Inv1
invoke	Prefix W-S	Inv5k'	Inv5k'	Inv5k'	Inv5k'	Inv5k'	Inv5
precise	Prefix W-S	prIs2s'	prEsis'	prIs2s'	pris'Es@ *	pris2s'	prIs2
promote	Prefix W-S	pr@m5t'	prQmt5'	pr@m5t'	prQm'EtI *	pr@m5t'	pr@r
protect	Prefix W-S	prQtEkt'	pr5tEkt'	'prQ_Ekt	pr5tEst'	pr5tEkt'	pr@r
reduce	Prefix W-S	rIdjus'	rIdjus'	rIdjus'	rIdVs'	rIdjus'	rIdju
reflect	Prefix W-S	rIfIEkt'	rIfIEkt'	rIfIEkt'	rIfIEks'	rIfIEkt'	rIfIE
regime	Prefix W-S	rI_2m'	ri_2m'	rI_im'	ri_2m'	'rE_1m	rIg2r
resist	Prefix W-S	rIsIst'	rizIst'	riInsIst'	rizIst'	rizIst'	rIsIs
revenge	Prefix W-S	rIvEn_'	rIvEn_'	rIvEn_'	rIvEn_'	rIvEn_'	rIvEn
submit	Prefix W-S	sVbmIt'	'sVbmIt	s@bmIt'	sVbmIt'	'sVbmIt	s@b
survive	Prefix W-S	s3v2n'	s3v2v'	s3v2v'	s3v2v'	s3v2v'	's3vI

## PREFIXES REPEL STRESS

## Appendix B. (continued)

Stimulus	Condition	Patient					RCO
		1	2	3	4	5	
adverse	Prefix S-W	{dv3s'	{dv3s'	{dv3s'	'{dv3s	{dv3s'	@dv
agate	Prefix S-W	{g1t'	@g1t'	{g1t'	Non Int. *	@g1t'	@g1
agent	Prefix S-W	{gEnt'	1_Ent'	'1_Ent	'1_@nt	1nJEnt'	@_E
atoll	Prefix S-W	@tQl'	@t5l'	'{tQl	@t5l'	@t5l'	@tQ
context	Prefix S-W	kQntEkst'	kQntEkst'	'kQntEkst	kQntEkst'	kQntEkst'	k@n
convent	Prefix S-W	kQnvEnt'	kQnvEnt'	'kQnv3t	kQnvEkt'	'kQnv@nt	k@n
depot	Prefix S-W	dEpQt'	dEpQt'	dIpQt'	'dEpQt	dip5t'	dIpQ
discord	Prefix S-W	dIsk9d'	dIsk9d'	dIsk9d'	dIsk9d'	dIsk9d'	dIsk9
entrails	Prefix S-W	Entr1lz'	Entr1ls'	Ent1lz'	Entr1ls'	Entr1ls'	Intr1
expert	Prefix S-W	Eksp3t'	Eksp3t'	Eksp3t'	Eksp3t'	Eksp3t'	Eksp
impulse	Prefix S-W	ImpVls'	ImpVls'	ImpVls'	ImpVls'	ImpVls'	ImpV
income	Prefix S-W	Ink5m'	InkVm'	'INkVm	InkVm'	InkVm'	Ink5
inland	Prefix S-W	'Inl{nd	Inl{nd'	'Inl@nd	'Inl@nd	'Inl{nd	Inl{r
input	Prefix S-W	'InpVt	InpVt'	'InpVt	pVt5t'	InpVt'	InpV
insight	Prefix S-W	Ins2t'	Ins2t'	'Ins2t	Ins2t'	Ins2t'	Ins2t
pretext	Prefix S-W	prItEkst'	prEtEkst'	prItEkst'	pri{ks'	priEkst'	prItE
product	Prefix S-W	prQdVkt'	'prQdjus	'prQdVkt	'prQdVkt	'prQdVkt	pr@

## PREFIXES REPEL STRESS

Appendix B. (continued)

Stimulus	Condition	Patient					RCO
		1	2	3	4	5	
program	Prefix S-W	prQgr{m'	'pr5gr@m	'prQgr@m	'pr5gr{m	'pr5gr@m	pr@
recent	Prefix S-W	risEnt'	risEnt'	'risEnt	risEnt'	rIsEnt'	rIsEn
reflex	Prefix S-W	rflEkt'	rflEks'	'rflEks	rflEks'	rflEks'	rflE
regent	Prefix S-W	ri_Ent'	ri_Ent'	ri_Ent'	ri_Ent'	'ri_Ent	ri_En
respite	Prefix S-W	rIsp2t'	risp2d'	rIsp2t'	risp2t'	risp2t'	rIsp2
retail	Prefix S-W	rIt1l'	rit1l'	'rit1l	rit1l'	rit1l'	rIt1l'
subway	Prefix S-W	s@bw1'	sVbw1'	'sVbw1	sVbw1'	'sVbw1	s@b
surface	Prefix S-W	s3f1s'	's3f *	's3f1s	's3f1s	s3f1s'	's3f1
adjunct	Prefix S-W	@d_VNkt'	{d_Vnkt'	{d_VNkt'	'{d *	{d_Vnkt'	@d_
apron	Prefix S-W	'{prQn	lprQn'	'lprQn	'lpr@n	'lprQn	@pr
aspect	Prefix S-W	{spEkt'	@spEkt'	{spEkt'	QspIt'	'{sbEkt	@sp
athlete	Prefix S-W	{Tlit'	{Tlit'	{TIE'tIk *	{Tlit'	{Tlit'	@Tl
concert	Prefix S-W	kQns3t'	kQns3t'	'kQns3t	'kQns3t	'kQns3t	k@n
congress	Prefix S-W	kQngrEs'	kQngrEs'	'kQngrEs	'kQngrEs	kQngrEs'	k@n
denim	Prefix S-W	'dEnIm	dEnIm'	'dEmIn	'dEmIn	dinIm'	dInIm
distant	Prefix S-W	dIst{nt'	dIst{nt'	'dIst@nt	dIst{nt'	'dIst@nt	dIst{
enzyme	Prefix S-W	Enz2m'	Enz2m'	Enz2m'	'Enz2m	Enz'2mIn*	'Enz

## PREFIXES REPEL STRESS

## Appendix B. (continued)

Stimulus	Condition	Patient					RCO
		1	2	3	4	5	
exile	Prefix S-W	Eks2l'	Ekz2l'	Eks2l'	Ekz2l'	Ekz2l'	Egz2l'
impasse	Prefix S-W	Imp{s'	Imp1s'	Imp{s'	Imp1s'	Imp{s'	Imp1s'
index	Prefix S-W	IndEks'	IndEks'	'IndEks	IndEks'	IndEks'	IndEks'
infant	Prefix S-W	Inf{nt'	Inf{nt'	'Inf@nt	'Inf@nt	'Inf@nt	Inf{nt'
injure	Prefix S-W	In_9'	In_j9'	In_j9'	'In_@	In_j9'	In_9'
instinct	Prefix S-W	InstInkt'	'InstItIt *	'Inst@nt	'InsVnt	'InstInkt	InstInkt'
prelude	Prefix S-W	prIljud'	priljud'	'prIljud	priljud'	priljud'	prIljud'
problem	Prefix S-W	'pr5blEm	'prQbl@m	'prQblEm	'prQbl@m	'prQbl@m	pr@blEm
prospect	Prefix S-W	prQspEkt'	PrQpQspEkt *	prQspEkt'	prQpEs'	prQspEkt'	pr@spEkt'
refuge	Prefix S-W	rIfju_'	rIfju_'	rIfju_'	rIfju'	rIfju_'	rIfju_'
relic	Prefix S-W	'rElIk	'rIlIk	ris2kl'	rElIk'	'rIlIk	rIlIk'
relish	Prefix S-W	rElIS'	rElIS'	'rElIS	'rElIS	rIlIS'	rIlIS'
rescue	Prefix S-W	rIskju'	rIskju'	riskj9'	rIskju'	'rEskju	rIskju'
revel	Prefix S-W	rIvEl'	rIvEl'	rIvEl'	rIv2l'	rIvEl'	rIvEl'
suburb	Prefix S-W	sVb3b'	sub3b'	'sVb3b	sub3b'	sub3b'	s@b3b'
survey	Prefix S-W	s3v1'	s3v1'	's3v1	's3v1	s3v1'	's3v2l'
bishop	NoPrefix S-W	'bIS@p	'bISQp	'bIS@p	'bISQp	'bISQp	'bISQp

## PREFIXES REPEL STRESS

## Appendix B. (continued)

Stimulus	Condition	Patient					RC0
		1	2	3	4	5	
borough	NoPrefix S-W	bQr6'	'b3@	'b{r@	NR *	'b3@	'b9@
chemist	NoPrefix S-W	'kEmIst	'kEmIst	'kEmIst	'tEmIst	NR *	'JEm
climate	NoPrefix S-W	kl2m1t'	klIm1t'	'kl2m1t	'kl2m@nt	'kl2m1t	klIm
donkey	NoPrefix S-W	'dQNki	'dQnki	'dQNkI	'dQnki	'dQnki	'dQN
famine	NoPrefix S-W	f@m2n'	'f{mIn	'f{mIn	'f{mIn	'f{mIn	'f{m
fortress	NoPrefix S-W	'f9trEs	f9trEs'	'f9trEs	'f9trEs	'f9trEs	f@tr
margin	NoPrefix S-W	'm#_In	'm#gIn	'm#_In	'm#_In	'm#_In	'm#g
merchant	NoPrefix S-W	'm3J{nt	'm3J@nt	'm3J@nt	'm3J@nt	'm3J@nt	'm3J
message	NoPrefix S-W	mEs1_'	'mEs1_	'mEsI_	'mEsI_	'mEs1_	'miss
monarch	NoPrefix S-W	mQn#J'	mQn#J'	'mQn#k	mQn'#ki *	'mQn@k	'mQn
orchard	NoPrefix S-W	'9J#d	'9J3d	'9J@d	'9J3d	'9J3d	'9J@
palace	NoPrefix S-W	p{11s'	'p{1@s	'p11s	'p{11s	'pV11s	'p{11
parent	NoPrefix S-W	'p#{nt	'p8@nt	'p{r@nt	'p8@nt	'p8@nt	'p#@
porridge	NoPrefix S-W	p9I_'	'pQrI_	'pQrI_	'pQrI_	'pQrI_	'p9rI
purpose	NoPrefix S-W	p3p5z'	'p3pQs	'p3p@s	'p3sp@s	'p3p@s	p3p5
silent	NoPrefix S-W	's2l@nt	's2lEnt	's2l@nt	's2lEnt	's2lEnt	's2l@
tactic	NoPrefix S-W	't{ktIk	't{ktIk	't{ktIk	NR *	NR *	't{ktl

## PREFIXES REPEL STRESS

## Appendix B. (continued)

Stimulus	Condition	Patient					RCO
		1	2	3	4	5	
talent	NoPrefix S-W	t@l{nt'	't{lEnt	't{l@nt	't{lEnt	't{lEnt	't{l@
textile	NoPrefix S-W	tEkst2l'	'tEkst2l	'tEkst2l	'Ekst2l	'tEkst2l	'tEks
toilet	NoPrefix S-W	't4lEt	't4lEt	't4lEt	't4lEt	't4lEt	't4lIt
twilight	NoPrefix S-W	'tw2l2t	tw2l2t'	'tw2l2t	'tw2l2t	'tw2l2t	'twIl
urban	NoPrefix S-W	'3b@n	'3bVn	'3b@n	'3bVn	NR *	'3b@
walnut	NoPrefix S-W	'w{lInVt	'w9lnVt	'w{lInVt	'w9lnVt	'w9lnVt	'w{lIn
welcome	NoPrefix S-W	'wElk@m	'wElk@m	wElk@m	'wElk@m	'wElk@m	'wElk
biscuit	NoPrefix S-W	bIskjut'	'bIsk@t	'bIskIts	'bIsk@t	NR *	'bIsk
campus	NoPrefix S-W	'k{mp@s	'k{mpVs	'k{mpVs	'k{mpVs	k{mpVs'	'k{m
chapel	NoPrefix S-W	'J{p@l	'_{p@l	'J{p@l	'_{p@l	'_{p@l	'JlP@
culture	NoPrefix S-W	kVltj9'	'kVIJ9	'kVIJ9	'kVIJ9	NR *	'kVlt
diamond	NoPrefix S-W	d2mQnd'	'd2j@mQnd *	d2m@nd	'd2m@nd	'd2mQnd	'dI@
fountain	NoPrefix S-W	f6nt1n'	f6nt1n'	'f6nt1n	f6nt1n'	'f6nt@n	'f6nt
fragile	NoPrefix S-W	fr{_2l'	'fr{_2l	'fr{_2l	'fr{_2l	'fr{_2l	'fr{g
frequent	NoPrefix S-W	frEkwEnt'	frEkwEnt'	'frikwEnt	'frEkwEnt	NR *	'frEk
garbage	NoPrefix S-W	k#b1_'	'g#r{_	'g#b1_	'g#bI_	'g#b1_	'g#bI
helmet	NoPrefix S-W	'hElmEt	'hElmEt	'hElm@t	'hElmEt	'hElmIt	'hElm

## PREFIXES REPEL STRESS

## Appendix B. (continued)

Stimulus	Condition	Patient					RCO
		1	2	3	4	5	
lettuce	NoPrefix S-W	'lEtVs	lEtjus'	'lEtVs	'lEt@s	'lEtIs	'lEtju
minute	NoPrefix S-W	mInjut'	mInjut'	'mIn@t	mInut'	'mInIt	mInj
mountain	NoPrefix S-W	m6nt1n'	'm6nt1n	'm6nt1n	'm6nt@n	'm6nt@n	'm6n
mustard	NoPrefix S-W	mVst#d'	'mVst#d	'mVst@d	'mVst#d	'mVst#d	'mVs
pattern	NoPrefix S-W	p{t3n'	'p{t3n	'p{t3n	'p{t3n	'p{t3n	'p{t3
pigeon	NoPrefix S-W	'pI_Vn	pI_Vn'	'pI_@n	'pI_Vn	'pI_@n	'p2_
portrait	NoPrefix S-W	p9tr1nt'	p9tr1t'	'p9tr@t	'p9tr1t	p9tr1t'	'p9tr
publish	NoPrefix S-W	'pVblIS	'blIS *	'pVblIS	'pQlIS	'pVblIS	'pVb
robot	NoPrefix S-W	'rQb@t	'r5bQt	r5bQt'	'r5bQt	'r5bQt	'rQb
scandal	NoPrefix S-W	'sk{nd@l	'sk{nd@l	'sk{nd@l	'sk{nd@l	'sk{nd@l	'sk{n
spinach	NoPrefix S-W	spIn{J'	spInIJ'	'spIn@S	'spInIJ	spIn1k'	'spIn
trolley	NoPrefix S-W	'tr5li	'trQli	'trQlI	'trQli	'trQli	'trQl
verdict	NoPrefix S-W	'v3dIkt	'v3dIk	'v3dIkt	'v3dIkt	'v3dIkt	'v3dI
volume	NoPrefix S-W	'vQljum	'vQlEm	'vQljum	'vQljum	'vQljum	'vQl
window	NoPrefix S-W	'wInd5	'wInd5	'wInd5	'wInd5	'wInd5	'wIn

Note. Non-Int. = Non-Intelligible; NR = No Response \* = Not included in the statistical analyses