



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in :
Journal of Lipid Research

Cronfa URL for this paper:
<http://cronfa.swan.ac.uk/Record/cronfa30987>

Paper:

Abdel-Khalik, J., Yutuc, E., Crick, P., Gustafsson, J., Warner, M., Roman, G., Talbot, K., Gray, E., Griffiths, W., Turner, M. & Wang, Y. (2017). Defective cholesterol metabolism in amyotrophic lateral sclerosis. *Journal of Lipid Research*, jlr.P071639
<http://dx.doi.org/10.1194/jlr.P071639>

This article is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Authors are personally responsible for adhering to publisher restrictions or conditions. When uploading content they are required to comply with their publisher agreement and the SHERPA RoMEO database to judge whether or not it is copyright safe to add this version of the paper to this repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

Defective cholesterol metabolism in amyotrophic lateral sclerosis^S

Jonas Abdel-Khalik,* Eylan Yutuc,* Peter J. Crick,* Jan-Åke Gustafsson,[†] Margaret Warner,[†] Gustavo Roman,[§] Kevin Talbot,** Elizabeth Gray,** William J. Griffiths,^{1,*} Martin R. Turner,^{1,**} and Yuqin Wang^{1,*}

Swansea University Medical School,* Swansea, United Kingdom; Department of Biology and Biochemistry,[†] Center for Nuclear Receptors and Cell Signaling, University of Houston, Houston, TX; Methodist Neurological Institute,[§] Methodist Hospital, Houston, TX; and Nuffield Department of Clinical Neurosciences,** University of Oxford, John Radcliffe Hospital, Oxford, United Kingdom

ORCID IDs: 0000-0002-4129-6616 (W.J.G.)

Abstract As neurons die, cholesterol is released in the central nervous system (CNS); hence, this sterol and its metabolites may represent a biomarker of neurodegeneration, including in amyotrophic lateral sclerosis (ALS), in which altered cholesterol levels have been linked to prognosis. More than 40 different sterols were quantified in serum and cerebrospinal fluid (CSF) from ALS patients and healthy controls. In CSF, the concentration of cholesterol was found to be elevated in ALS samples. When CSF metabolite levels were normalized to cholesterol, the cholesterol metabolite 3 β ,7 α -dihydroxycholest-5-en-26-oic acid, along with its precursor 3 β -hydroxycholest-5-en-26-oic acid and product 7 α -hydroxy-3-oxocholest-4-en-26-oic acid, were reduced in concentration, whereas metabolites known to be imported from the circulation into the CNS were not found to differ in concentration between groups. Analysis of serum revealed that (25R)26-hydroxycholesterol, the immediate precursor of 3 β -hydroxycholest-5-en-26-oic acid, was reduced in concentration in ALS patients compared with controls. We conclude that the acidic branch of bile acid biosynthesis, known to be operative in-part in the brain, is defective in ALS, leading to a failure of the CNS to remove excess cholesterol, which may be toxic to neuronal cells, compounded by a reduction in neuroprotective 3 β ,7 α -dihydroxycholest-5-en-26-oic acid.—Abdel-Khalik, J., E. Yutuc, P. J. Crick, J.-Å. Gustafsson, M. Warner, G. Roman, K. Talbot, E. Gray, W. J. Griffiths, M. R. Turner, and Y. Wang. **Defective cholesterol metabolism in amyotrophic lateral sclerosis.** *J. Lipid Res.* 2017. 58: 267–278.

Supplementary key words oxysterols • mass spectrometry • cytochrome P450 • nuclear receptors/LXR • brain lipids • bile acids and salts/biosynthesis • cholestenic acids • neurodegeneration.

Amyotrophic lateral sclerosis (ALS) is a heterogeneous, progressive, and fatal neurodegenerative disease characterized by variable loss of upper and lower motor neurons (1). Biomarkers that are sensitive to the progression of disease have the potential to shorten therapeutic trials and provide new drug targets. Study of the metabolome offers the potential to identify disease-specific patterns for ALS, possibly providing such biomarkers and new insights into the deranged biochemical pathways associated with ALS. Metabolic studies in blood and cerebrospinal fluid (CSF) have been performed utilizing proton nuclear magnetic resonance spectrometry (2–4), GC/MS, and LC/MS (5, 6). Significant differences between patients and controls have been observed, consistent with a range of pathogenic mechanisms that have been described in ALS, including mitochondrial dysfunction, oxidative stress, excitotoxicity, neuroinflammation, and hypermetabolism (7).

Abbreviations: ALS, amyotrophic lateral sclerosis; CSF, cerebrospinal fluid; CYP, cytochrome P450; DHC, dehydrocholesterol; diHC, dihydroxycholesterol; LXR, liver X receptor; PLS, primary lateral sclerosis; 3 β -HCA, 3 β -hydroxycholest-5-en-26-oic acid; 3 β ,7 α -diHCA, 3 β ,7 α -dihydroxycholest-5-en-26-oic acid; 7O-C, 7-oxocholesterol; 7 α -HC, 7 α -hydroxycholesterol; 7 α -HCO, 7 α -hydroxycholest-4-en-3-one; 7 α H,26-nor-C-3,24-diO, 7 α -hydroxy-26-norcholest-4-ene-3,24-dione; 7 α ,24-diH,3O-CA, 7 α ,24-dihydroxy-3-oxocholest-4-en-26-oic acid; 7 α ,25-diHC, 7 α ,25-dihydroxycholesterol; 7 α ,25-diHCO, 7 α ,25-dihydroxycholest-4-en-3-one; 7 α ,26-diHCO, 7 α ,26-dihydroxycholest-4-en-3-one; 7 α H,3O-CA, 7 α -hydroxy-3-oxocholest-4-en-26-oic acid; 7 β -HC, 7 β -hydroxycholesterol; 7-DHC, 7-dehydrocholesterol; 24-DHC, desmosterol; 24S-HC, 24S-hydroxycholesterol; 25-D₃, 25-hydroxyvitamin D₃; 25-HC, 25-hydroxycholesterol; 26-HC, (25R)26-hydroxycholesterol.

¹To whom correspondence should be addressed.

e-mail: w.j.griffiths@swansea.ac.uk (W.J.G.); martin.turner@ndcn.

ox.ac.uk (M.R.T.); y.wang@swansea.ac.uk (Y.W.)

^SThe online version of this article (available at <http://www.jlr.org>) contains a supplement.

This work was supported by UK Biotechnology and Biological Sciences Research Council Grants BB/I001735/1 (to W.J.G.) and BB/L001942/1 (to Y.W.); Welsh Government Grant A4B (to W.J.G. and Y.W.); a Swedish Science Council grant (to J.-Å.G.); and Robert A. Welch Fund Grant E-0044 (to J.-Å.G.). J.A.-K. was supported by a PhD studentship from Imperial College Healthcare Charities. M.R.T. and E.G. were funded by joint Medical Research Council / Motor Neuron Disease Association Fellowships G0701923 and MR/K01014X/1 (to M.R.T.).

*Author's Choice—Final version free via Creative Commons CC-BY license.

Manuscript received 25 August 2016 and in revised form 1 November 2016.

Published, JLR Papers in Press, November 3, 2016

DOI 10.1194/jlr.P071639

Lipids, distinct from other components of the metabolome (8), are nonpolar or amphipathic in nature and require separate analysis from more water-soluble metabolites. Cholesterol, both the nonesterified molecule and its esters with fatty acids, represents a major component of the total lipid content of cells in vertebrates. On a number of fronts, cholesterol and its metabolites represent potential biomarkers for ALS. High plasma levels of cholesterol have been suggested to be neuroprotective for ALS and to be associated with an increased survival time (9–11), but other data suggest that accumulation of cholesterol esters and ceramides mediate oxidative stress in motor neurons in ALS (12), while the gene cytochrome P450 27A1 (*CYP27A1*), encoding cholesterol (25R)26-hydroxylase (also known as sterol 27-hydroxylase), the first enzyme in the extrahepatic part of the bile acid biosynthesis pathway, was recently identified as a susceptibility gene for sporadic ALS (13). In addition, statins, the cholesterol-lowering drugs that inhibit HMG-CoA reductase, have been suggested to accelerate functional decline in ALS patients (14).

ALS is a neurodegenerative disease and as neurons die, cholesterol is released from cells. Cholesterol and its metabolites in the CNS are predominantly in a nonesterified form (15), and it the nonesterified molecules that are responsible for regulating cholesterol homeostasis (16, 17) and are the ligands to nuclear receptors (18, 19) and ligands to G-protein-coupled receptors (20, 21). We aimed to characterize and quantify nonesterified cholesterol and the widest range of nonesterified metabolites in CSF and serum from ALS patients compared with a group of healthy controls (see **Table 1** for molecules analyzed). An LC/MS approach was adopted to exploit charge-tagging to ensure maximum sensitivity and to gain structural information allowing identification of unexpected sterols (supplemental Fig. S1); a hydrolysis step was not included in the sample preparation protocol (22, 23).

MATERIALS AND METHODS

Human samples

Serum was from 35 patients diagnosed with ALS (24 male, 11 female, mean age 65), 6 patients diagnosed with primary lateral sclerosis (PLS; 2 male, 4 female, mean age 69), and 24 control samples (12 male, 12 female, mean age 58). CSF was from ALS patients ($n = 20$; 15 male, 5 female, mean age 61) and controls ($n = 15$; 12 male, 3 female, mean age 75). Serum and CSF samples were obtained from ALS and PLS patients and healthy volunteers (typically patient spouses and friends) as part of The Oxford Study for Biomarkers in Motor Neuron Disease (BioMOx). Two expert neurologists (M.R.T. and K.T.) made the diagnosis of ALS and PLS according to standard criteria. All participants provided informed consent, and the study was approved by the South Central Oxford Ethics Committee B. Additional control CSF samples for method development were from a study performed at Methodist Hospital, Houston, TX.

Methods

Serum. One hundred microliters of serum was added dropwise to a solution of 1,050 μ l of absolute ethanol containing 2 ng of 25-[26,26,26,27,27,27- 2 H $_6$]hydroxyvitamin D $_3$ (25-[2 H $_6$]D $_3$,

Toronto Research Chemicals, Ontario, Canada), 2 ng of 7 α ,25-[26,26,26,27,27,27- 2 H $_6$]dihydroxycholesterol (7 α ,25-[2 H $_6$]dihC), 20 ng of 7 α -[25,26,26,26,27,27,27- 2 H $_7$]hydroxycholesterol (7 α -[2 H $_7$]HC), 20 ng of 24R/S-[25,26,26,26,27,27,27- 2 H $_7$]hydroxycholesterol (24R/S-[2 H $_7$]HC), 20 ng of 22R-[25,26,26,26,27,27,27- 2 H $_7$]hydroxycholesterol-4-en-3-one (22R-[2 H $_7$]HCO), and 20 μ g of [25,26,26,26,27,27,27- 2 H $_7$]cholesterol (deuterated sterols from Avanti Polar Lipids Inc., Alabaster, AL) with sonication in an ultrasonic bath. After 5 min, 350 μ l of water was added to make the solution 70% ethanol. After a further 5 min of sonication, the solution was centrifuged at 17,000 g at 4°C for 30 min.

To separate cholesterol from the more polar oxysterols and steroid acids, the sample solution, now in 1.5 ml of 70% ethanol, was loaded onto a 200-mg Certified Sep-Pak C $_{18}$ column (Waters, Elstree, Herts, U.K.) preconditioned with 4 ml of absolute ethanol and 6 ml of 70% ethanol. The solvent flow rate through the column was at a rate of \sim 0.25 ml/min assisted by negative pressure at the column outlet generated by a vacuum manifold. The flow-through (1.5 ml) was combined with a column wash of 70% ethanol (5.5 ml) to give fraction SPE1-Fr1 (7 ml) in which 25-D $_3$, oxysterols, cholestenic, and cholenoic acids elute. A second fraction (SPE1-Fr2) was collected by eluting with a further 4 ml of 70% ethanol, before fraction 3 containing cholesterol and sterols of similar hydrophobicity was eluted with 2 ml of absolute ethanol (SPE1-Fr3). Finally, a fourth fraction eluting in 2 ml of absolute ethanol was collected, containing lipid more hydrophobic than cholesterol (SPE1-Fr4). Each fraction was divided into two equal subfractions, (A) and (B), and allowed to dry overnight under vacuum. Note that a hydrolysis step was not included in the sample preparation procedure, and by removing cholesterol from its metabolites in the first step of sample preparation, subsequent confusion with ex vivo oxidation products is eliminated.

Lyophilized material was reconstituted in 100 μ l of propanol-2-ol. Each sample was vortexed thoroughly. To subfractions (A) 1,000 μ l of 50 mM phosphate buffer (pH 7) was added, containing 3.0 μ l of cholesterol oxidase from *Streptomyces* sp. (2 mg/ml in H $_2$ O, 44 units/mg protein, Sigma-Aldrich Ltd, Dorset, England). The reaction mixture was left at 37°C for 1 h, after which the reaction was quenched with 2,000 μ l of methanol. Subfractions (B) were treated identically to subfractions (A), but in the absence of cholesterol oxidase. One hundred fifty microliters of glacial acetic acid was added to subfractions (A) and (B) and vortexed. One hundred ninety milligrams of [2 H $_5$]Girard P (GP) reagent, bromide salt (23), was added to subfractions (A), 150 mg of [2 H $_0$]GP reagent, chloride salt (TCI Europe, Zwiindrecht, Belgium), was added to subfractions (B) to allow for subsequent simultaneous LC/MS analysis; see below (supplemental Fig. S1). After vortexing, the derivatization reaction was left to proceed at room temperature, overnight in the dark.

To remove excess derivatization reagent, a 60-mg Oasis HLB (Waters) column was conditioned with 6 ml of 100% methanol, 6 ml of 10% methanol, and 4 ml of 70% methanol before the samples (3,250 μ l, \sim 70% organic) were loaded. The sample vessel was washed with 1 ml of 70% methanol, and the wash was applied to the column. One milliliter of 35% methanol was used to rinse the column, and the combined 5 ml of effluent was diluted with 4 ml of water to give 9 ml of 35% methanol solution. This solution was reapplied to the column, followed by a rinse of the column with 1 ml of 17.5% methanol. The 10 ml of combined effluent was diluted with 9 ml of water to give a 19-ml solution of 17.5% methanol. Reapplying the 19-ml solution to the column completed the recycling procedure by which sterols, oxysterols, cholestenic, and cholenoic acids were extracted by the column, whereas derivatization reagents eluted to waste. Six milliliters of 10% methanol was added to wash the column. Finally, 25-D $_3$, oxysterols, cholestenic and cholenoic acids, and more nonpolar sterols

TABLE 1. Sterols, oxysterols, cholestenic, and choleonic acids analyzed in serum and CSF

Systematic Name (trivial name)	Abbreviation	Lipid Maps ID	Authentic Standard
3 β -Hydroxycholestatriene ^a		NA	No
Cholesta-5,7-dien-3 β -ol (7-dehydrocholesterol) + cholesta-5,8-dien-3 β -ol ^b	7-DHC + 8-DHC	LMST01010069 /NA	Yes
Cholesta-5,24-dien-3 β -ol (desmosterol)	24-DHC	LMST01010016	Yes
Cholest-5-en-3 β -ol (cholesterol)	C	LMST01010001	Yes
Cholesta-5,8-diene-3 β ,24(or 25)-diol	24(or 25)H-8DHC	NA	No
Cholesta-5,8-diene-3 β ,26-diol	26H-8-DHC	NA	No
3 β -Hydroxycholest-5-en-7,8-epoxide	7,8-EC	NA	Yes
9,10-Secocholesta-5Z,7E,10-triene-3S,25-diol (25-Hydroxyvitamin D ₃)	25-D ₃	LMST03020246	Yes
9,10-Secocholesta-5Z,7E,10-triene-3S,24,25-triol (24,25-Dihydroxyvitamin D ₃)	24,25-D ₃	LMST03020273 or LMST03020274	Yes
Cholest-4(or 5)-ene-3 β ,6 β -diol (6 β -hydroxycholesterol) ^c	6 β -HC	NA	Yes
Cholest-5-ene-3 β ,7 α -diol (7 α -hydroxycholesterol)	7 α -HC	LMST01010013	Yes
7 α -Hydroxycholest-4-en-3-one	7 α -HCO	LMST01010271	Yes
Cholest-5-ene-3 β ,7 β -diol (7 β -hydroxycholesterol)	7 β -HC	LMST01010047	Yes
3 β -Hydroxycholest-5-en-7-one (7-oxocholesterol)	7O-C	LMST01010049	Yes
3 β -Hydroxycholest-5-en-24-one (24-oxocholesterol)	24O-C	LMST01010133	Yes
Cholest-5-ene-3 β ,24S-diol (24S-hydroxycholesterol)	24S-HC	LMST01010019	Yes
Cholest-5-ene-3 β ,25-diol (25-hydroxycholesterol)	25-HC	LMST01010018	Yes
Cholest-5-ene-3 β , (25R) 26-diol (27-hydroxycholesterol)	26-HC	LMST01010057	Yes
Cholest-5-ene-3 β ,12 α -diol (12 α -hydroxycholesterol) ^d	12 α -HC	NA	No
Cholest-5-ene-3 β ,7 α ,12 α -triol (7 α ,12 α -dihydroxycholesterol) ^d	7 α ,12 α -diHC	LMST04030165	No
7 α ,12 α -Dihydroxycholest-4-en-3-one ^d	7 α ,12 α -diHCO	LMST04030114	No
Cholest-5-ene-3 β ,7 α ,25-triol (7 α ,25-dihydroxycholesterol)	7 α ,25-diHC	LMST04030166	Yes
7 α ,25-Dihydroxycholest-4-en-3-one	7 α ,25-diHCO	LMST04030107	Yes
Cholest-5-ene-3 β ,7 α ,26-triol (7 α ,26-dihydroxycholesterol)	7 α ,26-diHC	LMST04030081	Yes
7 α ,26-Dihydroxycholest-4-en-3-one	7 α ,26-diHCO	LMST04030157	Yes
3 β ,22-Dihydroxycholest-5-en-24-one ^d	22-HC-24-O	NA	No
3 β ,5 α -Dihydroxycholest-7-en-6-one (dihydrocholestenone)		LMST01010260	Yes
Cholesta-5-en-3 β ,24,25-triol (24,25-dihydroxycholesterol)	24,25-diHC	NA	Yes
3 β -Hydroxycholest-5-en-26-oic acid	3 β -HCA	LMST04030072	Yes
3-Oxocholest-4-en-26-oic acid	3O-CA	NA	Yes
3 β -Hydroxycholesta-5,7-dien-26-oic acid ^f	3 β H-7-DHCA	NA	Yes
3-Oxocholesta-4,6-dien-26-oic acid ^f	3O-6-DHCA	NA	Yes
3 β ,7 α -Dihydroxycholest-5-en-26-oic acid ^f	3 β ,7 α -diHCA	LMST04030148	Yes
7 α -Hydroxy-3-oxocholest-4-en-26-oic acid ^f	7 α H,3O-CA	LMST04030149	Yes
Total 3 β ,7 α -Dihydroxycholest-5-en-26-oic acid ^g	3 β ,7 α -diHCA	LMST04030148	Yes
Total 7 α -Hydroxy-3-oxocholest-4-en-26-oic acid ^h	7 α H,3O-CA	LMST04030149	Yes
3 β ,7 β -Dihydroxycholest-5-en-26-oic acid	3 β ,7 β -diHCA	NA	Yes
3 β -Hydroxy-7-oxocholest-5-en-26-oic acid	3 β H,7O-CA	LMST04030215	Yes
3 β ,22,25-Trihydroxycholest-5-en-24-one ^d	22,25-diHC-24-O	NA	No
7 α ,x,y-Trihydroxycholest-4-en-3-one ^d	7 α ,x,y-triHCO	NA	No
7 α ,24(or 26),25-Trihydroxycholest-4-en-3-one ^d	7 α ,24(or 26),25-triHCO	NA	No
3 β ,7 α ,24-Trihydroxycholest-5-en-26-oic acid	3 β ,7 α ,24-triHCA	NA	Yes
7 α ,24-Dihydroxy-3-oxocholest-4-en-26-oic acid	7 α ,24-diH,3O-CA	NA	Yes
3 β ,7 β ,24-Trihydroxycholest-5-en-26-oic aci ⁱ	3 β ,7 β ,24-triHCA	NA	No
7 β ,24-Dihydroxy-3-oxocholest-4-en-26-oic aci ⁱ	7 β ,24-diH,3O-CA	NA	No
3 β ,7 α ,25-Trihydroxycholest-5-en-26-oic acid	3 β ,7 α ,25-triHCA	NA	Yes
7 α ,25-Dihydroxy-3-oxocholest-4-en-26-oic acid	7 α ,25-diH,3O-CA	NA	Yes
3 β ,7 β ,25-Trihydroxycholest-5-en-26-oic aci ⁱ	3 β ,7 β ,25-triHCA	NA	No
7 β ,25-Dihydroxy-3-oxocholest-4-en-26-oic acid ⁱ	7 β ,25-diH,3O-CA	NA	No
7 α ,x-Dihydroxy-3-oxocholest-4-en-26-oic acid ^d	7 α ,x-diH,3O-CA	NA	No
3 β ,7 α ,12 α -Trihydroxycholest-5-en-26-oic acid ^d	3 β ,7 α ,12 α -triHCA	NA	No
7 α ,12 α -Dihydroxy-3-oxocholest-4-en-26-oic acid ^d	7 α ,12 α -diH,3O-CA	LMST04030150	No
Trihydroxy-3-oxocholest-4-en-26-oic acid		NA	No
7 α -Hydroxy-3,24-bisoxocholest-4-en-26-oic acid ^d	7 α H,3,24-diO-CA	NA	No
7 α -Hydroxy-26-nor-cholest-4-ene-3,24-dione ^d	7 α H,26-nor-C-3,24-diO	NA	No
3 β ,7 α -Dihydroxychol-5-en-24-oic acid	3 β ,7 α - Δ^5 -BA	LMST04010217	Yes
7 α -Hydroxy-3-oxochol-4-en-24-oic acid	7 α H,3O- Δ^4 -BA	LMST04010239	Yes
3 β ,7 β -Dihydroxychol-5-en-24-oic acid	3 β ,7 β - Δ^5 -BA	LMST04010218	Yes

NA, not applicable.

^aLocation of double bonds unknown.

^bThe 7-DHC isomerizes to 8-DHC.

^cDehydration product of cholestane-3 β ,5 α ,6 β -triol.

^dAuthentic standard not available; annotation based on retention time, exact mass, and MS³ fragmentation.

^eThe 3 β ,7 α -diHCA dehydrates to 3 β H-7-DHCA.

^fThe 7 α H,3O-CA dehydrates to 3O-6-DHCA.

^gSum of intact 3 β ,7 α -diHCA and its dehydration product 3 β H-7-DHCA.

^hSum of intact 7 α H,3O-CA and its dehydration product 3O-6-DHCA.

ⁱIdentification based on comparison to 7 α -isomer.

were eluted from the column with 3×1 ml 100% methanol (SPE2-Fr1,2,3) and 1 ml of absolute ethanol (SPE2-Fr4). Oxysterols, 25-D₃, cholestenic, and choleonic acids elute in the first 2 ml from the column (SPE2-Fr1,2). Cholesterol and sterols of similar polarity elute over 3 ml of methanol (SPE2-Fr1,2,3).

For 25-D₃, oxysterol, cholestenic, and choleonic acid analysis, just before analysis by LC/MS, an equal aliquot of SPE2-Fr1,2 from subfractions (A) and (B) was combined together in one tube. For cholesterol analysis equal volumes of SPE2-Fr1,2,3 from subfractions (A) and (B) were combined. In both cases the combined effluent in methanol was diluted with water to achieve a concentration of 60% methanol, i.e., the initial composition of the mobile phases during LC analyses.

LC/MS and LC/MSⁿ analysis. Analysis was performed on a LTQ-Orbitrap Elite (Thermo Fisher Scientific, Hemel Hempstead, UK) equipped with an electrospray probe, and a Dionex Ultimate 3000 LC system (Dionex; now Thermo Fisher Scientific), essentially as described previously (22, 23). For each injection,

three to five scan events were performed: one high-resolution (120,000, full-width at half maximum height at m/z 400) MS scan event in the Orbitrap analyzer in parallel with two to four multi-stage fragmentation (MSⁿ) scan events in the LTQ linear ion trap. Quantification was performed by stable isotope dilution or by using isotope-labeled structural analogs.

CSF. Two hundred fifty microliters of CSF was added dropwise to 2,100 μ l of absolute ethanol containing 2 ng of 24R/S-[²H₇]HC, 2 ng of 7 α -[²H₇]HC, 2 ng of 22R-[²H₇]HCO, 0.4 ng of 7 α ,25-[²H₆]diHC, 4 ng of 25-[²H₆]D₃, and 0.8 μ g of [²H₇]cholesterol with sonication in an ultrasonic bath. Six hundred fifty microliters of water was added, and the solution was sonicated for 5 min, followed by centrifugation at 2,400 *g* at 4°C for 30 min.

The CSF sample solution in 3 ml of 70% ethanol was loaded onto a 200-mg Certified Sep-Pak C₁₈ column as in the serum protocol. However, the 3-ml flow-through was combined with a column wash of only 4 ml of 70% ethanol to give SPE1-Fr1 (7 ml) in which 25-D₃, oxysterols, cholestenic, and choleonic acids elute. The




Fig. 1. Concentration of 25-D₃ and cholesterol metabolites in serum. Box and whiskers plots showing the concentrations (ng/ml) of 25-D₃ (A), 7 α -HC (B), 7 α -HCO (C), 7 α ,25-diHCO (D), 26-HC (E), and 3 β -HCA (F) in serum from ALS (n = 35) and PLS (n = 6) patients and healthy controls (n = 24). The bottom and top of the box represents the first and third quartiles, and the band inside the box represents the median. The whiskers extend to the most extreme data points, which are no more than 1.5 times the range between first and third quartile distant from the box. Points beyond that are plotted individually. Data for other sterols can be found in supplemental Table S1. Univariate *t* tests were performed against the control group. * *P* < 0.05; ** *P* < 0.01.

remainder of the sample preparation procedure, including derivatization, was identical to that used for serum, with the exception that the final eluates from SPE2 were lyophilized and reconstituted in 60% methanol prior to sample injection on the LC system.

Statistics

An ANOVA was run for each sterol. Univariate *t* tests were performed against the control group: **P* < 0.05; ***P* < 0.01. Concentrations given in the text are means ± SDs.

RESULTS

Analysis of sterols, oxysterols, and cholestenic and choleonic acids in serum

Using enzyme-assisted derivatization for sterol analysis technology (22) in combination with LC/MS with MS³, we

analyzed serum from 35 patients diagnosed with ALS (24 male, 11 female, mean age 65) and 24 control samples (12 male, 12 female, mean age 58). Six patients diagnosed with the upper motor neuron-only, very slowly progressive variant of ALS, termed PLS (2 male, 4 female, mean age 69), were separately compared with the control group. Concentrations of nonesterified cholesterol, two of its precursors, and >40 metabolites were measured (see Table 1 for compounds and abbreviations and supplemental Table S1 for quantitative data; note that a hydrolysis step was not carried out, so all concentrations are for nonesterified compounds). We failed to find any sexual dimorphism in the control sample set and thus included both sexes in the current study.

There was no significant difference in the concentrations of cholesterol or its precursors desmosterol (24-DHC) or 7-dehydrocholesterol (7-DHC) between ALS, PLS, or




Fig. 2. Pathways of cholesterol metabolism initiated by the enzymes cholesterol 25-hydroxylase (CH25H) and CYP46A1. Changes in sterols concentration in CSF and serum are indicated by blue and red arrows, respectively. The direction of change corresponds to the direction of the arrow. Enzymes catalyzing the indicated reactions are shown where known. Univariate *t* tests were performed against the control group. * *P* < 0.05; ** *P* < 0.01.




Fig. 3. Pathway of cholesterol metabolism initiated by the enzymes CYP7A1 and CYP27A1. Changes in sterols concentration in CSF and serum are indicated by blue and red arrows, respectively. The direction of change corresponds to the direction of the arrow. Enzymes catalyzing the indicated reactions are shown where known. Enzyme abbreviations used are as follows: ACOX2, acyl-CoA oxidase 2, branched chain; AMACR, α -methylacyl-CoA racemase; BACS, bile acyl-CoA synthetase; DBP, D-bifunctional protein or multifunctional enzyme type 2 (HSD17B4); SCPx, sterol carrier protein x; VLCS, very long chain acyl-CoA synthetase. Univariate t tests were performed against the control group. * $P < 0.05$; ** $P < 0.01$.

control samples. Note we report here the concentration of 7-DHC as the sum of 7-DHC and its isomer 8-DHC, which are only partially resolved on our LC system. The 8-DHC is an enzymatic product of 7-DHC (24). Similarly, there was no significant difference in the concentration of the 7-DHC metabolite 25-hydroxyvitamin D₃ (25-D₃) in serum of the ALS patients compared with controls (Fig. 1).

The first steps of all cholesterol metabolism lead to hydroxycholesterol isomers, collectively known as oxysterols. A minor pathway, initiated in activated macrophages, leads to 25-hydroxycholesterol (25-HC), followed by subsequent metabolism to 7 α ,25-dihydroxycholesterol (7 α ,25-diHC) and 7 α ,25-dihydroxycholest-4-en-3-one (7 α ,25-diHCO, Fig. 2). Of these metabolites, 7 α ,25-diHCO was elevated in ALS serum ($P < 0.01$). 7 α -Hydroxycholesterol (7 α -HC) and 7 α -hydroxycholest-4-en-3-one (7 α -HCO) are the first members of the neutral pathway of bile acid biosynthesis (Fig. 3) (25); however, neither oxysterol showed a difference in concentration between ALS, PLS, or control samples, and nor did 7 β -hydroxycholesterol (7 β -HC) or 7-oxocholesterol (7O-C). 7O-C, 7 α -HC, and 7 β -HC can be formed via free radical oxidation of cholesterol, and these metabolites have been suggested as markers of oxidant stress (26).

Conversely, (25R)26-hydroxycholesterol (26-HC), the first member of the extrahepatic part of the acidic pathway of bile acid biosynthesis, was decreased in ALS and PLS serum compared with controls ($P < 0.01$), as was 3 β -hydroxycholest-5-en-26-oic acid (3 β -HCA) in ALS serum ($P < 0.05$). CYP27A1 is the enzyme responsible for hydroxylation of cholesterol at (25R)C-26 and then subsequent oxidation to the (25R)26 carboxylic acid. Note that we use the systematic nomenclature (27), where hydroxylation of the terminal carbon of the cholesterol side chain introducing 25R stereochemistry results in (25R)26-hydroxycholesterol (28). Unless stated otherwise, R stereochemistry is generally assumed at C-25. The nonsystematic name for 26-HC, widely used in the literature on account of the nomenclature of its synthesizing enzyme, CYP27A1, is 27-hydroxycholesterol. Levels of 26-HC in the circulation are often correlated to cholesterol, with high cholesterol levels associated with high levels of 26-HC (29). We thus decided to normalize for each sample the level of 26-HC and other metabolites to cholesterol to determine whether the observed differences were still maintained. When normalized to cholesterol (supplemental Table S2), elevation in concentration of 7 α ,25-diHCO was retained in ALS, as was a decrease in the concentration of 26-HC. Interestingly, in an earlier study, Wuolikainen et al. (30) found that “total” 26-HC in male ALS patients was similarly reduced. In combination, these results suggest reduced transcription, translation, or activity of cholesterol (25R)26-hydroxylase in ALS.

Analysis of sterols, oxysterols, and cholestenic and cholenic acids in CSF

As with serum, levels of cholesterol, 7-DHC (plus 8-DHC), and desmosterol were measured in CSF from ALS patients ($n = 20$; 15 male, 5 female, mean age 61) and controls ($n = 15$; 12 male, 3 female, mean age 75). The

concentrations of both desmosterol ($P < 0.05$) and cholesterol ($P < 0.01$) were found to be elevated in CSF from ALS patients (Fig. 4, supplemental Table S3). Because ALS is a neurodegenerative disease, we speculated that, when cholesterol is released by neurons as they die, it would be metabolized by CYP46A1 to 24S-hydroxycholesterol (24S-HC) and by CYP27A1 to members of the acidic pathway of bile acid biosynthesis (Figs. 2 and 3, respectively) (25). Surprisingly, when normalized to cholesterol, 24S-HC concentration was found to be reduced ($P < 0.01$) in CSF, as were members of the acidic pathway of bile acid biosynthesis (supplemental Table S4). Intriguingly, concentrations of 7 α -HC, 7 α -HCO, 7 β -HC, and 7O-C, which originate from the CYP7A1-initiated arm of the bile acid biosynthesis pathway and enter the CSF from the circulation, did not differ between ALS patients and controls (Fig. 5). This can




Fig. 4. Concentration of cholesterol and its precursors in CSF. Box and whisker plots showing concentrations (ng/ml) of 24-DHC (A), 7-DHC + 8-DHC (B), and cholesterol (C) in CSF from ALS ($n = 20$) patients and healthy controls ($n = 15$). Box and whiskers are as described in the Fig. 1 legend. Data for other sterols can be found in supplemental Table S3. Univariate t tests were performed against the control group. * $P < 0.05$; ** $P < 0.01$.

be explained because CYP7A1 is not expressed in the CNS, being liver-specific (Fig. 3) (25). The level of 26-HC was not significantly lower in CSF from ALS patients (0.09 ± 0.06 ng/ μ g cholesterol) compared with controls (0.13 ± 0.07 ng/ μ g cholesterol), but its downstream metabolites 3 β -HCA ($P < 0.01$), 3 β ,7 α -dihydroxycholest-5-en-26-oic acid (3 β ,7 α -diHCA; $P < 0.05$), and 7 α -hydroxy-3-oxocholest-4-en-26-oic acid (7 α H,3O-CA; $P < 0.01$) were all reduced in concentration (ng/ μ g cholesterol) in ALS CSF (Fig. 6). Whereas (25R)26-carboxylation by CYP27A1 occurs in the mitochondria, 7 α -hydroxylation by CYP7B1, the oxysterol 7 α -hydroxylase, and oxidation of the 3 β -hydroxy group by hydroxysteroid dehydrogenase (HSD) 3B7 occur in the endoplasmic reticulum, further metabolism to bile acids proceeds in the peroxisome (31). Peroxisomal metabolites found in CSF include 7 α ,24-dihydroxy-3-oxocholest-4-en-26-oic acid (7 α ,24-diH,3O-CA); 7 α -hydroxy-3,24-bisoxocholest-4-en-26-oic acid (7 α H,3,24-diO-CA), which is also observed as the decarboxylated

dione; 7 α -hydroxy-26-norcholest-4-ene-3,24-dione (7 α H,26-nor-C-3,24-diO); and the β -oxidation product 7 α -hydroxy-3-oxocholest-4-en-24-oic acid (7 α H,3O- Δ^4 -BA). With the exception of the latter compound, all of the peroxisomal intermediates were decreased ($P < 0.05$ or 0.01 ; ng/ μ l cholesterol) in CSF from ALS patients.

DISCUSSION

The major route for cholesterol metabolism in the CNS is conversion to 24S-HC via the neuron-specific enzyme CYP46A1 (32, 33), accounting for approximately two-thirds of cholesterol metabolism in brain, and elevated levels of total 24S-HC have been found in CSF of patients with some neurodegenerative diseases, e.g., Alzheimer's disease (AD) (34, 35). The term total describes the sum of both esterified and nonesterified sterol. In most studies, concentrations of total cholesterol or of total hydroxycholesterols are




Fig. 5. Concentration of CYP7A1, CYP46A1, and CH25H pathway metabolites in CSF. Box and whisker plots showing concentrations (ng/ μ g cholesterol) of 7 α -HC (A), 7 β -HC (B), 7O-C (C), 24S-HC (D), 7 α ,25-diHCO (E), and 7 α ,26-diHCO (F) in CSF from ALS (n = 20) patients and healthy controls (n = 15). Box and whiskers are as described in the Fig. 1 legend. Data for other sterols can be found in supplemental Table S4. Univariate *t* tests were performed against the control group. * $P < 0.05$; ** $P < 0.01$.




Fig. 6. Concentration of acidic pathway metabolites in CSF. Box and whisker plots showing concentrations (ng/μg cholesterol) of 3β-HCA (A), 3β,7α-diHCA (B), 7αH,3O-CA (C), 3β,7β-diHCA (D), 7α,24-diH,3O-CA (E), and 7αH,26-nor-C3,24-diO (F) in CSF from ALS (n = 20) patients and healthy controls (n = 15). Box and whiskers are as described in the Fig. 1 legend. Data for other sterols can be found in supplemental Table S4. Univariate *t* tests were performed against the control group. * *P* < 0.05; ** *P* < 0.01.

measured where sterols esterified with fatty acids are hydrolyzed and the total sterol is assayed, although it is the nonesterified sterols that are biologically active. Usually the ratio of esterified to nonesterified sterol in plasma is approximately 10:1 (36). Surprisingly, levels of total 24S-HC are decreased in blood plasma of patients with advanced AD, presumably as a consequence of a reduced number of active neurons producing 24S-HC (37). In a recent study, Wuolikainen et al. found total 24S-HC plasma levels to be increased in female ALS patients, but not in male patients, whereas total 26-HC was reduced in male ALS patients, but not females (30). Side-chain hydroxycholesterols, i.e., 22R-HC, 24S-HC, 25-HC, and 26-HC, and also cholestenic acids, are ligands to liver X receptors (LXRs), the β-isoform being particularly expressed in brain, and, interestingly, LXRβ^{-/-} mice show adult-onset motor neuron pathology

(38, 39). Importantly, LXRβ is involved in control of lipogenesis and cholesterol homeostasis, and these mice showed increased cholesterol levels in spinal cord, gliosis, and inflammation preceding motor neuron loss and clinical disease onset (40). In the present study, a loss of LXR signaling may result from reduced CNS levels of oxysterol and cholestenic acid LXR-ligands, ultimately leading to motor neuron pathology.

Although cholesterol cannot cross the blood-brain barrier and be imported to, or exported from, the brain, 24S-HC, 26-HC, and other cholesterol metabolites can cross the blood-brain barrier (35, 41). Heverin et al. (42) have shown that 26-HC is imported to the brain, whereas Meaney et al. (43) have shown that its acidic metabolite 7αH,3O-CA is exported from the brain to circulation. Further studies by Iuliano et al. (44) have shown that a precursor of this

conversely, reduced levels of 24S-HC in the CNS, as observed in ALS, should result in enhanced cholesterol biosynthesis and *Il1b* transcription, inflammasome activity, and neuroinflammation.

In summary, we have identified elevated cholesterol concentrations in the CSF of ALS patients. When normalized to cholesterol, concentrations of glial-derived acidic metabolites are reduced, but cholesterol metabolites imported from the circulation are not. These data point to impaired activity of the CYP27A1 enzyme leading to a failure of the CNS to remove excess cholesterol, which may be toxic to neuronal cells, compounded by a reduction in neuroprotective $3\beta,7\alpha$ -diHCA and other LXR ligands (Fig. 7) **■**

The authors thank members of the European Network for Oxysterol Research (<http://oxysterols.com/>) for informative discussions.

REFERENCES

- Kiernan, M. C., S. Vucic, B. C. Cheah, M. R. Turner, A. Eisen, O. Hardiman, J. R. Burrell, and M. C. Zoing. 2011. Amyotrophic lateral sclerosis. *Lancet*. **377**: 942–955.
- Gray, E., J. R. Larkin, T. D. Claridge, K. Talbot, N. R. Sibson, and M. R. Turner. 2015. The longitudinal cerebrospinal fluid metabolomic profile of amyotrophic lateral sclerosis. *Amyotroph. Lateral Scler. Frontotemporal Degener.* **16**: 456–463.
- Blasco, H., P. Corcia, C. Moreau, S. Veau, C. Fournier, P. Vourc'h, P. Emond, P. Gordon, P. F. Pradat, J. Praline, et al. 2010. 1H-NMR-based metabolomic profiling of CSF in early amyotrophic lateral sclerosis. *PLoS One*. **5**: e13223.
- Kumar, A., L. Bala, J. Kalita, U. K. Misra, R. L. Singh, C. L. Khetrapal, and G. N. Babu. 2010. Metabolomic analysis of serum by (1) H NMR spectroscopy in amyotrophic lateral sclerosis. *Clin. Chim. Acta*. **411**: 563–567.
- Wuolikainen, A., T. Moritz, S. L. Marklund, H. Antti, and P. M. Andersen. 2011. Disease-related changes in the cerebrospinal fluid metabolome in amyotrophic lateral sclerosis detected by GC/TOFMS. *PLoS One*. **6**: e17947.
- Blasco, H., P. Corcia, P. F. Pradat, C. Bocca, P. H. Gordon, C. Veyrat-Durebex, S. Mavel, L. Nadal-Desbarats, C. Moreau, D. Devos, et al. 2013. Metabolomics in cerebrospinal fluid of patients with amyotrophic lateral sclerosis: an untargeted approach via high-resolution mass spectrometry. *J. Proteome Res.* **12**: 3746–3754.
- Turner, M. R., R. Bowser, L. Bruijn, L. Dupuis, A. Ludolph, M. McGrath, G. Manfredi, N. Maragakis, R. G. Miller, S. L. Pullman, et al. 2013. Mechanisms, models and biomarkers in amyotrophic lateral sclerosis. *Amyotroph. Lateral Scler. Frontotemporal Degener.* **14**(Suppl 1): 19–32.
- Sud, M., E. Fahy, D. Cotter, K. Azam, I. Vadivelu, C. Burant, A. Edison, O. Fiehn, R. Higashi, K. S. Nair, et al. 2016. Metabolomics Workbench: An international repository for metabolomics data and metadata, metabolite standards, protocols, tutorials and training, and analysis tools. *Nucleic Acids Res.* **44**: D463–D470.
- Dupuis, L., P. Corcia, A. Fergani, J. L. Gonzalez De Aguilar, D. Bonnefont-Rousselot, R. Bittar, D. Seilhean, J. J. Hauw, L. Lacomblez, J. P. Loeffler, et al. 2008. Dyslipidemia is a protective factor in amyotrophic lateral sclerosis. *Neurology*. **70**: 1004–1009.
- Sutedja, N. A., Y. T. van der Schouw, K. Fischer, E. M. Sizoo, M. H. Huisman, J. H. Veldink, and L. H. Van den Berg. 2011. Beneficial vascular risk profile is associated with amyotrophic lateral sclerosis. *J. Neurol. Neurosurg. Psychiatry*. **82**: 638–642.
- Dorst, J., P. Kuhnlein, C. Hendrich, J. Kassubek, A. D. Sperfeld, and A. C. Ludolph. 2011. Patients with elevated triglyceride and cholesterol serum levels have a prolonged survival in amyotrophic lateral sclerosis. *J. Neurol.* **258**: 613–617.
- Cutler, R. G., W. A. Pedersen, S. Camandola, J. D. Rothstein, and M. P. Mattson. 2002. Evidence that accumulation of ceramides and cholesterol esters mediates oxidative stress-induced death of motor neurons in amyotrophic lateral sclerosis. *Ann. Neurol.* **52**: 448–457.
- Diekstra, F. P., C. G. Saris, W. van Rhenen, L. Franke, R. C. Jansen, M. A. van Es, P. W. van Vught, H. M. Blauw, E. J. Groen, S. Horvath, et al. 2012. Mapping of gene expression reveals CYP27A1 as a susceptibility gene for sporadic ALS. *PLoS One*. **7**: e35333.
- Zheng, Z., L. Sheng, and H. Shang. 2013. Statins and amyotrophic lateral sclerosis: a systematic review and meta-analysis. *Amyotroph. Lateral Scler. Frontotemporal Degener.* **14**: 241–245.
- Dietschy, J. M., and S. D. Turley. 2004. Thematic review series: brain Lipids. Cholesterol metabolism in the central nervous system during early development and in the mature animal. *J. Lipid Res.* **45**: 1375–1397.
- Zhang, Y., K. M. Lee, L. N. Kinch, L. Clark, N. V. Grishin, D. M. Rosenbaum, M. S. Brown, J. L. Goldstein, and A. Radhakrishnan. 2016. Direct demonstration that Loop1 of Scap binds to Loop7: a CRUCIAL EVENT IN CHOLESTEROL HOMEOSTASIS. *J. Biol. Chem.* **291**: 12888–12896.
- Radhakrishnan, A., Y. Ikeda, H. J. Kwon, M. S. Brown, and J. L. Goldstein. 2007. Sterol-regulated transport of SREBPs from endoplasmic reticulum to Golgi: oxysterols block transport by binding to Insig. *Proc. Natl. Acad. Sci. USA*. **104**: 6511–6518.
- Evans, R. M., and D. J. Mangelsdorf. 2014. Nuclear Receptors, RXR, and the Big Bang. *Cell*. **157**: 255–266.
- Gustafsson, J. A. 2015. Control of gene expression by novel metabolic intermediates. *J. Steroid Biochem. Mol. Biol.* **153**: 102–104.
- Hannedouche, S., J. Zhang, T. Yi, W. Shen, D. Nguyen, J. P. Pereira, D. Guerini, B. U. Baumgarten, S. Roggo, B. Wen, et al. 2011. Oxysterols direct immune cell migration via EBI2. *Nature*. **475**: 524–527.
- Liu, C., X. V. Yang, J. Wu, C. Kuei, N. S. Mani, L. Zhang, J. Yu, S. W. Sutton, N. Qin, H. Banie, et al. 2011. Oxysterols direct B-cell migration through EBI2. *Nature*. **475**: 519–523.
- Griffiths, W. J., P. J. Crick, Y. Wang, M. Ogundare, K. Tuschl, A. A. Morris, B. W. Bigger, P. T. Clayton, and Y. Wang. 2013. Analytical strategies for characterization of oxysterol lipidomes: liver X receptor ligands in plasma. *Free Radic. Biol. Med.* **59**: 69–84.
- Crick, P. J., T. William Bentley, J. Abdel-Khalik, I. Matthews, P. T. Clayton, A. A. Morris, B. W. Bigger, C. Zerbinati, L. Tritapepe, L. Iuliano, et al. 2015. Quantitative charge-tags for sterol and oxysterol analysis. *Clin. Chem.* **61**: 400–411.
- Griffiths, W. J., J. Abdel-Khalik, P. J. Crick, M. Ogundare, C. H. Shackleton, K. Tuschl, M. K. Kwok, B. W. Bigger, A. A. Morris, A. Honda, et al. 2016. Sterols and oxysterols in plasma from Smith-Lemli-Opitz syndrome patients. *J. Steroid Biochem. Mol. Biol.* Epub ahead of print. March 11, 2016; doi: 10.1016/j.jsmb.2016.03.018.
- Russell, D. W. 2003. The enzymes, regulation, and genetics of bile acid synthesis. *Annu. Rev. Biochem.* **72**: 137–174.
- Iuliano, L., F. Micheletta, S. Natoli, S. Ginanni Corradini, M. Iappelli, W. Elisei, L. Giovannelli, F. Violi, and U. Diczfalusy. 2003. Measurement of oxysterols and alpha-tocopherol in plasma and tissue samples as indices of oxidant stress status. *Anal. Biochem.* **312**: 217–223.
- Fahy, E., S. Subramaniam, H. A. Brown, C. K. Glass, A. H. Merrill, Jr., R. C. Murphy, C. R. Raetz, D. W. Russell, Y. Seyama, W. Shaw, et al. 2005. A comprehensive classification system for lipids. *J. Lipid Res.* **46**: 839–861.
- Fakheri, R. J., and N. B. Javitt. 2012. 27-Hydroxycholesterol, does it exist? On the nomenclature and stereochemistry of 26-hydroxylated sterols. *Steroids*. **77**: 575–577.
- Babiker, A., S. Dzeletovic, B. Wiklund, N. Pettersson, J. Salonen, K. Nyssonson, M. Eriksson, U. Diczfalusy, and I. Björkhem. 2005. Patients with atherosclerosis may have increased circulating levels of 27-hydroxycholesterol and cholestenic acid. *Scand. J. Clin. Lab. Invest.* **65**: 365–375.
- Wuolikainen, A., J. Acimovic, A. Lovgren-Sandblom, P. Parini, P. M. Andersen, and I. Björkhem. 2014. Cholesterol, oxysterol, triglyceride, and coenzyme Q homeostasis in ALS. Evidence against the hypothesis that elevated 27-hydroxycholesterol is a pathogenic factor. *PLoS One*. **9**: e113619.
- Ferdinandusse, S., S. Denis, P. L. Faust, and R. J. Wanders. 2009. Bile acids: the role of peroxisomes. *J. Lipid Res.* **50**: 2139–2147.
- Russell, D. W., R. W. Halford, D. M. Ramirez, R. Shah, and T. Kotti. 2009. Cholesterol 24-hydroxylase: an enzyme of cholesterol turnover in the brain. *Annu. Rev. Biochem.* **78**: 1017–1040.
- Björkhem, I. 2007. Rediscovery of cerebrosterol. *Lipids*. **42**: 5–14.

34. Leoni, V., T. Masterman, F. S. Mousavi, B. Wretling, L. O. Wahlund, U. Diczfalusy, J. Hillert, and I. Björkhem. 2004. Diagnostic use of cerebral and extracerebral oxysterols. *Clin. Chem. Lab. Med.* **42**: 186–191.
35. Björkhem, I. 2013. Five decades with oxysterols. *Biochimie*. **95**: 448–454.
36. Dzeletovic, S., O. Breuer, E. Lund, and U. Diczfalusy. 1995. Determination of cholesterol oxidation products in human plasma by isotope dilution-mass spectrometry. *Anal. Biochem.* **225**: 73–80.
37. Bretillon, L., A. Siden, L. O. Wahlund, D. Lutjohann, L. Minthon, M. Crisby, J. Hillert, C. G. Groth, U. Diczfalusy, and I. Björkhem. 2000. Plasma levels of 24S-hydroxycholesterol in patients with neurological diseases. *Neurosci. Lett.* **293**: 87–90.
38. Andersson, S., N. Gustafsson, M. Warner, and J. A. Gustafsson. 2005. Inactivation of liver X receptor beta leads to adult-onset motor neuron degeneration in male mice. *Proc. Natl. Acad. Sci. USA*. **102**: 3857–3862.
39. Kim, H. J., X. Fan, C. Gabbi, K. Yakimchuk, P. Parini, M. Warner, and J. A. Gustafsson. 2008. Liver X receptor beta (LXRbeta): a link between beta-sitosterol and amyotrophic lateral sclerosis-Parkinson's dementia. *Proc. Natl. Acad. Sci. USA*. **105**: 2094–2099.
40. Bigini, P., K. R. Steffensen, A. Ferrario, L. Diomedea, G. Ferrara, S. Barbera, S. Salzano, E. Fumagalli, P. Ghezzi, T. Mennini, et al. 2010. Neuropathologic and biochemical changes during disease progression in liver X receptor beta^{-/-} mice, a model of adult neuron disease. *J. Neuropathol. Exp. Neurol.* **69**: 593–605.
41. Saeed, A. A., G. Genove, T. Li, D. Lutjohann, M. Olin, N. Mast, I. A. Pikuleva, P. Crick, Y. Wang, W. Griffiths, et al. 2014. Effects of a disrupted blood-brain barrier on cholesterol homeostasis in the brain. *J. Biol. Chem.* **289**: 23712–23722.
42. Heverin, M., S. Meaney, D. Lutjohann, U. Diczfalusy, J. Wahren, and I. Björkhem. 2005. Crossing the barrier: net flux of 27-hydroxycholesterol into the human brain. *J. Lipid Res.* **46**: 1047–1052.
43. Meaney, S., M. Heverin, U. Panzenboeck, L. Ekstrom, M. Axelsson, U. Andersson, U. Diczfalusy, I. Pikuleva, J. Wahren, W. Sattler, et al. 2007. Novel route for elimination of brain oxysterols across the blood-brain barrier: conversion into 7alpha-hydroxy-3-oxo-4-cholestenic acid. *J. Lipid Res.* **48**: 944–951.
44. Iuliano, L., P. J. Crick, C. Zerbinati, L. Tritapepe, J. Abdel-Khalik, M. Poirot, Y. Wang, and W. J. Griffiths. 2015. Cholesterol metabolites exported from human brain. *Steroids*. **99**: 189–193.
45. Ogunbare, M., S. Theofilopoulos, A. Lockhart, L. J. Hall, E. Arenas, J. Sjoval, A. G. Brenton, Y. Wang, and W. J. Griffiths. 2010. Cerebrospinal fluid steroidomics: are bioactive bile acids present in brain? *J. Biol. Chem.* **285**: 4666–4679.
46. Zelcer, N., and P. Tontonoz. 2006. Liver X receptors as integrators of metabolic and inflammatory signaling. *J. Clin. Invest.* **116**: 607–614.
47. Evans, M. C., Y. Couch, N. Sibson, and M. R. Turner. 2013. Inflammation and neurovascular changes in amyotrophic lateral sclerosis. *Mol. Cell. Neurosci.* **53**: 34–41.
48. Theofilopoulos, S., W. J. Griffiths, P. J. Crick, S. Yang, A. Meljon, M. Ogunbare, S. S. Kitambi, A. Lockhart, K. Tuschl, P. T. Clayton, et al. 2014. Cholestenic acids regulate motor neuron survival via liver X receptors. *J. Clin. Invest.* **124**: 4829–4842.
49. Zhang, J., Y. Akwa, M. el-Etr, E. E. Baulieu, and J. Sjoval. 1997. Metabolism of 27-, 25- and 24-hydroxycholesterol in rat glial cells and neurons. *Biochem. J.* **322**: 175–184.
50. Clayton, P. T. 2011. Disorders of bile acid synthesis. *J. Inherit. Metab. Dis.* **34**: 593–604.
51. Reboldi, A., E. V. Dang, J. G. McDonald, G. Liang, D. W. Russell, and J. G. Cyster. 2014. Inflammation. 25-Hydroxycholesterol suppresses interleukin-1-driven inflammation downstream of type I interferon. *Science*. **345**: 679–684.
52. Wang, Y., S. Muneton, J. Sjoval, J. N. Jovanovic, and W. J. Griffiths. 2008. The effect of 24S-hydroxycholesterol on cholesterol homeostasis in neurons: quantitative changes to the cortical neuron proteome. *J. Proteome Res.* **7**: 1606–1614.
53. Pfrieger, F. W., and N. Ungerer. 2011. Cholesterol metabolism in neurons and astrocytes. *Prog. Lipid Res.* **50**: 357–371.