

Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in:

Sports Medicine

Cronfa URL for this paper:

<http://cronfa.swan.ac.uk/Record/cronfa34215>

Paper:

Eddolls, W., McNarry, M., Stratton, G., Winn, C. & Mackintosh, K. (2017). High-Intensity Interval Training Interventions in Children and Adolescents: A Systematic Review. *Sports Medicine*

<http://dx.doi.org/10.1007/s40279-017-0753-8>

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/iss/researchsupport/cronfa-support/>

SYSTEMATIC REVIEW

High-Intensity Interval Training Interventions in Children and Adolescents: A Systematic Review

William T. B. Eddolls¹  · Melitta A. McNarry¹  · Gareth Stratton^{1,2}  · Charles O. N. Winn¹  · Kelly A. Mackintosh¹ 

© The Author(s) 2017. This article is an open access publication

Abstract

Background Whilst there is increasing interest in the efficacy of high-intensity interval training in children and adolescents as a time-effective method of eliciting health benefits, there remains little consensus within the literature regarding the most effective means for delivering a high-intensity interval training intervention. Given the global health issues surrounding childhood obesity and associated health implications, the identification of effective intervention strategies is imperative.

Objectives The aim of this review was to examine high-intensity interval training as a means of influencing key health parameters and to elucidate the most effective high-intensity interval training protocol.

Methods Studies were included if they: (1) studied healthy children and/or adolescents (aged 5–18 years); (2) prescribed an intervention that was deemed high intensity; and (3) reported health-related outcome measures.

Results A total of 2092 studies were initially retrieved from four databases. Studies that were deemed to meet the criteria were downloaded in their entirety and independently assessed for relevance by two authors using the pre-

determined criteria. From this, 13 studies were deemed suitable. This review found that high-intensity interval training in children and adolescents is a time-effective method of improving cardiovascular disease biomarkers, but evidence regarding other health-related measures is more equivocal. Running-based sessions, at an intensity of >90% heart rate maximum/100–130% maximal aerobic velocity, two to three times a week and with a minimum intervention duration of 7 weeks, elicit the greatest improvements in participant health.

Conclusion While high-intensity interval training improves cardiovascular disease biomarkers, and the evidence supports the effectiveness of running-based sessions, as outlined above, further recommendations as to optimal exercise duration and rest intervals remain ambiguous owing to the paucity of literature and the methodological limitations of studies presently available.

Key Points

High-intensity interval training can improve certain cardiovascular health parameters in children and adolescents.

Evidence supporting the overall effectiveness of high-intensity interval training as a means of eliciting improvements to other health outcomes, specifically body composition and blood pressure, remains unclear.

While this review enables the establishment of suggested guidelines for high-intensity interval training protocols, recommendations for some protocol details remain unclear.

✉ William T. B. Eddolls
636442@swansea.ac.uk

¹ Applied Sports Science Technology and Medicine Research Centre (A-STEM), College of Engineering, Swansea University, A101 Engineering East, Bay Campus, Fabian Way, Swansea SA1 8EN, UK

² School of Sport Health and Exercise Science, University of Western Australia, Perth, WA, Australia

1 Introduction

Worldwide, the prevalence of childhood and adolescent obesity has reached unparalleled levels [1, 2]. Specifically, in the UK, approximately 28% of children are classified as overweight or obese [2, 3], representing a significant burden on health services. Indeed, current estimates suggest that obesity and its deleterious health consequences, such as type 2 diabetes mellitus [4] and coronary heart disease [5], cost the National Health Service £5.1 billion per year [6]. Whilst some reports suggest a plateau in paediatric obesity over the last decade [7], others suggest that UK obesity levels have increased between 1980 and 2014 by 48 and 39% in boys and girls, respectively [2]. Of concern, paediatric obesity has been associated with an increased prevalence of cardiometabolic risk factors [8], which have been shown to track into later life [9] and increase the risk of premature mortality [9, 10]. Although the causes of obesity and cardiometabolic risk are multifaceted, low physical activity levels, as well as high engagement in sedentary pursuits, have been identified as key contributory factors [9, 11–13]. Specifically, according to the latest statistics, in England, only 21% of boys and 16% of girls meet UK physical activity guidelines of at least 60 min of moderate-to-vigorous physical activity every day [3], with physical activity levels further declining with age [14–16]. Effective interventions targeted at increasing youth physical activity levels are therefore imperative.

While traditional interventions designed to increase physical activity and improve health have principally used moderate-intensity continuous exercise [17–19], the relevance of such programmes to the sporadic high-intensity nature of children's habitual play patterns has been questioned [20]. Consequently, high-intensity interval-based programmes have recently been investigated as a potentially potent and time-efficient form of physical activity and health promotion [21]. Indeed, whilst exercise, a sub-component of physical activity, is structured and conducted for the health-associated benefits, it could provide a necessary mediatory step to provoke positive long-term behavioural change. However, to date, studies have demonstrated varied success at eliciting significant improvements [22–25], potentially owing to a lack of consensus regarding an optimum high-intensity interval training (HIIT) intervention protocol with regard to exercise intensity, frequency and duration. Nonetheless, recent systematic reviews [26, 27] highlighted that HIIT can elicit greater improvements in health-related parameters (i.e. cardiometabolic health and body composition) in adolescents, compared with traditional programmes [23, 28–30]. However, no systematic reviews have investigated the effects of HIIT on primary school-aged children, the

identified potential differences between pre-pubertal and pubertal youth in the adaptations elicited or provided optimal HIIT protocol recommendations. Therefore, the purpose of this review was to systematically synthesise the scientific literature regarding HIIT on improving body composition, cardiometabolic health and cardiovascular health in children and adolescents and to establish an optimal HIIT protocol with regard to session structure, intensity, frequency and duration.

2 Methodology

In line with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses [31], the following methodology details the review's inclusion criteria, search strategy, data collection and study analysis protocols.

2.1 Inclusion Criteria

2.1.1 Types of Study

The present review included studies involving interventions targeted at reducing obesity-related physiological parameters with a principal focus on one of the following: high-intensity physical activity, high-intensity exercise/training or high-intensity intermittent/interval exercise/training interventions.

2.1.2 Types of Participant

Studies incorporating children and adolescents between the ages of 5 and 18 years were included. These age constraints were applied to all children and adolescents without disability, irrespective of whether they were a healthy weight, overweight or obese. Studies that used physical activity interventions as part of a treatment for specific illnesses were excluded.

2.1.3 Intervention Variables and Outcome Measures

To be included in the review, studies were required to report a minimum of one intervention exercise session intensity variable and one outcome measure, measured at baseline and post-intervention and compared with either a moderate-intensity exercise intervention or control group.

2.1.4 Intervention Intensity Variables

Interventions were defined as high intensity if: (1) the intensity was $\geq 90\%$ peak oxygen uptake [32]; (2) had an intensity that was $\geq 100\%$ maximal aerobic speed [33]; and/or (3) ensured that the participant's heart rate was

$\geq 90\%$ of their peak heart rate [34, 35]. There were no restrictions applied regarding the duration of the intervention.

2.1.5 Primary Outcomes

Primary outcomes included cardiometabolic health markers, namely mean systolic (SBP) and diastolic (DBP) blood pressure, body composition in the form of body mass index (BMI), body fat percentage (BF%) and fat-free mass, and cardiovascular disease (CVD) biomarker analysis including at least one of the following: glucose, insulin, triglyceride and total cholesterol, as well as its sub-fractions, low-density lipoprotein cholesterol and high-density lipoprotein cholesterol.

2.2 Search Strategy

Electronic databases were searched until September 2016, with no restriction set on the publication year. The PubMed and SCOPUS databases were explored using the following keyword search strategy, devised by the research team and verified by a subject librarian: (high intensity training OR high intensity exercise* OR high intensity activit* OR high intensity intermittent training OR intensity intermittent exercise* OR high intensity intermittent activit* or high intensity interval training OR high intensity interval exercise* OR high intensity interval activit*) AND (child* OR children OR pediatric OR paediatric OR adolescen* OR juvenile*) AND (health OR healthy). Inclusion of at least one of the keywords was required in the study title for it to be considered. Studies were excluded based on language; only studies written in English were included. Additional studies were identified by searching the reference lists of included studies. Google Scholar and ResearchGate were also searched to identify studies that were potentially overlooked by the database searches.

2.3 Data Collection and Analysis

Pertinent study abstracts from the stated search strategies were downloaded and independently screened. Studies that were deemed to meet the criteria were downloaded in their entirety and independently assessed for relevance by two authors using the pre-determined criteria. When study information was missing, the research team attempted to contact the primary author of the incomplete study. If the author failed to respond, the study was excluded.

2.4 Effect Size

Cohen's d was used to determine the standardised mean effect of HIIT on the previously outlined health-related

outcome measures compared with a control group or a moderate- or light-intensity group [36]. Confidence intervals (CIs) were calculated by applying an equation recommended by Nakagawa and Cuthill [37], employing standard error calculations [38]. For studies that provided values for both moderate and control groups, moderate group values were included as the comparison. Additionally, effect size was not calculated for studies that failed to disclose post-intervention mean values.

2.5 Risk of Bias Assessment

Risk of bias was assessed independently by two reviewers using the Cochrane risk of bias tool (RoB 2.0) [39]. Using the RoB 2.0 tool, studies were awarded an overall risk of bias grade of either high, some or low risk of bias. This overall grade was calculated by assessing five domains: (A) bias arising from the randomisation process; (B) bias owing to deviations from intended interventions; (C) bias owing to missing outcome data; (D) bias in measurement of the outcome; and (E) bias in selection of the reported result.

2.6 Heterogeneity Assessment

Because of the variation of the study characteristics in this review, for example between interventions, outcome measures and cohort populations, it was deemed unsuitable to amalgamate the results for a meta-analysis. Therefore, the results in this review were analysed narratively.


3 Results

The database search generated 2092 studies. Once duplicates were removed, 54 title/abstracts were screened for eligibility, with the reference list search producing three further studies. From this, 13 studies were deemed suitable. The screening process is shown in Fig. 1. In total, 7 of the 13 studies included in this review examined the effects of HIIT interventions on pre-pubertal participants, the characteristics of which are summarised in Table 1, with the remaining six studies examining the effects of HIIT interventions on pubertal participants (Table 2).

3.1 Risk of Bias

The methodological rigour of studies included in this review, according to the risk of bias assessment, is presented in Table 3. Seven studies were considered to have a high risk of bias [22, 24, 25, 29, 40, 41, 45], whereas only two [28, 43] and four [23, 30, 42, 44] studies were considered to have some or low risk of bias, respectively. In

Fig. 1 Phases of study selection during data collection



studies deemed to have a high or some risk of bias [22, 24, 25, 28, 29, 40, 41, 43, 45], the bias arose from the randomisation process (domain A).

3.2 Body Mass and Composition

All 13 included studies reported the effect of HIIT compared with moderate-intensity exercise or a control group on BMI ($n = 9$), BF% ($n = 9$) or fat-free mass ($n = 1$). The results revealed little evidence to suggest that HIIT can elicit significant changes in body composition (Table 4), although Tjønna et al. [23] and Racil et al. [30], both of which were assessed to be at low risk of bias, reported significant improvements in BMI and BF% associated with a medium-to-large effect sizes following a 3-month intervention.

It is pertinent to note that whilst other studies failed to find a significant improvement in measures of body mass or composition, there was a general trend for a greater change in body mass and composition in the HIIT group [22, 28, 29, 43, 45]. The exception to this was Baquet et al. [25], who saw significant increases to BMI and BF% for both the HIIT protocol and the control group; however, this study was deemed to have a high risk of bias. While no significant benefits were reported in pre-pubertal children, significant improvements in body mass and composition

have been demonstrated in pubertal children [23, 30], suggesting a potential maturational effect. There were insufficient data to examine potential sex differences regarding the efficacy of HIIT in eliciting significant changes in body composition.

3.3 Cardiovascular Health

3.3.1 Blood Pressure

Five studies investigated the effect of HIIT on SBP and DBP, with the majority concluding significant benefits were obtained (Table 5). Specifically, two studies that were assessed to be of low risk of bias, Tjønna et al. [23] and Racil et al. [30], reported significant improvements in both SBP and DBP following the intervention, although it is pertinent to note the low effect sizes associated with the improvements reported for SBP and DBP in Racil et al. [30] (Table 5). Whilst the remaining studies [28, 29, 44] reported no significant differences, they demonstrated a trend towards a lower SBP and DBP. Interestingly, Boddy et al. [44], demonstrating methodological rigour through a low risk of bias, found an increase in both DBP and SBP in the HIIT group and reductions in the control group, though not significant. Regarding maturation differences, no studies examining pre-pubertal children reported

Table 1 Characteristics of studies examining pre-pubertal participants

References	Sample population	Maturation	INT duration (wk)	INT type	Group size (n)	Modality/intensity	Repeated bouts/frequency	Exercise bout/recovery duration	Protocol duration (including recovery)	Total exercise INT duration
Baquet et al. [22]	Pre-pubertal primary school children; N = 53 (23 boys; 8–11 years)	Maturation measured, but not reported	7	HIT protocol	33	Shuttle runs (100–130 % MAS)	Bouts: 5–10 Sets: 1–4	10–20 s/ 10–20 s	30 min	7 h
Baquet et al. [40]	Primary school children; N = 77 (43 boys; 9.6 ± 1.0 years)	Stage 1 = 40 boys, 29 girls Stage 3 = 3 boys, 5 girls combined <3	7	HIT protocol	20	Shuttle runs (100–130 % MAS)	Bouts: 5–10 Sets: 1–4	3-min rest between set (2 times weekly)	25–35 min	10 h and 30 min
Baquet et al. [41]	Pre-pubertal children; N = 100 (46 boys; 9.7 ± 0.8 years)	Stage 1 = 46 boys, 25 girls Stage 3 = 29 girls combined <3	7	HIT protocol	22	Shuttle runs (100–130 % MAS)	Bouts: 5–10 Sets: 1–4	3-min rest between set (3 times weekly)	18–39 min	7 h and 21 min
Lambbrick et al. [42]	Obese and normal weight children; N = 55 (32 boys)	Peak height velocity at baseline: INT = 2.7 ± 0.1 CON = 2.65 ± 0.05	6	HIT protocol	22	Shuttle runs (80–85 % MAS)	Bouts: 1–4 (3 times weekly)	6–18 min/ 5 min	30 min	7 h
Lau et al. [24]	Overweight primary school children; N = 48 (36 boys; 10.4 ± 0.9 years)	Maturation not reported	6	HIT protocol	19	Shuttle runs (100–130 % MAS)	Bouts: 5–10 Sets: 1–4	10–20 s/ 10–20 s	60 min	12 h
Nourry et al. [43]	Pre-pubertal children; N = 18 (11 boys; 10.0 ± 0.8 years)	Stage 1 = 18	8	LIT protocol	53	Child specific games (93% mean HR _{max})	Bouts: 7 (2 times weekly)	6 min/2 min	60 min	12 h
Rosenkranz et al. [28]	Pre-pubertal children; N = 16 (2 boys; 7–12 years)	Stage 1 = 16	8	HIT protocol	27	Shuttle runs (120 % MAS)	Bouts: 12 (3 times weekly)	15 s/15 s	6 min	1 h and 48 min
			8	LIT protocol	21	Shuttle runs (100 % MAS)	Bouts: 16 (3 times weekly)	15 s/15 s	8 min	2 h and 24 min
			8	HIT protocol	12	Shuttle runs (100–130 % MAS)	Bouts: 10 Sets: 4 times	10–20 s/ 10–20 s	30 min	8 h
			9	Control	9	Shuttle runs (100–130 % MAS)	Bouts: 5–10 Sets: 4 times	(2 times weekly)	30 min	8 h
			8	HIT protocol	8	Shuttle runs (100–130 % MAS)	Bouts: 5–10 Sets: 4 times	10–20 s/ 10–20 s	30 min	8 h
			8	Control	8					

HIT high-intensity interval training, HR_{max} heart rate maximum, INT intervention, LIT light-intensity training, MAS maximal aerobic speed

Table 2 Characteristics of studies examining pubertal participants

References	Sample population	Maturation	INT duration (wk)	INT type	Group size (n)	Modality/ intensity	Repeated bouts/ frequency	Exercise bout/ recovery duration	Protocol duration (including recovery)	Total exercise INT duration
Baquet et al. [25]	Secondary school children; N = 551 (290 boys; 11–16 years)	Not reported	10	HIIT protocol	503	Shuttle runs (100–120 % MAS)	Bouts: 10 Sets: 3 3-min rest between set (1 times weekly)	10 s/10 s	60 min	10 h
Boddy et al. [44]	Secondary school girls; N = 16 (11.8 ± 0.3 years)	Peak height velocity at baseline: CON = 0.028 ± 0.427	3	Control HIIT protocol	48	Dance class (>93.4 mean % HR _{max})	Bouts: 6 (4 times weekly)	30 s/45 s	20 min	4 h
Racil et al. [30]	Obese female adolescents; N = 47 (14.2 ± 1.2 years)	Not reported	12	HIIT protocol	17	Various types of interval training (100% MAS)	Bouts: 8–16 (3 times weekly)	15 s/15 s	4–8 min	3 h and 36 min
Sperlich et al. [45]	Male soccer players; N = 19 (13.5 ± 0.4 years)	Maturation not reported	5	Control HIIT protocol	16	Various types of interval training (80% MAS)	Bouts: 8–16 (3 times weekly)	15 s/15 s	4–8 min	3 h and 36 min
Tjonna et al. [23]	Overweight and obese adolescents; N = 54 (26 boys; 14.0 ± 0.3 years)	Maturation not reported	12	HIIT protocol	14	Various types of interval training (90–95 % HR _{max})	Bouts: 4–12 (3–4 times weekly)	30 s to 4 min/30 s to	30 min	8 h and 45 min
Weston et al. [29]	Adolescent secondary school children; N = 101 (62 boys; 14.0 ± 0.3 years)	Peak height velocity at baseline: INT = 0.3 ± 1.0	10	Control HIIT protocol	28	Various types of interval training (50–70 % HR _{max}) Treadmill walking/ running (90–95 % HR _{max})	Bouts: 1–5 (3–4 times weekly) Bouts: 4 (2 times weekly)	10–30 min/1–3 min 4 min/3 min	40–60 min	14 h and 35 min
		Control = 0.5 ± 1.3		Control	26	Games based (>90% mean HR _{max})	Bouts: 4–7 (3 times weekly)	45 s/90 s	7 min 30 s to 14 min 15 s	5 h and 30 min

HIIT high-intensity interval training, HR_{max} heart rate maximum, INT intervention, MAS maximal aerobic speed

Table 3 Risk of bias assessment

References	Domain A	Domain B	Domain C	Domain D	Domain E	Total
Baquet et al. [22]	High	Low	Low	Low	Low	High
Baquet et al. [25]	High	Low	Low	Low	Low	High
Baquet et al. [40]	High	Low	Low	Low	Low	High
Baquet et al. [41]	High	Low	Low	Low	Low	High
Boddy et al. [44]	Low	Low	Low	Low	Low	Low
Lambrick et al. [42]	Low	Low	Low	Low	Low	Low
Lau et al. [24]	High	Low	Low	Low	Low	High
Nourry et al. [43]	Some	Low	Low	Low	Low	Some
Racil et al. [30]	Low	Low	Low	Low	Low	Low
Rosenkranz et al. [28]	Some	Low	Low	Low	Low	Some
Sperlich et al. [45]	High	Low	Low	Low	Low	High
Tjønna et al. [23]	Low	Low	Low	Low	Low	Low
Weston et al. [29]	High	Low	Low	Low	Low	High

Domain: (A) bias arising from the randomisation process, (B) bias owing to deviations from intended interventions, (C) bias owing to missing outcome data, (D) bias in measurement of the outcome, (E) bias in selection of the reported result. Total overall risk of bias grade was calculated by assessing the five domains [A–E]

Table 4 Baseline and post-intervention changes to body mass/composition and effect size between high-intensity interval training (HIIT) and control/moderate protocols

References	Outcome measure	HIIT (mean change from baseline)	Control/moderate (mean change from baseline)	Effect size (Cohen's <i>d</i>)	95% CI
Baquet et al. [25]	BMI (kg/m^2)	0.40**	0.60**	-0.14	-0.78 to 0.49
Baquet et al. [40]	BMI (kg/m^2)	0.10	-0.30/0.20	-0.10	-0.78 to 0.58
Boddy et al. [44]	BMI (kg/m^2)	-0.50	0.20	0.93	0.30 to 1.56
Lambrick et al. [42]	BMI (kg/m^2)	0.00	0.00	0.23	-0.40 to 0.87
Lau et al. [24]	BMI (kg/m^2)	0.20	0.10/0.40*	0.42	-0.18 to 1.01
Racil et al. [30]	BMI (kg/m^2)	-3.20*	0.30/-1.70*	-1.41	-3.52 to 0.69
Rosenkranz et al. [28]	BMI (kg/m^2)	-1.40	0.00	-0.06	-0.73 to 0.61
Tjønna et al. [23]	BMI (kg/m^2)	-0.70**	-0.20	-1.50	-2.06 to -0.94
Weston et al. [29]	BMI (kg/m^2)	-0.60 ^a	0.80 ^a	N/A	N/A
Baquet et al. [25]	Body fat (%)	1.60*	1.30*	-0.31	-0.94 to 0.33
Baquet et al. [22]	Body fat (%)	-0.90	-0.70	0.10	-0.51 to 0.71
Baquet et al. [41]	Body fat (%)	0.10	0.10	0.02	-0.66 to 0.69
Boddy et al. [44]	Body fat (%)	0.22	0.46	0.61	0.14 to 1.07
Lambrick et al. [42]	Body fat (%)	-0.10	0.40	0.08	-0.49 to 0.65
Nourry et al. [43]	Body fat (%)	1.50	0.50	0.14	-0.56 to 0.84
Racil et al. [30]	Body fat (%)	-3.90*	-0.50/-3.40*	-0.59	-1.06 to -0.12
Rosenkranz et al. [28]	Body fat (%)	-2.20	-1.00	-0.17	-0.80 to 0.45
Tjønna et al. [23]	Body fat (%)	-0.90**	-0.30	3.00	2.35 to 3.65
Sperlich et al. [45]	FFM (kg)	1.00	0.90	0.71	0.28 to 1.13

BMI body mass index, CI confidence interval, FFM fat-free mass, N/A effect size not calculated because of no reported post-intervention means

* $p < 0.05$, ** $p < 0.01$, significantly different from baseline

^a Values adjusted for sex, baseline value and maturity offset

significant changes, whereas significant improvements were reported in pubertal populations [23, 30]. Sex differences could not be investigated because of insufficient data.

3.3.2 Cardiovascular Disease Biomarker Health

Four studies examined the effect of HIIT on CVD biomarkers, specifically blood glucose ($n = 4$), total

Table 5 Baseline to post-intervention changes in systolic blood pressure/diastolic blood pressure (SBP/DBP) and effect size between high-intensity interval training (HIIT) and control/moderate protocols

References	Outcome measure (mmHg)	HIIT (mean change from baseline)	Control/moderate (mean change from baseline)	Effect size (Cohen's <i>d</i>)	95% CI
Boddy et al. [44]	SBP	10.10	-1.40	0.34	0.08 to 0.61
Racil et al. [30]	SBP	-0.60*	0.00/-0.40*	0.00	-0.26 to 0.26
Rosenkranz et al. [28]	SBP	-2.20	-2.50	-0.54	-0.82 to -0.26
Tjønna et al. [23]	SBP	-9.40**	-2.50*	-2.00	-2.31 to -1.69
Weston et al. [29]	SBP	-5.00 ^a	-1.00 ^a	N/A	N/A
Boddy et al. [44]	DBP	5.90	-4.10	1.14	0.75 to 1.52
Racil et al. [30]	DBP	-6.00*	-1.00/-4.00*	-0.32	-0.64 to 0.01
Rosenkranz et al. [28]	DBP	-2.50	-1.70	-0.83	-1.18 to -0.48
Tjønna et al. [23]	DBP	-5.50**	1.80	-1.50	-1.89 to -1.11
Weston et al. [29]	DBP	-6.00 ^a	-4.00 ^a	N/A	N/A

CI confidence interval, N/A effect size not calculated because of no reported post-intervention means

* $p < 0.05$; ** $p < 0.01$ significantly different from baseline

^a Values adjusted for sex, baseline value and maturity offset

cholesterol ($n = 2$), high-density lipoprotein cholesterol ($n = 3$), low-density lipoprotein cholesterol ($n = 1$), blood triglycerides ($n = 3$) and insulin ($n = 2$). The results, in Table 6, support HIIT as an effective strategy for improving CVD biomarker health. All of the studies demonstrated significant [23, 28, 30] or clinically substantial [29] improvements in multiple CVD biomarker

outcome measures. Tjønna et al. [23] reported greater significant improvements for blood glucose in favour of the HIIT group ($p < 0.01$) compared with the control group ($p < 0.05$), with an effect size indicating large clinically important differences ($d = -1.43$, 95% CI -3.01 to 0.16). Additionally, results from Rosenkranz et al. [28] suggest large significant reductions in total cholesterol compared

Table 6 Baseline and post-intervention changes in cardiovascular disease biomarkers and effect size between high-intensity interval training (HIIT) and control/moderate protocols

References	Outcome measure	HIIT (mean change from baseline)	Control/moderate (mean change from baseline)	Effect size (Cohen's <i>d</i>)	95% CI
Racil et al. [30]	Glucose (mmol•L ⁻¹)	-0.20*	0.00/-0.20*	-0.32	-0.44 to 0.13
Rosenkranz et al. [28]	Glucose (mg/dL)	5.20	0.40	-0.16	-1.80 to 1.17
Tjønna et al. [23]	Glucose (mmol•L ⁻¹)	-0.30**	-0.10	-1.43	-3.01 to 0.16
Weston et al. [29]	Glucose (mmol•L ⁻¹)	-0.10 ^a	-0.03 ^a	N/A	N/A
Rosenkranz et al. [28]	Total cholesterol (mg/dL)	-22.00*	2.40	-0.93	-1.16 to -0.70
Weston et al. [29]	Total cholesterol (mmol•L ⁻¹)	-0.24 ^a	0.00 ^a	N/A	N/A
Rosenkranz et al. [28]	HDL-cholesterol (mg/dL)	9.90	3.60	0.42	0.02 to 0.81
Tjønna et al. [23]	HDL-cholesterol (mmol•L ⁻¹)	0.11*	0.09	0.35	-0.50 to 1.20
Weston et al. [29]	HDL-cholesterol (mmol•L ⁻¹)	-0.14 ^a	-0.24 ^a	N/A	N/A
Rosenkranz et al. [28]	LDL-cholesterol (mg/dL)	-34.80*	-5.60	-1.67	-2.03 to -1.31
Rosenkranz et al. (2012) [28]	Triglycerides (mg/dL)	23.50	3.50	-0.07	-0.33 to 0.19
Tjønna et al. [23]	Triglycerides (mmol•L ⁻¹)	-0.50	-0.10	-0.71	-1.57 to 0.14
Weston et al. [29]	Triglycerides (mmol•L ⁻¹)	-0.05 ^a	0.18 ^a	N/A	N/A
Racil et al. [30]	Insulin (IU mL ⁻¹)	-5.70*	-0.80/-4.30*	-0.82	-1.55 to -0.10
Tjønna et al. [23]	Insulin (pmol/L)	-54.30*	-33.00*	-0.46	-0.70 to -0.22

CI confidence interval, HDL high-density lipoprotein cholesterol, LDL low-density lipoprotein cholesterol, N/A effect size not calculated because of reported post-intervention means

* $p < 0.05$; ** $p < 0.01$ significantly different from baseline

^a Values adjusted for sex, baseline value and maturity offset

with the control group. Further, the associated effect size ($d = -0.93$, 95% CI -1.16 to -0.70) suggested highly clinically important differences. Rosenkranz et al. [28] saw significant reductions ($p < 0.05$) in low-density lipoprotein cholesterol as a result of HIIT, with a large and clinically important difference ($d = -1.67$, 95% CI -2.03 to -1.31) compared with a small non-significant increase in the control group. Examining blood insulin results in Tjønna et al. [23], larger significant reductions in the HIIT group were found compared with the control group, with further significant reductions after a 12-month follow-up. In addition, high-density lipoprotein cholesterol significantly improved in the HIIT group compared with a non-significant increase in the control group. Furthermore, Racil et al. [30] reported significant reductions in blood glucose and insulin in both HIIT and moderate-intensity protocols compared with a control group, with a greater improvement reported in the HIIT group for both measures. Effect sizes for both blood glucose and insulin were low to moderate ($d = 0.32$, 95% CI -0.44 to 0.13) and large ($d = -0.82$, 95% CI -1.55 to -0.10), respectively.

Notwithstanding the limited improvement in blood triglycerides, Weston et al. [29] reported clinically substantial beneficial effects as a result of HIIT despite increased triglyceride levels reported in the control group. Despite these encouraging findings, it is pertinent to note that the studies by Rosenkranz et al. [28] and Weston et al. [29] were assessed to have some and a high risk of bias, respectively; therefore, caution should be taken when interpreting these studies as methodological limitations may have confounded the results. There was no effect owing to maturation on CVD biomarkers, with significant or clinically substantial improvements found in both pre-pubertal [28] and pubertal [23, 29, 30] children. Sex differences were not reported and, therefore, their effect is unknown.

3.4 High-Intensity Interval Training Intervention Protocol

All studies included in this review provided a detailed description of their intervention protocol in terms of session structure, duration, intensity and frequency, in addition to intervention duration; key details of these are summarised in Tables 1 and 2. Based on the four studies that demonstrated significant health improvements [23, 28–30], a running-based HIIT intervention at an intensity of $>90\%$ heart rate maximum/100–130 % maximal aerobic velocity, two to three times a week with a minimum intervention duration lasting 7 weeks could be considered the suggested practice. However, suggested exercise session duration and rest intervals remain ambiguous owing to the variance across the studies, a notion also supported by Baquet et al. [46].

4 Discussion

The aim of the current review was to synthesise previous literature that examined HIIT in children and adolescents and establish its potential effect on body composition, cardiometabolic health and cardiovascular health. In addition, this review aimed to identify an optimal HIIT protocol with regard to session structure, intensity, frequency and duration. In accord with this aim, 13 studies were evaluated providing evidence suggesting that HIIT can significantly improve certain health parameters in children and adolescents. However, evidence supporting the overall effectiveness of HIIT as a means of eliciting improvements to all the specified health outcomes remains unclear. Some guidelines for a HIIT protocol were established, though recommendations for certain protocol details remain unidentified.

Advancing previous reviews [26, 27], the findings of the current review suggest that pubertal children may achieve a greater benefit as a result of HIIT when compared with pre-pubertal children, a topic that has been widely debated [47–50]. However, it is pertinent to note that this may be a consequence of several methodological factors that limit the interpretation of previous studies. First, the duration of the HIIT interventions examining pubertal participants tended to be longer than those in pre-pubertal children. Given the present findings suggesting that a minimum of 7 weeks is required for significant adaptations to be manifest, this shorter intervention duration may lead to erroneous conclusions regarding the efficacy of HIIT in this population.

Furthermore, differences in the participant characteristics between pre-pubertal and pubertal studies with regard to baseline body mass or body composition and health status may confound interpretation of inter-study differences and their attribution to maturity per se. Specifically, the majority of studies in pre-pubertal children used those of a normal weight compared with the inclusion of overweight or obese participants in pubertal studies, which may predispose these latter studies to demonstrating greater health benefits, irrespective of their biological age. Moreover, both Tjønna et al. [23] and Racil et al. [30] did not report maturation stages, subsequently casting ambiguity over the cohort's true maturational stage. Despite this, when focusing on the additional study involving pubertal children [29], it generally elicited greater improvements in outcome measures when compared with the study involving pre-pubertal children [28]. Moreover, only two of the studies involving pubertal children that demonstrated positive significant results [23, 30] considered dietary intake; a failure to account for changes in dietary intake, which is strongly associated with cardiometabolic health [51] and

obesity [52] in children and adolescents, confounds the interpretation of the results and their attribution to the exercise per se. Finally, in addition to the aforementioned methodological limitations, the interpretation of the overall findings of this study may also be limited by a mixed risk of bias between the significant studies.

An additional factor that may contribute to the collective ambiguity regarding the overall effectiveness of HIIT is the ‘compensation effect’. Specifically, a recent study has suggested that school children appear to compensate for increased physical activity levels, with a reduction in physical activity undertaken the following day [53]. Additionally, the ‘activitystat’ hypothesis suggests that increased levels of physical activity during one part of the day may result in a compensatory decrease in physical activity in another part [54]. Therefore, there is scope to suggest that prescribing exercise that reflects the characteristics of children’s comparatively elevated levels of habitual play [55] may result in a decrease in habitual physical activity levels that day or the subsequent day. This therefore highlights the need to measure habitual physical activity, as was the case for only four of the studies included within this review [23, 28, 29, 44], alongside the previously outlined study outcomes. Regarding the effect of HIIT between sexes, no conclusions could be drawn as none of the studies included in this review provided a breakdown of between sex differences. This could have implications for future research, given the possibility that effects of exercise interventions on body size [56, 57], cardiorespiratory health [58] and cardiometabolic health [57] may be sex dependent in children. Therefore, future studies should endeavour to report a more expansive breakdown of results, thus providing clarification as to possible sex and maturational differences associated with HIIT.

Whilst the studies within this review have advanced our understanding regarding the influence of HIIT in children under laboratory-based conditions, the relatively small sample sizes and intervention delivery methods highlight potential issues regarding larger scale implementation of HIIT. Schools have frequently been used as a foundation in the implementation of physical activity interventions [59, 60] because of their access to a greater population of children, who spend 40% of their waking hours there [61], and are widely accepted as one of the most effective locations to promote physical activity and health. Furthermore, previous studies have demonstrated how short-term HIIT interventions have been successfully embedded within the school timetable [44, 62]. It has been suggested that HIIT can allow for greater class control compared with conventional physical education lessons, and can be adapted to include specific movements related to different

sports [63]. Therefore, given the promising findings regarding the effectiveness of HIIT, future research may wish to consider how long-term HIIT interventions could be incorporated within the school environment. A key consideration in the development of future interventions, and participants’ engagement in and adherence to the devised program, is a participant’s perceived enjoyment. Whilst HIIT has been suggested to be a preferable exercise modality to more conventional aerobic exercise [26], further research is required, with only one study in the present review considering this aspect of intervention development and implementation [42]. Finally, the sustained post-intervention efficacy of HIIT interventions that reported significant improvements to body composition and cardiometabolic and cardiovascular health in children and adolescents remains indeterminate owing to a predominant scarcity of studies reporting a post-intervention follow-up. Therefore, future studies should incorporate a follow-up period within their study design to assess the long-term post-intervention sustainability of positive HIIT elicited benefits.

5 Conclusion

High-intensity interval training is a time-effective method of improving CVD biomarker health in children and adolescents. However, evidence supporting its effectiveness in additional health measures remains equivocal. This review suggests that running-based sessions, at an intensity of >90% maximum heart rate/100–130% maximal aerobic velocity, two to three times a week and with a minimum intervention duration lasting >7 weeks, elicit improvements in health markers; however, these findings are limited by the mixed risk of bias between the significant studies. Further recommendations as to exercise duration and rest intervals remain ambiguous owing to the paucity and methodological limitations of studies presently available.

Compliance with Ethical Standards

Funding William Eddolls, Melitta McNarry and Kelly Mackintosh were supported by Commando Joe’s Fun and Fitness Ltd. The funders had no role in the study design, collection, analysis and interpretation of data, writing of the manuscript or submission of the article for publication.

Conflict of interest William Eddolls, Melitta McNarry, Gareth Stratton, Charles Winn and Kelly Mackintosh have no conflicts of interest directly relevant to the content of this review.

Ethics approval Approval to conduct this research was granted by Swansea University A-STEM Ethical Advisory Committee.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Lobstein T, Jackson-Leach R, Moodie ML, et al. Child and adolescent obesity: part of a bigger picture. *Lancet*. 2015;385:2510–20.
- Ng M, Fleming T, Robinson M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the global burden of disease study 2013. *Lancet*. 2014;384:766–81.
- Ryley A. Children's BMI, overweight and obesity. In: Mindell CR, editor. Health survey for England - 2012. London: NHS England; 2013.
- American Diabetes Association. Type 2 diabetes in children and adolescents. *Pediatrics*. 2000;105:671–80.
- American Heart Association. Obesity and heart disease. *Circulation*. 1997;96:3248–50.
- Scarborough P, Bhatnagar P, Wickramasinghe KK, et al. The economic burden of ill health due to diet, physical inactivity, smoking, alcohol and obesity in the UK: an update to 2006–07 NHS costs. *J Public Health (Bangkok)*. 2011;33:527–35.
- Rokholm B, Baker JL, Sørensen TIA. The levelling off of the obesity epidemic since the year 1999: a review of evidence and perspectives. *Obes Rev*. 2010;11:835–46.
- Skinner AC, Perrin EM, Moss LA, et al. Cardiometabolic risks and severity of obesity in children and young adults. *N Engl J Med*. 2015;373:1307–17.
- Freedman DS, Mei Z, Srinivasan SR, et al. Cardiovascular risk factors and excess adiposity among overweight children and adolescents: the Bogalusa Heart Study. *J Pediatr*. 2007;150(12–17):e2.
- Franks PW, Hanson RL, Knowler WC, et al. Childhood obesity, other cardiovascular risk factors and premature death. *N Engl J Med*. 2010;362:485–93.
- Janssen I, Katzmarzyk PT, Boyce WF, et al. Comparison of overweight and obesity prevalence in school-aged youth from 34 countries and their relationships with physical activity and dietary patterns. *Obes Rev*. 2005;6:123–32.
- Trost SG, Kerr LM, Ward DS, et al. Physical activity and determinants of physical activity in obese and non-obese children. *Int J Obes Relat Metab Disord*. 2001;25:822–9.
- Kopelman PG. Obesity as a medical problem. *Nature*. 2000;404:635–43.
- Sallis JF, Prochaska JJ, Taylor WC. A review of correlates of physical activity of children and adolescents. *Med Sci Sports Exerc*. 2000;32:963–75.
- Heitzler C, Lytle L, Erickson D, et al. Physical activity and sedentary activity patterns among children and adolescents: a latent class analysis approach. *J Phys Act Health*. 2011;8:457–67.
- Troiano RP, Berrigan D, Dodd KW, et al. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc*. 2008;40:181–8.
- Mcmurray RG, Harrell JS, Bangdiwala SI, et al. A school-based intervention can reduce body fat and blood pressure in young adolescents. *J Adolesc Health*. 2002;31:125–32.
- Hansen HS, Froberg K, Hyldebrandt N, et al. A controlled study of eight months of physical training and reduction of blood pressure in children: the Odense schoolchild study. *BMJ*. 1991;303:682–5.
- Meyer AA, Kundt G, Lenschow U, et al. Improvement of early vascular changes and cardiovascular risk factors in obese children after a six-month exercise program. *J Am Coll Cardiol*. 2006;48:1865–70.
- Bailey RC, Olson J, Pepper SL, et al. The level and tempo of children's physical activities: an observational study. *Med Sci Sport Exerc*. 1995;27:1033–41.
- Gibala MJ. High-intensity interval training: a time-efficient strategy for health promotion? *Curr Sports Med Rep*. 2007;6:211–3.
- Baquet G, Berthoin S, Dupont G, et al. Effects of high intensity intermittent training on peak VO₂ in prepubertal children. *Int J Sports Med*. 2002;23:439–44.
- Tjønna AE, Stølen TO, Bye A, et al. Aerobic interval training reduces cardiovascular risk factors more than a multitreatment approach in overweight adolescents. *Clin Sci*. 2009;116:317–26.
- Lau PWC, Wong DP, Ngo JK, et al. Effects of high-intensity intermittent running exercise in overweight children. *Eur J Sport Sci*. 2014;11:1–9.
- Baquet G, Berthoin S, Gerbeaux M, et al. High-intensity aerobic training during a 10 week one-hour physical education cycle: effects on physical fitness of adolescents aged 11 to 16. *Int J Sports Med*. 2001;22:295–300.
- Logan GRM, Harris N, Duncan S, et al. A review of adolescent high-intensity interval training. *Sports Med*. 2014;44:1071–85.
- Costigan SA, Eather N, Plotnikoff RC, et al. High-intensity interval training for improving health-related fitness in adolescents: a systematic review and meta-analysis. *Br J Sports Med*. 2015;49:1253–61.
- Rosenkranz SK, Rosenkranz RR, Hastmann TJ, et al. High-intensity training improves airway responsiveness in inactive nonasthmatic children: evidence from a randomized controlled trial. *J Appl Physiol*. 2012;112:1174–83.
- Weston KL, Azevedo LB, Bock S, et al. Effect of novel, school-based high-intensity interval training (HIT) on cardiometabolic health in adolescents: Project FFAB (Fun Fast Activity Blasts): an exploratory controlled before-and-after trial. *PLoS One*. 2016;11:e0159116.
- Racil G, Coquart JB, Elmontassar W, et al. Greater effects of high- compared with moderate-intensity interval training on cardio-metabolic variables, blood leptin concentration and ratings of perceived exertion in obese adolescent females. *Biol Sport*. 2016;33:145–52.
- Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med*. 2009;6:e1000100.
- Gibala MJ, McGee SL. Metabolic adaptations to short-term high-intensity interval training: a little pain for a lot of gain? *Exerc Sport Sci Rev*. 2008;36:58–63.
- Baker D. Recent trends in high-intensity aerobic training for field sports. *Prof Strength Cond*. 2011;27:4–8.
- Hood MS, Little JP, Tarnopolsky MA, et al. Low-volume interval training improves muscle oxidative capacity in sedentary adults. *Med Sci Sports Exerc*. 2011;43:1849–56.
- Little JP, Gillen JB, Percival ME. Low-volume high-intensity interval training reduces hyperglycemia and increases muscle mitochondrial capacity in patients with type 2 diabetes. *J Appl Physiol*. 2011;111:1554–60.
- Cohen J. Statistical power analysis for the behavioral sciences. New York: Routledge Academic; 1988.
- Nakagawa S, Cuthill IC. Effect size, confidence interval and statistical significance: a practical guide for biologists. *Biol Rev Camb Philos Soc*. 2007;82:591–605.

38. Hunter JE, Schmidt FL. Methods of meta-analysis: correcting error and bias in research findings. 2nd ed. Thousand Oaks: Sage; 2004.
39. Higgins JPT, Sterne JAC, Savović J, et al. A revised tool for assessing risk of bias in randomized trials. In: Chandler J, McKenzie J, Boutron I, et al., editors. Cochrane methods. Cochrane Database Syst Rev. 2016;10(Suppl. 1):CD201601.
40. Baquet G, Gamelin F, Mucci P, et al. Continuous vs. interval aerobic training in 8- to 11-year-old children. *J Strength Cond Res.* 2010;24:1381–8.
41. Baquet G, Guinhouya C, Dupont G, et al. Effects of a short-term interval training program on physical fitness in prepubertal children. *J Strength Cond Res.* 2004;18:708–13.
42. Lambrick D, Westrupp N, Kaufmann S, et al. The effectiveness of a high-intensity games intervention on improving indices of health in young children. *J Sports Sci.* 2016;34:190–8.
43. Nourry C, Deruelle F, Guinhouya C, et al. High-intensity intermittent running training improves pulmonary function and alters exercise breathing pattern in children. *Eur J Appl Physiol.* 2005;94:415–23.
44. Boddy LM, Stratton G, Hackett AF, et al. The effectiveness of a “short, sharp, shock” high intensity exercise intervention in 11- and 12-year-old Liverpool schoolgirls. *Arch Exerc Health Dis.* 2010;44:19–25.
45. Sperlich B, De Marées M, Koehler K, et al. Effects of 5 weeks of high-intensity interval training vs. volume training in 14-year-old soccer players. *J Strength Cond Res.* 2011;25:1271–8.
46. Baquet G, van Praagh E, Berthoin S. Endurance training and aerobic fitness in young people. *Sports Med.* 2003;33:1127–43.
47. McNarry MA, Mackintosh KA, Stoegefialke K. Longitudinal investigation of training status and cardiopulmonary responses in pre- and early-pubertal children. *Eur J Appl Physiol.* 2014;114:1573–80.
48. Ostoicic SM, Stojanovic M, Jukic I, et al. The effects of six weeks of training on physical fitness and performance in teenage and mature top-level soccer players. *Biol Sport.* 2009;26:379–87.
49. Mirwald RL. Longitudinal comparison of aerobic power in active and inactive boys aged 7.0 to 17.0 years. *Ann Hum Biol.* 1981;8:405–14.
50. McNarry MA, Welsman JR, Jones AM. The influence of training and maturity status on girls’ responses to short-term, high-intensity upper- and lower-body exercise. *Appl Physiol Nutr Metab.* 2011;36:344–52.
51. Funtikova AN, Navarro E, Bawaked RA, et al. Impact of diet on cardiometabolic health in children and adolescents. *Nutr J.* 2015;14:118.
52. Gillis LJ, Kennedy LC, Gillis AM, et al. Relationship between juvenile obesity, dietary energy and fat intake and physical activity. *Int J Obes.* 2002;26:458–63.
53. Ridgers ND, Timperio A, Cerin E, et al. Compensation of physical activity and sedentary time in primary school children. *Med Sci Sports Exerc.* 2014;46:1564–9.
54. Rowland TW. The biological basis of physical activity. *Med Sci Sports Exerc.* 1998;30:392–9.
55. Brodersen NH, Steptoe A, Boniface DR, et al. Trends in physical activity and sedentary behaviour in adolescence: ethnic and socioeconomic differences. *Br J Sports Med.* 2007;41:140–4.
56. Lazaar N, Aucouturier J, Ratel S, et al. Effect of physical activity intervention on body composition in young children: influence of body mass index status and gender. *Acta Paediatr.* 2007;96:1315–20.
57. Martínez-Vizcaíno V, Sánchez-López M, Notario-Pacheco B, et al. Gender differences on effectiveness of a school-based physical activity intervention for reducing cardiometabolic risk: a cluster randomized trial. *Int J Behav Nutr Phys Act.* 2014;11:154.
58. Obert P, Mandigouts S, Nottin S, et al. Cardiovascular responses to endurance training in children: effect of gender. *Eur J Clin Investig.* 2003;33:199–208.
59. Christian D, Todd C, Hill R, et al. Active children through incentive vouchers-evaluation (ACTIVE): a mixed-method feasibility study. *BMC Public Health.* 2016;16:890.
60. Mackintosh KA, Niezen G, Eslambolchilar P. Mission possible: using ubiquitous social goal sharing technology to promote physical activity in children. *Movement Health Exerc.* 2016;5:1–14.
61. Fox KR, Cooper A, McKenna J. The school and promotion of children’s health-enhancing physical activity: perspectives from the United Kingdom. *J Sch Health.* 2004;23:338–58.
62. Costigan SA, Eather N, Plotnikoff RC, et al. Preliminary efficacy and feasibility of embedding high intensity interval training into the school day: a pilot randomized controlled trial. *Prev Med Rep.* 2015;2:973–9.
63. Ratel S, Lazaar N, Dore E, et al. High-intensity intermittent activities at school: controversies and facts. *J Sports Med Phys Fit.* 2004;44:272–80.