



Swansea University
Prifysgol Abertawe



Cronfa - Swansea University Open Access Repository

This is an author produced version of a paper published in:
Annals of Pure and Applied Logic

Cronfa URL for this paper:
<http://cronfa.swan.ac.uk/Record/cronfa36024>

Paper:

Brattka, V., de Brecht, M. & Pauly, A. (2012). Closed choice and a Uniform Low Basis Theorem. *Annals of Pure and Applied Logic*, 163(8), 986-1008.
<http://dx.doi.org/10.1016/j.apal.2011.12.020>

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/library/researchsupport/ris-support/>

Closed Choice and a Uniform Low Basis Theorem[★]

Vasco Brattka

*Laboratory of Foundational Aspects of Computer Science
Department of Mathematics & Applied Mathematics
University of Cape Town, South Africa
Vasco.Brattka@uct.ac.za*

Matthew de Brecht

*Graduate School of Informatics
Kyoto University, Japan
matthew@iip.ist.i.kyoto-u.ac.jp*

Arno Pauly

*Computer Laboratory
University of Cambridge, UK
Arno.Pauly@cl.cam.ac.uk*

Abstract

We study closed choice principles for different spaces. Given information about what does not constitute a solution, closed choice determines a solution. We show that with closed choice one can characterize several models of hypercomputation in a uniform framework using Weihrauch reducibility. The classes of functions which are reducible to closed choice of the singleton space, of the natural numbers, of Cantor space and of Baire space correspond to the class of computable functions, of functions computable with finitely many mind changes, of weakly computable functions and of effectively Borel measurable functions, respectively. We also prove that all these classes correspond to classes of non-deterministically computable functions with the respective spaces as advice spaces. The class of limit computable functions can be characterized with parallelized choice on natural numbers. On top of these results we provide further insights into algebraic properties of closed choice. In particular, we prove that closed choice on Euclidean space can be considered as “locally compact choice” and it is obtained as product of closed choice on the natural numbers and on Cantor space. We also prove a Quotient Theorem for compact choice which shows that single-valued functions can be “divided” by compact choice in a certain sense. Another result is the Independent Choice Theorem, which provides a uniform proof that many choice principles are closed under composition. Finally,

we also study the related class of low computable functions, which contains the class of weakly computable functions as well as the class of functions computable with finitely many mind changes. As one main result we prove a uniform version of the Low Basis Theorem that states that closed choice on Cantor space (and the Euclidean space) is low computable. We close with some related observations on the Turing jump operation and its initial topology.

Key words: Computable analysis, Borel complexity, Weihrauch reducibility.

1 Introduction

The basic task to be studied in the present paper is the following:

Given information about what does not constitute a solution, find a solution.

The difficulty of this task depends strongly on the structure of the set of potential solutions. In general, each represented space (X, δ) induces a topology, where a set $U \subseteq X$ is open, if its characteristic function

$$\chi_U : X \rightarrow \mathbb{S}, x \mapsto \begin{cases} 1 & \text{if } x \in U \\ 0 & \text{otherwise} \end{cases}$$

is continuous with respect to the representation δ and a standard representation $\delta_{\mathbb{S}}$ of Sierpiński space $\mathbb{S} = \{0, 1\}$ (which is equipped with the topology $\{\emptyset, \{1\}, \{0, 1\}\}$). Such a standard representation of \mathbb{S} can be defined by

$$\delta_{\mathbb{S}}(p) = 1 : \iff (\exists n) p(n) = 0$$

for all $p \in \mathbb{N}^{\mathbb{N}}$. Intuitively, the open sets are those for which membership can be continuously confirmed. Each represented space then comes naturally with a representation δ° of the open sets, defined by

$$\delta^{\circ}(p) = U : \iff [\delta \rightarrow \delta_{\mathbb{S}}](p) = \chi_U$$

for all $p \in \mathbb{N}^{\mathbb{N}}$. Here $[\delta \rightarrow \delta_{\mathbb{S}}]$ denotes the canonical function space representation (see [36]) of δ and $\delta_{\mathbb{S}}$ (which is the exponential in the category of

* This work has been supported by the National Research Foundation of South Africa (NRF) and the Japanese Society for Promotion of Sciences (JSPS)

represented spaces). The representation δ° in turn induces a representation ψ_-^X of the closed sets by $\psi_-^X(p) = X \setminus \delta^\circ(p)$. The restriction to closed sets as solution sets arises from the fact that they are exactly those sets for which one can continuously confirm membership in the complement.

We give some intuitive descriptions of equivalent versions of this very general representation for concrete spaces that we will consider.

- $\mathbb{N} = \{0, 1, 2, \dots\}$, the set of natural numbers: the standard representation is defined by $\delta_{\mathbb{N}}(p) := p(0)$ and an equivalent way of defining $\psi_-^{\mathbb{N}}$ is by $\psi_-^{\mathbb{N}}(p) = \{n \in \mathbb{N} : n + 1 \notin \text{range}(p)\}$. That is $\psi_-^{\mathbb{N}}(p) = A$, if p is an enumeration of all points that are not in A .
- $\{0, 1\}^{\mathbb{N}}$, the Cantor space: the standard representation can be obtained by restricting the identity on Baire space to Cantor space $\delta_{\{0,1\}^{\mathbb{N}}} := \text{id}_{\mathbb{N}^{\mathbb{N}}}|_{\{0,1\}^{\mathbb{N}}}$. In this case one can think that $\psi_-^{\{0,1\}^{\mathbb{N}}}(p) = A$ if p is a (potentially empty) enumeration of words $w_i \in \{0, 1\}^*$ such that $A = \{0, 1\}^{\mathbb{N}} \setminus \bigcup_{i=0}^{\infty} w_i \{0, 1\}^{\mathbb{N}}$. That is p is a (potentially empty) enumeration of words w_i such that the corresponding balls exhaust the exterior of A .
- $\mathbb{N}^{\mathbb{N}}$, the Baire space: this case can be handled analogously to Cantor space, except that the representation $\delta_{\mathbb{N}^{\mathbb{N}}}$ is just the identity.
- \mathbb{R} , the Euclidean real number line (and \mathbb{R}^n in general): for convenience we assume that we use some standard numbering $\bar{\cdot} : \mathbb{N} \rightarrow \mathbb{Q}$. Then the Cauchy representation $\rho : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{R}$ can be defined by $\rho(p) := \lim_{n \rightarrow \infty} \overline{p(n)}$, where the domain $\text{dom}(\rho)$ contains only rapidly converging sequences, i.e. p with $|\overline{p(i)} - \overline{p(j)}| < 2^{-j}$ for all $i > j$. Thus, a real number x is represented by a rapidly converging sequence of rational numbers. The representation $\psi_-^{\mathbb{R}}$ can then be considered as follows: a name p of a set A is a sequence $(\langle a_i, b_i \rangle)_{i \in \mathbb{N}}$ such that $A = \mathbb{R} \setminus \bigcup_{i=0}^{\infty} (\overline{a_i}, \overline{b_i})$. That is, intuitively, p is a list of rational intervals that exhaust the complement of A .
- $\mathbb{I} := [0, 1]$, the real unit interval (and \mathbb{I}^n in general): this can be treated by restricting the case of \mathbb{R}^n .

For most spaces, closed choice is not computable. Thus, our interest lies on classifying the degree of incomputability, that is the Weihrauch degree of closed choice, depending on the underlying space. Some of the arising Weihrauch degrees are associated with certain models of type-2 hypercomputation, giving an independent justification for our interest in closed choice. Additionally, as already demonstrated in [6], several important mathematical theorems share a Weihrauch degree with an appropriate version of closed choice.

In recursion theory, a question closely related to our notion of closed choice has been studied. Given a Π_1^0 -class of Cantor space (which is a co-c.e. closed set in our terminology), what can we say about its elements? It is known that a co-c.e. closed set may contain no computable points, but always contains a low point [16]. We present a stronger result, which takes the form that closed choice

for Cantor space is computable, if we replace the standard representation of the elements with another one, which just renders the low points computable. On the side, we present a few results on the initial topology of the Turing jump operator (called Π -topology by Joseph Miller, see [21]).

2 Weihrauch Reducibility

This section serves to give a brief introduction into represented spaces, realizers, Weihrauch reducibility and several associated operations. The basic reference for this section is [36]. While the study of (variants of) Weihrauch-reducibility has commenced over a decade ago ([31], [34], [35], [15]), the relevant sources for this section are [7], [6] and [28].

A significant ingredient of the theory of represented spaces is Baire space $\mathbb{N}^{\mathbb{N}}$, i.e. the set of natural number sequences, equipped with the topology derived from the metric $d_{\mathbb{N}^{\mathbb{N}}}$ which is defined by $d_{\mathbb{N}^{\mathbb{N}}}(u, u) = 0$ and $d_{\mathbb{N}^{\mathbb{N}}}(u, v) = 2^{-\min\{n \mid u_n \neq v_n\}}$ for $u \neq v$. A useful property of Baire space to be exploited frequently is the existence of an effective and bijective pairing function $\langle \cdot, \cdot \rangle : \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$. In the following we will denote partial functions using the symbol \subseteq as prefix and multi-valued function using the double function arrow \rightrightarrows . The term “function” or “map” might refer to any of those but often we will indicate totality or single-valuedness, if relevant.

Definition 2.1 (Representation) A *representation* δ of a set X is a surjective single-valued (potentially partial) function $\delta : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow X$. A *represented space* (X, δ) is a set X together with a representation δ of it.

Using represented spaces we can define the concept of a realizer. We denote the composition of two (multi-valued) functions f and g either by $f \circ g$ or by fg .

Definition 2.2 (Realizer) Let $f : \subseteq (X, \delta_X) \rightrightarrows (Y, \delta_Y)$ be a multi-valued function between represented spaces. A *realizer* of f is a single-valued function $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ satisfying $\delta_Y \circ F(p) \in f \circ \delta_X(p)$ for all $p \in \text{dom}(f \delta_X)$. We use the notation $F \vdash f$ for expressing that F is a realizer of f .

As realizers are single-valued by definition, the statement that some function F is a realizer always implies its single-valuedness. Realizers allow us to transfer the notions of computability and continuity and other notions available for Baire space to any represented space; a function between represented spaces will be called *computable*, if it has a computable realizer, etc. Now we have

gathered the necessary provision to define Weihrauch reducibility (\leq_W):

Definition 2.3 (Weihrauch reducibility) Let $f : \subseteq X \rightrightarrows Y$ and $g : \subseteq U \rightrightarrows V$ be multi-valued functions between represented spaces. Define $f \leq_W g$, if there are computable single-valued functions $K, H : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ satisfying $K \circ \langle \text{id}, G \circ H \rangle \vdash f$ for all $G \vdash g$.

We note that the relations \leq_W and \vdash implicitly refer to the underlying representations, which we will only mention explicitly if necessary. The relation \leq_W is reflexive and transitive, thus it induces a partial order on the set of its equivalence classes (which we refer to as *Weihrauch degrees*). This partial order will be denoted by \leq_W , as well. In this sense, \leq_W is a distributive bounded lattice (for details see [28] and [7]). We use \equiv_W to denote equivalence regarding \leq_W , $<_W$ for strict reducibility and $|_W$ for incomparability. There is a slightly stronger version of Weihrauch reducibility where the condition $K \circ \langle \text{id}, G \circ H \rangle \vdash f$ is replaced by $K \circ G \circ H \vdash f$. This *strong Weihrauch reducibility* is denoted by $f \leq_{sw} g$.

We mention that the symbol \leq_W is also used to denote Wadge reducibility, which is in some sense a counterpart of Weihrauch reducibility for sets and has been studied since the early 1970s, see [32,33,29]. The double usage of \leq_W should not lead to confusion since Wadge reducibility is defined for sets and Weihrauch reducibility for functions. We mention that some further information on the history of Weihrauch reducibility is given in [7] and not repeated here.

We proceed to define a couple of useful operations. While all definitions are given in terms of functions between represented spaces, they transfer directly to the according Weihrauch degrees.

The first operation is the coproduct, which plays the role of the supremum in the Weihrauch lattice. By $X \amalg Y := (\{0\} \times X) \cup (\{1\} \times Y)$ we denote the disjoint sum of two sets X and Y and if these spaces are represented spaces, then we assume that $X \amalg Y$ is equipped with the canonical coproduct representation (see [28] for details).

Definition 2.4 (Coproduct) Let $f : \subseteq X \rightrightarrows Y$ and $g : \subseteq W \rightrightarrows Z$ be two multi-valued functions on represented spaces. Then we define $f \amalg g : \subseteq X \amalg W \rightrightarrows Y \amalg Z$ by $(f \amalg g)(0, u) := \{0\} \times f(u)$ and $(f \amalg g)(1, u) := \{1\} \times g(u)$.

One obtains that $H \vdash (f \amalg g)$ holds for exactly those H satisfying $H(0w) = F(w)$ and $H(1w) = G(w)$ for some realizers $F \vdash f$ and $G \vdash g$ (that can depend on w). We assume that the product $X \times Y$ of represented spaces (X, δ_X) and (Y, δ_Y) is represented with the canonical product representation $[\delta_X, \delta_Y]$ (see

[36] for details).

Definition 2.5 (Products) Let $f : \subseteq X \rightrightarrows Y$ and $g : \subseteq W \rightrightarrows Z$ be two multi-valued functions on represented spaces. Then we define $f \times g : \subseteq X \times W \rightrightarrows Y \times Z$ by $(f \times g)(x, w) := f(x) \times g(w)$.

One obtains that $H \vdash (f \times g)$ holds for exactly those H satisfying $H(\langle u, v \rangle) = \langle F(u), G(v) \rangle$ for some realizers $F \vdash f$ and $G \vdash g$ (that might depend on u, v).

We say that a multi-valued map f on represented spaces is *pointed*, if it contains at least one computable point in its domain and we say that it is *idempotent*, if $f \times f \equiv_W f$. In some cases the product and the coproduct are closely related. If $f \times g$ is pointed and $f \amalg g$ is idempotent, then $f \amalg g \equiv_W f \times g$, since

$$f \amalg g \leq_W f \times g \leq_W (f \amalg g) \times (f \amalg g) \leq_W (f \amalg g), \quad (1)$$

where pointedness of $f \times g$ is only required for the first reduction and idempotency of $f \amalg g$ only for the last one. It is useful to consider a countable product of a multi-valued function with itself, which has been introduced in [7].

Definition 2.6 (Parallelization) Let $f : \subseteq X \rightrightarrows Y$ be a multi-valued function on represented spaces. We define the *parallelization* $\hat{f} : \subseteq X^{\mathbb{N}} \rightrightarrows Y^{\mathbb{N}}$ by $\hat{f}(x_i)_{i \in \mathbb{N}} := \mathbf{X}_{i=0}^{\infty} f(x_i)$.

We obtain that $H \vdash \hat{f}$ holds for exactly those H satisfying $H(\langle u_1, u_2, \dots \rangle) = \langle F_1(u_1), F_2(u_2), \dots \rangle$ for some realizers $F_i \vdash f$ for $i \in \mathbb{N}$ (that might depend on u_i). We use the notation $\langle x_1, x_2, \dots \rangle$ for the canonical countable pairing on Baire space. In [28] a finite type of parallelization was introduced. For any represented space (X, δ) we denote by $X^* = \bigcup_{i=0}^{\infty} (\{i\} \times X^i)$ the set of all finite sequences over X and we assume that X^* is denoted by its canonical standard representation δ^* . For $f : \subseteq X \rightrightarrows Y$, we use f^i to denote the i -fold product of f with itself; and understand f^0 to be Weihrauch-equivalent to $\text{id}_{\mathbb{N}}$.

Definition 2.7 (Finite parallelization) Let $f : \subseteq X \rightrightarrows Y$ be a multi-valued function on represented spaces. We define the *finite parallelization* $f^* : \subseteq X^* \rightrightarrows Y^*$ by $f^* := \amalg_{i=0}^{\infty} f^i$ with $f^*(i, x) := f^i(x)$ for all $(i, x) \in X^*$.

Both types of parallelization form closure operators for the Weihrauch lattice, which means $f \leq_W \hat{f}$ and $\hat{f} \equiv_W \hat{\hat{f}}$, and $f \leq_W g$ implies $\hat{f} \leq_W \hat{g}$ and analogously for finite parallelization (see [28,26] and [7] for details). It is easy to see that for pointed multi-valued functions idempotency is equivalent to $f \equiv_W f^*$. It is interesting to mention that some variant of the (continuous) Weihrauch degrees has recently be proved to be undecidable (see [20]).

3 Closed Choice

Now we define the general version of closed choice for a represented space.

Definition 3.1 (Closed Choice) Let (X, δ) be a represented space. Then the *closed choice* operation of this space is defined by

$$C_X : \subseteq \mathcal{A}_-(X) \rightrightarrows X, A \mapsto A$$

with $\text{dom}(C_X) := \{A \in \mathcal{A}_-(X) : A \neq \emptyset\}$.

Here we assume that $\mathcal{A}_-(X)$ is the set of closed subsets of X equipped with the negative information representation ψ_-^X as defined in the introduction. The computable points in $\mathcal{A}_-(X)$ are called *co-c.e. closed sets*. Intuitively, C_X takes as input a non-empty closed set in negative description (i.e. by some form of enumeration of its complement) and it produces an arbitrary point of this set as output. Hence, if we write $A \mapsto A$, then we mean that the multi-valued map C_X maps the input A (as a point in $\mathcal{A}_-(X)$) to the set A as a subset of X , namely the set of possible function values.

Closed choice for particular spaces can characterize certain classes of functions or degrees of mathematical theorems. In [14] it was proved that $C_{\{0,1\}^{\mathbb{N}}}$ is equivalent to the Hahn-Banach Theorem and to Weak König's Lemma and in [6] it was shown that $C_{\mathbb{N}}$ is equivalent to the Baire Category Theorem, Banach's Inverse Mapping Theorem and several other theorems from functional analysis. The following example shows that also many other classes that have been considered can be characterized as classes of closed choice for certain spaces.

Example 3.2 We obtain $C_{\{0\}} \equiv_W C_{\mathbb{S}} \equiv_W \text{id}$, $C_{\{0,1\}} \equiv_W \text{LLPO}$ and, more generally, $C_{\{0,1,\dots,n\}} \equiv_W \text{MLPO}_{n+1}$.

Here MLPO_n and $\text{LLPO} = \text{MLPO}_2$ are taken from [34]. For $n \geq 1$ we consider $\text{MLPO}_n : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}$ as a multi-valued map with

$$\text{dom}(\text{MLPO}_n) := \{\langle p_1, \dots, p_n \rangle : (\exists i = 1, \dots, n) p_i = \widehat{0}\}$$

and

$$\text{MLPO}_n \langle p_1, \dots, p_n \rangle := \{i : p_i = \widehat{0}\}.$$

Since LLPO is not idempotent (see [7]), it follows that closed choice is not necessarily idempotent. However, it is a straightforward observation that closed

choice is always pointed, since X is always a co-c.e. closed subset of itself.

Lemma 3.3 (Pointedness) *If X is a non-empty represented space, then \mathbf{C}_X is pointed.*

We get the following first result.

Proposition 3.4 (Products) *Let X and Y be non-empty represented spaces. We obtain $\mathbf{C}_X \amalg \mathbf{C}_Y \leq_W \mathbf{C}_X \times \mathbf{C}_Y \leq_W \mathbf{C}_{X \times Y}$.*

Proof. As mentioned in the introduction, coproducts are reducible to products for all pointed functions. It is easy to prove that the Cartesian product $P : \mathcal{A}_-(X) \times \mathcal{A}_-(Y) \rightarrow \mathcal{A}_-(X \times Y), (A, B) \mapsto A \times B$ is computable and we obtain $\mathbf{C}_X \times \mathbf{C}_Y = \mathbf{C}_{X \times Y} \circ P$. Hence $\mathbf{C}_X \times \mathbf{C}_Y \leq_W \mathbf{C}_{X \times Y}$. \square

We will see in Corollary 5.7 that the inverse of the first reduction does not hold in general. Also the second reduction cannot be reversed in general, as the following result shows. We denote by $\hat{n} := nnn\dots \in \mathbb{N}^{\mathbb{N}}$ the constant sequence with value n .

Proposition 3.5 (Products of choice for finite spaces) *Let A and B be finite sets, each with at least two elements and equipped with the discrete representation and topology. Then $\mathbf{C}_A \times \mathbf{C}_B <_W \mathbf{C}_{A \times B}$.*

Proof. We assume that $A = \{0, \dots, n\}$ and $B = \{0, \dots, k\}$ with $n, k \geq 1$ and we assume that A is represented by $\delta_A(n\hat{0}) := n$ with $\text{dom}(\delta_A) = A \times \{\hat{0}\}$. Moreover, we assume $\psi_-^A(p) = \{i : i+1 \notin \text{range}(p)\}$ with $\text{range}(p) \subseteq \{0, \dots, n+1\}$. This representation is computably equivalent to the generic definition of ψ_-^A given above. Analogous assumptions are made for the representations δ_B and ψ_-^B and $\delta_{A \times B}$ and $\psi_-^{A \times B}$. We have $\mathbf{C}_A \times \mathbf{C}_B \leq_W \mathbf{C}_{A \times B}$ by Proposition 3.4.

Let us now assume that $\mathbf{C}_{A \times B} \leq_W \mathbf{C}_A \times \mathbf{C}_B$ holds. Then there are computable functions H, K such that $F = H\langle \text{id}, GK \rangle$ is a realizer of $\mathbf{C}_{A \times B}$ for any realizer G of $\mathbf{C}_A \times \mathbf{C}_B$. Now we consider $\hat{0} = 000\dots$ which represents $\psi_-^{A \times B}(\hat{0}) = A \times B$. Then $(L, R) := [\psi_-^A, \psi_-^B]K(\hat{0})$ is a pair of finite sets.¹ For all m and for all $p \in \text{dom}(\psi_-^{A \times B})$ we have $\psi_-^{A \times B}(p) = \psi_-^{A \times B}(0^m p)$. Moreover, by continuity of K and since $A \times B$ is finite, there is $m \in \mathbb{N}$ such that for all $p \in \text{dom}(\psi_-^{A \times B})$, we obtain that $(L', R') = [\psi_-^A, \psi_-^B]K(0^m p)$ implies $L' \subseteq L$ and $R' \subseteq R$. By continuity of H and since $A \times B$ is equipped with the discrete representation, this m can be taken such that $H\langle 0^m p, q \rangle$ is identical to $H\langle \hat{0}, q \rangle$ for any fixed name q of an element of $L' \times R'$. Finally, since there are only finitely many

¹ We are thankful to one of the referees for providing a version of this paragraph that clarified and corrected the earlier version of it.

such q , this m can be selected as satisfying this property for all those q . Hence, for such m we obtain $F(0^m p) = H(\widehat{0}, GK(0^m p))$. Since $\psi_-^{A \times B}$ is computably equivalent to the representation ψ_m given by $\psi_m(0^m p) := \psi_-^{A \times B}(p)$, we can assume without loss of generality that there are computable functions H, K such that $F = HGK$ is a realizer of $\mathbb{C}_{A \times B}$ for any realizer G of $\mathbb{C}_A \times \mathbb{C}_B$.

Let $M_j \subsetneq M_{j-1} \subsetneq \dots \subsetneq M_0$ now be a strictly decreasing sequence of non-empty subsets $M_i \subseteq A \times B$. Due to continuity of K there is a monotone sequence of words $w_0 \sqsubseteq w_1 \sqsubseteq \dots \sqsubseteq w_j$ such that $\psi_-^{A \times B}(p_i) = M_i$ for $p_i := w_i \widehat{0}$ and such that the sets $(L_i, R_i) := [\psi_-^A, \psi_-^B]K(p_i)$ are component wise monotone as well. That is $\emptyset \neq L_j \subseteq L_{j-1} \subseteq \dots \subseteq L_0$ and $\emptyset \neq R_j \subseteq R_{j-1} \subseteq \dots \subseteq R_0$. The cardinality of $A \times B$ is $(n+1)(k+1)$ and hence the longest strictly decreasing chain (M_i) of non-empty sets is one with length $j+1 = (n+1)(k+1)$. The longest decreasing chain (L_i, R_i) with the property that for each $i < j$ the left component or the right component is strictly decreasing, i.e. $L_{i+1} \subsetneq L_i$ or $R_{i+1} \subsetneq R_i$, has length $n+k+1$. For $n, k \geq 1$ we have that $n+k+1 < (n+1)(k+1)$. Hence, there has to be at least one $i < j$ such that $(L_i, R_i) = (L_{i+1}, R_{i+1})$. By assumption there is some element $x \in M_i \setminus M_{i+1}$. For each element $y \in L_i \times R_i$ there is a realizer G_y of $\mathbb{C}_A \times \mathbb{C}_B$ with $y = [\delta_A, \delta_B]G_y K(p_{i+1})$ and by assumption $z := [\delta_A, \delta_B]HG_y K(p_{i+1}) \in M_{i+1}$ and hence $z \neq x$. By continuity of K there is an extension w of w_i such that $\psi_-^{A \times B}(p) = \{x\}$ for $p := w \widehat{0}$ and $[\psi_-^A, \psi_-^B]K(p) \subseteq (L_i, R_i)$ (where the inclusion is meant component wise). Hence any realizer G of $\mathbb{C}_A \times \mathbb{C}_B$ selects an element $y = [\delta_A, \delta_B]GK(p) \in L_i \times R_i$ and thus $[\delta_A, \delta_B]HGK(p) \neq x$ in contrast to the fact that HGK is supposed to be a realizer of $\mathbb{C}_{A \times B}$. Contradiction! \square

Alternatively, one could prove this result by considering the level of the respective operations, a concept that has been introduced by Hertling [15]. For one, one can prove directly $\text{MLPO}_{n+1} \times \text{MLPO}_{k+1} \leq_W \text{LPO}^{n+k}$, which implies that $n+k$ is an upper bound on the level of $\text{MLPO}_{n+1} \times \text{MLPO}_{k+1}$. On the other hand, $\text{LPO}^{(n+1)(k+1)-1}$ can be reduced to any realizer of $\text{MLPO}_{(n+1)(k+1)}$ (see Theorem 5.2.2 in [35]), which implies that the level of $\text{MLPO}_{(n+1)(k+1)}$ is at least $(n+1)(k+1)-1$. Since Hertling proved that the level is preserved downwards by Weihrauch reducibility, the desired result follows also from these observations. We do not work out the details here. For the simplest case of the set $\{0, 1\}$ we get the following conclusion.

Corollary 3.6 $\mathbb{C}_{\{0,1\}} \times \mathbb{C}_{\{0,1\}} <_W \mathbb{C}_{\{0,1\} \times \{0,1\}}$.

We will see, however, that for many infinite spaces we get a nicer behavior of products. This is partially due to the following result.

Proposition 3.7 (Surjections) *Let A and B be represented spaces and let $s : \subseteq A \rightarrow B$ be a computable surjection with a co-c.e. closed domain $\text{dom}(s)$.*

Then $\mathcal{C}_B \leq_W \mathcal{C}_A$.

Proof. If $s : \subseteq A \rightarrow B$ is computable and $\text{dom}(s)$ is co-c.e. closed in A , then $S : \mathcal{A}_-(B) \rightarrow \mathcal{A}_-(A), M \mapsto s^{-1}(M)$ is computable too and if s is surjective, then we obtain $\mathcal{C}_B = s \circ \mathcal{C}_A \circ S$, i.e. $\mathcal{C}_B \leq_W \mathcal{C}_A$. \square

As a consequence of this observation and Proposition 3.4 we obtain the following sufficient criterion for idempotency of choice.

Corollary 3.8 (Idempotency) *Let A be a represented space. If there is a computable surjection $s : A \rightarrow A^2$, then $\mathcal{C}_A \times \mathcal{C}_A \equiv_W \mathcal{C}_{A \times A} \equiv_W \mathcal{C}_A$ and, in particular, \mathcal{C}_A is idempotent and hence also $\mathcal{C}_A^* \equiv_W \mathcal{C}_A$.*

Since the spaces $\mathbb{N}, \{0, 1\}^{\mathbb{N}}, \mathbb{N}^{\mathbb{N}}$ and $\mathbb{N} \times \{0, 1\}^{\mathbb{N}}$ admit computable and bijective pairing functions, we get the following conclusion.

Corollary 3.9 *The choice principles $\mathcal{C}_{\mathbb{N}}, \mathcal{C}_{\{0,1\}^{\mathbb{N}}}, \mathcal{C}_{\mathbb{N}^{\mathbb{N}}}$ and $\mathcal{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}$ are idempotent.*

We close this section with the following example that shows that in some cases choice commutes with parallelization and finite parallelization and in other cases it does not.

Example 3.10 *We obtain $\widehat{\mathcal{C}}_{\{0,1\}} \equiv_W \widehat{\text{LLPO}} \equiv_W \mathcal{C}_{\{0,1\}^{\mathbb{N}}}$, but $\widehat{\mathcal{C}}_{\mathbb{N}} \equiv_W \lim <_W \mathcal{C}_{\mathbb{N}^{\mathbb{N}}}$ and $\mathcal{C}_{\mathbb{N}}^* \equiv_W \mathcal{C}_{\mathbb{N}} \equiv_W \mathcal{C}_{\mathbb{N}^*}$, but $\mathcal{C}_{\{0,1\}}^* \equiv_W \text{LLPO}^* <_W \mathcal{C}_{\mathbb{N}} \equiv_W \mathcal{C}_{\{0,1\}^*}$.*

4 Choice on Computable Metric Spaces

In this section we want to study choice on certain large classes of computable metric spaces. We recall that a *computable metric space* (X, d, α) is a separable metric space (X, d) together with a numbering $\alpha : \mathbb{N} \rightarrow X$ of a countable dense subset with respect to which the metric is computable. By a *computable Polish space* we mean a computable metric space that is also complete. Usually, we will assume that computable metric spaces are represented by their Cauchy representations δ_X (see [36]). We use two different representation κ_- and κ to represent the set $\mathcal{K}(X)$ of compact subsets of a computable metric space X (see [9] for details). Roughly speaking, a κ_- -name of a compact set $K \subseteq X$ is a list of all finite covers of K by rational open balls, whereas a κ -name comes with the additional requirement that all open balls in the cover actually have non-empty intersection with K . That is, κ_- provides negative information on the set K (each cover allows to exclude points) and κ provides full information

(each ball in the cover meets the set). By $\mathcal{K}_-(X)$ and $\mathcal{K}(X)$ we denote the set of compact subsets represented by κ_- and κ , respectively. The compact sets that are computable with respect to κ_- and κ are called *co-c.e. compact* and *computably compact*, respectively. We mention that a computable metric space is computably compact in itself if and only if it is co-c.e. compact in itself.

Computable Polish spaces X admit total computable and admissible representations $\delta : \mathbb{N}^{\mathbb{N}} \rightarrow X$ (see, for instance, Corollary 4.4.12 in [2]) and computably compact computable metric spaces X admit computable representations $\delta : \{0, 1\}^{\mathbb{N}} \rightarrow X$ as we will prove next. Two representations δ_1, δ_2 of the same set are said to be (*computably*) *reducible* to each other, in symbols $\delta_1 \leq \delta_2$, if there exists a computable function $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $\delta_1 = \delta_2 \circ F$. Moreover, δ_1 and δ_2 are said to be (*computably*) *equivalent*, in symbols $\delta_1 \equiv \delta_2$, if $\delta_1 \leq \delta_2$ and $\delta_2 \leq \delta_1$ hold. We recall that a representation of a computable metric space is called *computably admissible* if it is computably equivalent to the Cauchy representation of the space.

Proposition 4.1 *Let X be a computably compact computable metric space. Then there is a surjective computable map $\varphi : \{0, 1\}^{\mathbb{N}} \rightarrow X$ that is also computably admissible.*

Proof. Let (X, d, α) be a computably compact computable metric space. We use a version $\delta_X : \subseteq \{0, 1\}^{\mathbb{N}} \rightarrow X$ of the Cauchy representation, defined as follows

$$\delta_X(01^{n_0+1}01^{n_1+1}0\dots) := \lim_{i \rightarrow \infty} \alpha(n_i)$$

where $\text{dom}(\delta_X)$ contains only those sequences of the given type which, additionally, converge rapidly, i.e. such that $d(\alpha(n_i), \alpha(n_j)) < 2^{-j}$ for all $i \geq j$. It is known that there exists a computably proper and computably admissible representation $\delta : \subseteq \{0, 1\}^{\mathbb{N}} \rightarrow X$ that is a restriction of δ_X , see Corollary 4.6 in [37]. Such a map is, in particular, computable and surjective and the fact that it is computably proper implies that $\delta^{-1}(K)$ is co-c.e. compact for any co-c.e. compact $K \subseteq X$. If X itself is co-c.e. compact, then $A := \text{dom}(\delta) = \delta^{-1}(X)$ is also co-c.e. compact. We claim that there is a total computable map $\iota : \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 1\}^{\mathbb{N}}$ such that $A \subseteq \text{range}(\iota) \subseteq \text{dom}(\delta_X)$. A machine computing ι works as follows: given an input $p \in \{0, 1\}^{\mathbb{N}}$ the machine checks in steps longer and longer prefixes w of p for the property

$$w\{0, 1\}^{\mathbb{N}} \subseteq \{0, 1\}^{\mathbb{N}} \setminus A. \tag{2}$$

Since A is co-c.e. closed, this property is c.e. in w . As long as the property cannot be verified, the machine simultaneously checks whether the input is of

the form $p = 01^{n_0+1}01^{n_1+1}0\dots$ and whether the property $d(\alpha(n_i), \alpha(n_j)) < 2^{-j}$ is satisfied for all $i \geq j$ such that $01^{n_i+1}0$ is completely included in w . If the latter property is positively verified, then the output is extended such that it matches p up to the corresponding $01^{n_i+1}0$. If, at any time, property (2) is positively verified, then it is clear that $p \notin A$ and the processing of the input is stopped and the output is extended just by infinitely many repetitions of the last block $1^{n_i+1}0$ (if no block has been written at this stage, then an arbitrary block 01^{n_0+1} is repeated infinitely often as output). If the input p is not of the form $p = 01^{n_0+1}01^{n_1+1}0\dots$, then the test for property (2) will eventually be positive. It is clear that altogether this machine computes a function $\iota : \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 1\}^{\mathbb{N}}$ such that $A \subseteq \text{range}(\iota) \subseteq \text{dom}(\delta_X)$. This guarantees that $\varphi := \delta_X \circ \iota$ is computable, total and surjective. Since $\text{dom}(\delta) = A \subseteq \text{range}(\iota)$ it follows that $\delta = \varphi \circ \iota^{-1}$. Since ι is computable, also ι^{-1} is computable (see Corollary 6.7) and hence it follows that φ is computably admissible. \square

Hence we obtain the following corollary. The first statement is a consequence of Proposition 3.7 and the second statement a consequence of the previous Proposition 4.1.

Corollary 4.2 *Let X be a computable Polish space. Then $\mathcal{C}_X \leq_W \mathcal{C}_{\mathbb{N}^{\mathbb{N}}}$. If, additionally, X is computably compact, then $\mathcal{C}_X \leq_W \mathcal{C}_{\{0,1\}^{\mathbb{N}}}$.*

We say that $\iota : A \rightarrow B$ is a *computable embedding*, if ι is computable and injective and its partial inverse ι^{-1} is computable too. Now we can use the Embedding Theorem 3.7 from [8] in order to obtain the following proposition.

Proposition 4.3 *Let A and B be computable metric spaces and let $\iota : A \rightarrow B$ be a computable embedding such that $\text{range}(\iota)$ is co-c.e. closed in B . Then $\mathcal{C}_A \leq_W \mathcal{C}_B$.*

Proof. From Theorem 3.7 in [8] it follows that for a computable embedding $\iota : A \rightarrow B$ with co-c.e. closed range $\iota(A)$ the map $J : \mathcal{A}_-(A) \rightarrow \mathcal{A}_-(B), M \mapsto \iota(M)$ is computable. We obtain $\mathcal{C}_A = \iota^{-1} \circ \mathcal{C}_B \circ J$ and hence $\mathcal{C}_A \leq_W \mathcal{C}_B$. \square

We recall that a metric space is called perfect, if it has no isolated points. In Proposition 6.2 in [8] it has been proved that any non-empty perfect computable Polish space is rich, i.e. admits a computable embedding $\iota : \{0, 1\}^{\mathbb{N}} \rightarrow X$ and in this case $\text{range}(\iota)$ is automatically co-c.e. closed. Hence we obtain the following corollary.

Corollary 4.4 *Let X be a computable Polish space. If X is rich and, in particular, if X is non-empty and has no isolated points, then $\mathcal{C}_{\{0,1\}^{\mathbb{N}}} \leq_W \mathcal{C}_X$.*

Together with Corollary 4.2 we get the following corollary (which has essentially been proved in [14] already).

Corollary 4.5 *Let X be a computably compact metric space, which is non-empty and has no isolated points, then $C_{\{0,1\}^{\mathbb{N}}} \equiv_W C_X$.*

Thus, $C_{\{0,1\}^{\mathbb{N}}}$ can be identified with “compact choice” for a very large class of compact spaces. In particular, we obtain the following corollary.

Corollary 4.6 $C_{\{0,1\}^{\mathbb{N}}} \equiv_W C_{[0,1]} \equiv_W C_{[0,1]^{\mathbb{N}}}$.

We would like to show that $C_{\mathbb{N}} \times C_{\{0,1\}^{\mathbb{N}}}$ plays a similar role for locally compact spaces as $C_{\{0,1\}^{\mathbb{N}}}$ does for compact spaces. The following lemma plays a role in the proof of the next result and it is worth being formulated separately.

Lemma 4.7 *Let K be a non-empty computably compact computable metric space. Then $C_K : \subseteq \mathcal{A}_-(K) \rightrightarrows K$ has a total extension $C'_K : \mathcal{A}_-(K) \rightrightarrows K$ with $C_K \equiv_W C'_K$.*

Proof. The set $\{A \in \mathcal{A}_-(K) : A = \emptyset\}$ is c.e. open for co-c.e. compact K . Since K is computably compact, we can assume by Proposition 4.1 without loss of generality that K is represented by a total representation $\delta : \{0,1\}^{\mathbb{N}} \rightarrow K$. Hence C_K can be extended to a suitable C'_K as follows: a realizer F of C_K is modified to a map G such that never anything else but zeros and ones are written on the output tape and as soon as the empty set is detected as input, the output is just continued with constant zeros. In any other respect, the map G behaves exactly as F . Due to totality of δ , this output of G is in the domain of δ . The modification guarantees that the empty set as input leads to some infinite output and non-empty sets are treated by G exactly as by F . The construction shows that C'_K is reducible to C_K . The reverse direction follows since C'_K is an extension of C_K . \square

We note that not every multi-valued operation has a total equivalent extension (as robust division shows, see [26]).

Classically, a space X is called σ -compact or K_σ -space, if it can be written as a countable union of compact sets. For many spaces this property is somewhat weaker than local compactness, this holds in particular for represented Hausdorff spaces. The induced topology of every represented space is known to be hereditarily Lindelöf (see Lemma 2.5 in [1]) and this means that if it is, additionally, a Hausdorff space, then local compactness implies σ -compactness. This is the reason why we speak about “locally compact choice” for short. We say that X is a *computable K_σ -space*, if X is a computable metric space, such

that there exists a computable sequence $(K_i)_{i \in \mathbb{N}}$ of non-empty computably compact sets with $X = \bigcup_{i=0}^{\infty} K_i$.

Proposition 4.8 (Locally compact choice) *Let X be a computable K_σ -space. Then $\mathbf{C}_X \leq_W \mathbf{C}_{\mathbb{N}} \times \mathbf{C}_{\{0,1\}^{\mathbb{N}}}$.*

Proof. We consider the total extensions \mathbf{C}'_{K_i} of choice that exist according to a uniform version of Lemma 4.7. Using a uniform version of Corollary 4.2, we obtain

$$F := \mathbf{C}'_{K_0} \times \mathbf{C}'_{K_1} \times \mathbf{C}'_{K_2} \times \dots \leq_W \widehat{\mathbf{C}_{\{0,1\}^{\mathbb{N}}}} \equiv_W \mathbf{C}_{\{0,1\}^{\mathbb{N}}}.$$

Given a closed set $A \subseteq X$ we can compute the sequence $(A \cap K_n)_{n \in \mathbb{N}}$ of co-c.e. compact sets and hence we can enumerate the set $\{n \in \mathbb{N} : A \cap K_n = \emptyset\}$. This implies that we can find an n such that $A \cap K_n \neq \emptyset$ with the help of $\mathbf{C}_{\mathbb{N}}$. Moreover, $F((A \cap K_n)_{n \in \mathbb{N}})$ can be obtained with the help of $\mathbf{C}_{\{0,1\}^{\mathbb{N}}}$, as indicated above. Altogether, this shows $\mathbf{C}_X \leq_W \mathbf{C}_{\mathbb{N}} \times \mathbf{C}_{\{0,1\}^{\mathbb{N}}}$. \square

This result can even be generalized to the case that the K_σ -space is only co-c.e. compact in the sense that the sequence $(K_i)_{i \in \mathbb{N}}$ is only a computable sequence of co-c.e. compact sets. However, in this case the uniform version of Lemma 4.7 needs some extra attention since the extensions \mathbf{C}'_K might not always produce a value in K (but only some infinite sequence). By Proposition 3.4 we have $\mathbf{C}_{\mathbb{N}} \times \mathbf{C}_{\{0,1\}^{\mathbb{N}}} \leq_W \mathbf{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}$. On the other hand, we can apply the previous proposition to the K_σ -space $\mathbb{N} \times \{0,1\}^{\mathbb{N}}$ (with $K_n := \{n\} \times \{0,1\}^{\mathbb{N}}$) and we get the inverse reduction. We can also apply the previous proposition to \mathbb{R}^k (with $K_n := [-n, n]^k$).

Corollary 4.9 $\mathbf{C}_{\mathbb{R}^k} \equiv_W \mathbf{C}_{\mathbb{R}} \equiv_W \mathbf{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}} \equiv_W \mathbf{C}_{\mathbb{N}} \times \mathbf{C}_{\{0,1\}^{\mathbb{N}}}$ for all $k \geq 1$.

We mention that by the Theorem of Hurewicz (see Theorem 7.10 in [18]) any Polish space which is not K_σ admits an embedding $\iota : \mathbb{N}^{\mathbb{N}} \rightarrow X$ such that $\text{range}(\iota)$ is closed. Using relativized topological versions of Propositions 4.3 and 4.8 and Corollary 4.2 we obtain the following dichotomy.

Corollary 4.10 (Dichotomy) *If X is a Polish space, then there is an oracle such that either $\mathbf{C}_X \leq_W \mathbf{C}_{\mathbb{R}}$ or $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}} \equiv_W \mathbf{C}_X$, relatively to that oracle (i.e. with continuous reductions).*

In other words, topologically the interval between $\mathbf{C}_{\mathbb{R}}$ and $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$ is not inhabited by choice principles of Polish spaces. It is not too hard to see that for many computable metric spaces X that are not K_σ , such as $\mathbb{R}^{\mathbb{N}}$, $\mathcal{C}[0,1]$ and ℓ_p , there

is a computable embedding $\iota : \mathbb{N}^{\mathbb{N}} \rightarrow X$ with a co-c.e. closed image. Hence we get the following corollary of Proposition 4.3.

Corollary 4.11 $C_{\mathbb{N}^{\mathbb{N}}} \equiv_{\text{w}} C_{\mathbb{R}^{\mathbb{N}}} \equiv_{\text{w}} C_{\ell_p} \equiv_{\text{w}} C_{C[0,1]}$ for all computable real $p \geq 1$.

The results mentioned so far in this section are mostly applicable to Polish spaces. We mention two further examples for non-Polish spaces. Any sequence $(x_n)_{n \in \mathbb{N}}$ in X can be seen as a surjection from \mathbb{N} onto the range of the sequence. Hence we obtain the following corollary.

Corollary 4.12 Let X be a represented space and let $(x_n)_{n \in \mathbb{N}}$ be a computable sequence in X with $R := \{x_n : n \in \mathbb{N}\}$. Then $C_R \leq_{\text{w}} C_{\mathbb{N}}$.

This can, in particular, be applied to the rational numbers as a subspace of Euclidean space.

Corollary 4.13 $C_{\mathbb{Q}} \equiv_{\text{w}} C_{\mathbb{N}}$, independently of whether \mathbb{Q} is equipped with the discrete representation and topology or with the Euclidean one.

The irrational numbers are computably homeomorphic to Baire space (with respect to the Euclidean topology and via their continued fraction representation) and hence we get the following conclusion.

Corollary 4.14 $C_{\mathbb{R} \setminus \mathbb{Q}} \equiv_{\text{w}} C_{\mathbb{N}^{\mathbb{N}}}$.

5 Compact Choice, Quotients and Join-Irreducibility

The following theorem shows that any single-valued function f that can be computed from compact choice and another function g can already be computed from g alone. Thus, we can “divide” by compact choice in such a situation. This result generalizes Corollary 8.8 in [7].

Theorem 5.1 (Quotients) Let X be a represented space and Y be a computable metric space and let g be a multi-valued function on represented spaces. If $f : \subseteq X \rightarrow Y$ is single-valued and $f \leq_{\text{w}} C_{\{0,1\}^{\mathbb{N}}} \times g$, then $f \leq_{\text{w}} g$.

Proof. We use the Cauchy representation δ_Y for Y and canonical projections $\pi_i : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ with $\pi_1 \langle p, q \rangle = p$ and $\pi_2 \langle p, q \rangle = q$. Now let $f : \subseteq X \rightarrow Y$ be such that $f \leq_{\text{w}} C_{\{0,1\}^{\mathbb{N}}} \times g$. Hence there are computable functions H and K such that $K \langle \text{id}, PH \rangle$ is a realizer of f for any realizer P of $C_{\{0,1\}^{\mathbb{N}}} \times g$. Since H and K are computable, as well as the Cartesian product on compact sets, it follows

from Theorem 3.3 in [37] that there is a computable function $S : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ with

$$\kappa_- S\langle p, q \rangle = \delta_Y K\langle \{p\} \times \langle \kappa_- \pi_1 H(p) \times \{q\} \rangle \rangle$$

for all $p \in \text{dom}(f\delta_X)$ and suitable q . We now consider the function $T : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ with $T(p) = S\langle p, G\pi_2 H(p) \rangle$. Whenever G is a realizer of g , then T is a realizer of the function $F : \subseteq X \rightarrow \mathcal{K}_-(Y), x \mapsto \{f(x)\}$. Hence, $F \leq_W g$. If the space Y is a computable metric space, then $\text{in} : Y \rightarrow \mathcal{K}_-(Y), x \mapsto \{x\}$ has a computable inverse (see Lemma 6.4 in [5]) and it follows that $f = \text{in}^{-1} \circ F$. That implies $f \leq_W F$. \square

We note that this theorem can be generalized to larger classes of spaces Y . The only property that is exploited is that the injection $\text{in} : Y \rightarrow \mathcal{K}_-(Y)$ has a computable inverse. We obtain some straightforward corollaries.

Corollary 5.2 *Let X be a represented space and Y be a computable metric space. If $f : \subseteq X \rightarrow Y$ is single-valued and $f \leq_W \mathbf{C}_{\{0,1\}^{\mathbb{N}}}$, then f is computable.*

This is just Corollary 8.8 from [7]. Together with Corollary 4.9 we obtain the following result, which is new.

Corollary 5.3 *Let X be a represented space and Y be a computable metric space. If $f : \subseteq X \rightarrow Y$ is single-valued and $f \leq_W \mathbf{C}_{\mathbb{R}}$, then $f \leq_W \mathbf{C}_{\mathbb{N}}$.*

By exploiting the distributivity of the Weihrauch lattice discovered in [28], a restricted version of Theorem 5.1 could be obtained, using coproducts instead of products. Combined with the observation that coproducts are the suprema in the Weihrauch lattice, and the usefulness of the decomposition into products presented in Corollary 4.9, it seems sensible to explore whether any of our principles of closed choice can be expressed as a supremum of other degrees. The negative answer is a consequence of the next result. To formulate it, we define the concept of join-irreducibility in the Weihrauch lattice.

Definition 5.4 (Join-irreducibility) A multi-valued function f on represented spaces is called *join-irreducible*, if $f \equiv_W \coprod_{n \in \mathbb{N}} f_n$ implies the existence of an $n_0 \in \mathbb{N}$ with $f \equiv_W f_{n_0}$.

We note that for finitely many f_n , this is exactly the ordinary lattice theoretic concept of join-irreducibility. For countably many f_n , this concept might be called σ -join-irreducibility (see [29]). However, this is also not quite appropriate since the coproduct $\coprod_{n \in \mathbb{N}} f_n$ is not necessarily the supremum of the f_n . This is correct for continuous reducibility, but not for the computable

case. We refrain to introduce another name and call the above concept just join-irreducibility, which is justified since we will basically only apply it in a situation with finitely many f_n .

If $f : \subseteq X \rightrightarrows Y$ is a function between represented spaces, with representation δ of X , then we define f_A for each set $A \subseteq \mathbb{N}^{\mathbb{N}}$ as follows. We let $(X_A, \delta|_A)$ be the represented space with $X_A := \delta(A)$ and the restriction $\delta|_A$ of δ to A . Then $f_A : \subseteq X_A \rightrightarrows Y$ is the restriction of f to the represented space $(X_A, \delta|_A)$. That is, we obtain $F|_A \vdash f_A$ if $F \vdash f$. Using this concept, we get the following sufficient criterion for join-irreducibility.

Lemma 5.5 (Join-irreducibility) *Let (X, δ_X) and Y be represented spaces. Assume that for some multi-valued function $f : \subseteq X \rightrightarrows Y$ the equivalence $f \equiv_{\mathbb{W}} f_A$ holds for each non-empty set $A \subseteq \mathbb{N}^{\mathbb{N}}$ that is clopen in $\text{dom}(f\delta_X)$. Then f is join-irreducible.*

Proof. Assume $f \leq_{\mathbb{W}} \coprod_{n \in \mathbb{N}} f_n$. Then there exists a computable function $\mathcal{N} : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}$ with $f_{\mathcal{N}^{-1}(n)} \leq_{\mathbb{W}} f_n$ and $\text{dom}(\mathcal{N}) = \text{dom}(f\delta_X)$. There has to be an $n_0 \in \mathbb{N}$, so that $\mathcal{N}^{-1}(n_0) \neq \emptyset$, and due to continuity of \mathcal{N} , this set is closed and open in $\text{dom}(f\delta_X)$. Thus, by the assumption, we have $f \leq_{\mathbb{W}} f_{\mathcal{N}^{-1}(n_0)} \leq_{\mathbb{W}} f_{n_0}$. The other direction is trivial. \square

If we take away finitely many small open rational balls from \mathbb{N} , $\{0, 1\}^{\mathbb{N}}$, $\mathbb{N}^{\mathbb{N}}$ or \mathbb{R} , respectively, such that the remainder is non-empty, then the remainder is still large enough to simulate closed choice of the entire space within this subspace. This is why closed choice for all these spaces satisfies the above criterion for join-irreducibility.

Corollary 5.6 $\mathbb{C}_{\mathbb{N}}$, $\mathbb{C}_{\{0,1\}^{\mathbb{N}}}$, $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ and $\mathbb{C}_{\mathbb{R}}$ are join-irreducible.

Another consequence is that the coproduct (i.e. the supremum) of $\mathbb{C}_{\mathbb{N}}$ and $\mathbb{C}_{\{0,1\}^{\mathbb{N}}}$ is strictly below the product.

Corollary 5.7 $\mathbb{C}_{\mathbb{N}} \coprod \mathbb{C}_{\{0,1\}^{\mathbb{N}}} <_{\mathbb{W}} \mathbb{C}_{\mathbb{N}} \times \mathbb{C}_{\{0,1\}^{\mathbb{N}}}$.

This corollary also shows that the coproduct of two idempotent functions is not necessarily idempotent (see Equation (1)).

Corollary 5.8 $\mathbb{C}_{\mathbb{N}} \coprod \mathbb{C}_{\{0,1\}^{\mathbb{N}}}$ is not idempotent.

6 Unique Choice and Inversion

In this section we briefly discuss a variant of choice, which we call unique choice. This is choice restricted to the special case of singletons. We only formulate unique choice for Hausdorff spaces in order to guarantee that singletons are closed.

Definition 6.1 (Unique Closed Choice) Let (X, δ) be a represented Hausdorff space. We consider the injection $\text{in}_X : X \hookrightarrow \mathcal{A}_-(X), x \mapsto \{x\}$. The partial inverse $\text{UC}_X : \subseteq \mathcal{A}_-(X) \rightarrow X$ of this injection is called *unique closed choice* operation of the space X .

Since unique choice UC_X is a restriction of choice C_X , it is clear that $\text{UC}_X \leq_W \text{C}_X$ holds. In some cases we can say more. In case of \mathbb{N} it turns out that unique choice is not easier than full choice. The proof idea is very similar to the proof idea of Proposition 3.3 in [6], where $\text{C}_{\mathbb{N}}$ is reduced to finite choice. We only describe it informally here.

Proposition 6.2 $\text{UC}_{\mathbb{N}} \equiv_W \text{C}_{\mathbb{N}}$.

Proof. It is clear that $\text{UC}_{\mathbb{N}} \leq_W \text{C}_{\mathbb{N}}$. We prove $\text{C}_{\mathbb{N}} \leq_W \text{UC}_{\mathbb{N}}$ by an intuitive description of a suitable algorithm. Given an enumeration n_0, n_1, \dots of the complement of a set $A \subseteq \mathbb{N}$, we choose $c = 0$ as starting candidate for a potential element in A and we choose $j = 0$ as starting position to keep track of where we have to change our mind. In steps $i = 0, 1, \dots$ we inspect the enumeration n_i in order to find the candidate c and simultaneously we start to generate as output a negative description of $\{j\}$ by enumerating all numbers $k > j$. Whenever some i with $c = n_i$ is found, we choose as new candidate c the minimal element $c = \min(\mathbb{N} \setminus \{n_0, \dots, n_i\})$. Whenever that happens, we choose $j = \max\{i, m + 1\}$ as new position, where m is the largest number that has been produced on the output and now we start to produce as output a negative description of $\{j\}$ by enumerating all numbers $m + 1, \dots, j - 1$ (if there are any) and then all numbers $k > j$, while we continue to inspect the sequence n_{i+1}, n_{i+2}, \dots to find the new candidate c . If we continue like this, then eventually we will find a candidate c that is actually in A and hence not in the enumeration of the n_i . The output will then be a negative description of $\{j\}$ for some number j that is larger than or equal to the last position in the enumeration where we had to change our candidate. That is, the number j together with the original enumeration n_0, n_1, \dots allows to identify the candidate c . The number j can be obtained from the output with the help of unique choice $\text{UC}_{\mathbb{N}}$. \square

In case of Baire space we formulate the following conjecture.

Conjecture 6.3 $\text{UC}_{\mathbb{N}^{\mathbb{N}}} \equiv_{\text{W}} \text{C}_{\mathbb{N}^{\mathbb{N}}}$.

On the other hand, unique choice $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ is also not too simple. One can easily see that $\lim \leq_{\text{W}} \text{UC}_{\mathbb{N}^{\mathbb{N}}}$ holds for the limit map

$$\lim : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}, \langle p_0, p_1, p_2, \dots \rangle \mapsto \lim_{i \rightarrow \infty} p_i$$

and with the methods of the next section it also follows that the cone below $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ is closed under composition. Hence, $\text{UC}_{\mathbb{N}^{\mathbb{N}}}$ cannot be located on any finite level of the Borel hierarchy. This can also be deduced from the fact that there are co-c.e. closed singletons $\{p\} \subseteq \mathbb{N}^{\mathbb{N}}$ such that p is hyperarithmetical, but not arithmetical (see Propositions 1.8.62 and 1.8.70 in [23]). We obtain the following corollary as a direct consequence of Corollaries 5.2 and 5.3 and the previous proposition and the observation that $\text{UC}_{\mathbb{N}} \leq_{\text{W}} \text{UC}_{\mathbb{R}}$ holds.

Corollary 6.4 $\text{UC}_{\{0,1\}^{\mathbb{N}}} \equiv_{\text{W}} \text{C}_{\{0\}} \equiv_{\text{W}} \text{id}$ and $\text{UC}_{\mathbb{R}} \equiv_{\text{W}} \text{C}_{\mathbb{N}}$.

We will use the inversion and the graph map as follows

- $\text{Inv}_{X,Y} : \subseteq \mathcal{C}(X, Y) \times Y \rightarrow X, (f, y) \mapsto f^{-1}(y)$, where $\text{dom}(\text{Inv}_{X,Y}) := \{(f, y) : f \text{ injective and } y \in \text{dom}(f^{-1})\}$
- $\text{graph}_{X,Y} : \mathcal{C}(X, Y) \rightarrow \mathcal{A}_-(X \times Y), f \mapsto \text{graph}(f)$, where $\text{graph}(f) := \{(x, y) \in X \times Y : f(x) = y\}$.

For computable metric spaces X and Y the map $\text{graph}_{X,Y}$ is known to be computable (see [5]). It turns out that the map $\text{Inv}_{X,Y}$ is reducible to unique choice of X .

Theorem 6.5 (Inversion operator) *Let X and Y be computable metric spaces. Then $\text{Inv}_{X,Y} \leq_{\text{W}} \text{UC}_X$.*

Proof. In [5] we have established the formula

$$f^{-1}(y) = \text{in}_X^{-1} \circ \text{sec}(\text{graph}_{X,Y}(f), y),$$

where $\text{sec} : \mathcal{A}_-(X \times Y) \times Y \rightarrow X, (A, y) \mapsto A_y := \{x \in X : (x, y) \in A\}$ is the computable section map (see [5]). Altogether, this shows $\text{Inv}_{X,Y} \leq_{\text{W}} \text{in}_X^{-1} = \text{UC}_X$. \square

As a corollary we get that in particular any specific inverse of a computable map is reducible to unique choice. We can generalize this non-uniform result

even to the case of non-injective maps and ordinary choice. We note that the inverse $f^{-1} : \subseteq Y \rightrightarrows X, y \mapsto f^{-1}\{y\}$ exists as a multi-valued map for any single-valued $f : X \rightarrow Y$.

Theorem 6.6 (Inversion) *Let X and Y be computable metric spaces. If $f : X \rightarrow Y$ is computable, then $f^{-1} \leq_W \mathbf{C}_X$ and if f is also injective, then $f^{-1} \leq_W \mathbf{UC}_X$.*

Proof. If $f : X \rightarrow Y$ is computable, then $F : \mathcal{A}_-(Y) \rightarrow \mathcal{A}_-(X), A \mapsto f^{-1}(A)$ is computable too and we obtain

$$f^{-1}(y) = \mathbf{C}_X \circ F \circ \text{in}_Y(y),$$

i.e. $f^{-1} \leq \mathbf{C}_X$ and if f is also injective, then we obtain $f^{-1}(y) = \text{in}_X^{-1} \circ F \circ \text{in}_Y(y)$, i.e. $f^{-1} \leq_W \mathbf{UC}_X$. \square

We mention that a multi-valued function f on represented spaces is called *weakly computable*, if $f \leq_W \mathbf{C}_{\{0,1\}^{\mathbb{N}}}$ and f is called *computable with finitely many mind changes* if it can be computed on a Turing machine that revises its output at most finitely many times for each particular input. In Theorem 7.11 we will show that the latter is equivalent to $f \leq_W \mathbf{C}_{\mathbb{N}}$. We get the following result as a corollary of Theorem 6.6 and Corollary 6.4.

Corollary 6.7 (Compact inversion) *Let X and Y be computable metric spaces and let X be computably compact. If $f : X \rightarrow Y$ is computable, then f^{-1} is weakly computable, if f is also injective, then f^{-1} is even computable.*

The second part of the statement was known as such (see, for instance, [5]). The following corollary is also a consequence of Theorem 6.6 and Corollary 6.4.

Corollary 6.8 (Locally compact inversion) *Let X be a computable K_σ -space and let Y be a computable metric space. If $f : X \rightarrow Y$ is computable, then $f^{-1} \leq_W \mathbf{C}_{\mathbb{R}}$ and if f is also injective then $f^{-1} \leq_W \mathbf{C}_{\mathbb{N}}$, hence f^{-1} is computable with finitely many mind changes.*

These results are not necessarily optimal. For instance, it is known that the inverse of an injective computable map $f : \mathbb{R} \rightarrow \mathbb{R}$ is even computable. However, for this result one has to exploit additional properties of \mathbb{R} , such as connectedness properties (see [5]). We give some example that shows that the inversion results do not hold true for arbitrary represented spaces. By $\mathbb{R}_<$ we denote the set of real numbers equipped with the left cut representation $\rho_<$,

which represents a real number x by an enumeration of all rational numbers $q < x$ (see [36]).

Example 6.9 *Let \mathbb{R} denote the real number represented with the Cauchy representation and let $\mathbb{R}_{<}$ denote the real number denoted with the left cut representation. The identity $f : \mathbb{R} \rightarrow \mathbb{R}_{<}, x \mapsto x$ is computable and its inverse $f^{-1} : \mathbb{R}_{<} \rightarrow \mathbb{R}$ is known to be equivalent to lim (see Proposition 3.7 in [6] and Exercise 8.2.12 in [36]). In particular, f^{-1} is not reducible to $\mathbb{C}_{\mathbb{R}}$.*

7 Choice on Baire Space and Non-Deterministic Computability

In this section we will compare the power of choice for certain spaces with models of hypercomputation that have been considered. This approach to classify models of hypercomputation in terms of Weihrauch reducibility has been started in [27]. Here, the relevant models of hypercomputation are non-deterministically computable functions and functions computable with revising computations in the sense of Martin Ziegler [38,39]. The latter ones are also known as functions computable with finitely many mind-changes, for instance in learning theory [11,12].

In [38] Martin Ziegler has introduced a concept of non-deterministically computable functions. We generalize this concept to advice spaces that are subsets of Baire space and we prove that this concept can be characterized by choice for the advice space. This characterization yields some interesting consequences.

Definition 7.1 (Non-deterministic computability) Let $(X, \delta_X), (Y, \delta_Y)$ be represented spaces and let $A \subseteq \mathbb{N}^{\mathbb{N}}$. A function $f : \subseteq X \rightrightarrows Y$ is said to be *non-deterministically computable with advice space A* , if there exist two computable functions $F_1, F_2 : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $\langle \text{dom}(f\delta_X) \times A \rangle \subseteq \text{dom}(F_2)$ and for each $p \in \text{dom}(f\delta_X)$ the following hold:

- (1) $(\exists r \in A) \delta_{\mathbb{S}} F_2 \langle p, r \rangle = 0$,
- (2) $(\forall r \in A) (\delta_{\mathbb{S}} F_2 \langle p, r \rangle = 0 \implies \delta_Y F_1 \langle p, r \rangle \in f\delta_X(p))$.

Here $A \subseteq \mathbb{N}^{\mathbb{N}}$ is considered as subspace of Baire space. Intuitively, the set A serves as a set of possible advices that can give extra support to the computation. Any computation can be successful or it can fail, which is indicated by the output of F_2 (where “1” means the advice is recognized to fail after finite time and “0” means the advice is successful in the long run). That is F_2 can be considered as a realizer of a function $f_2 : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{S}$. The set

$$A_p := \{r \in A : \delta_{\mathbb{S}} F_2 \langle p, r \rangle = 0\} = \{r \in A : f_2 \langle p, r \rangle = 0\}$$

is the set of successful advices for input $p \in \text{dom}(f\delta_X)$. Intuitively, F_2 is a method to recognize unsuccessful advices and F_1 is a method to determine the output of the computation for successful advices. The two conditions then express intuitively that for each fixed admissible input the following hold:

- (1) There exists a successful advice for this input.
- (2) Each successful advice produces a correct output.

Functions that are non-deterministically computable in the sense of [38] are non-deterministically computable with full Baire space $\mathbb{N}^{\mathbb{N}}$ as advice space². Now we can prove the following equivalence.

Theorem 7.2 (Non-deterministic computability) *Let X and Y be represented spaces, $A \subseteq \mathbb{N}^{\mathbb{N}}$ and let $f : \subseteq X \rightrightarrows Y$ be a multi-valued function. Then the following are equivalent:*

- (1) $f \leq_W \mathbf{C}_A$,
- (2) f is non-deterministically computable with advice space A .

Proof. We consider the represented spaces (X, δ_X) and (Y, δ_Y) . Let f be non-deterministically computable with advice space A . Then there are computable functions F_1, F_2 according to Definition 7.1. By type conversion and since $\langle \text{dom}(f\delta_X) \times A \rangle \subseteq \text{dom}(F_2)$ we can transfer F_2 into a computable function

$$h : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathcal{C}(A, \mathbb{S}), p \mapsto (r \mapsto \delta_{\mathbb{S}} F_2 \langle p, r \rangle).$$

Hence, for each $p \in \text{dom}(f\delta_X)$ the function $h(p) = \chi_{A \setminus A_p}$ is a characteristic function of the closed set $A_p \in \mathcal{A}_-(A)$ of successful advices. Here h can also be considered as computable function of type $h : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathcal{A}_-(A), p \mapsto A_p$. By condition (1) of Definition 7.1 we obtain that $A_p \neq \emptyset$ for any $p \in \text{dom}(f\delta_X)$ and by condition (2) we obtain $\delta_Y F_1 \langle p, \mathbf{C}_A h(p) \rangle \subseteq f\delta_X(p)$. Let H be a computable realizer of h . Then $F_1 \langle \text{id}, GH \rangle$ is a realizer of f for any realizer G of \mathbf{C}_A and hence $f \leq_W \mathbf{C}_A$.

On the other hand, let $f \leq_W \mathbf{C}_A$. Then any realizer G of \mathbf{C}_A computes some realizer F of f , i.e. there are computable functions H, K such that for all realizers G of \mathbf{C}_A there is some realizer F of f such that $F(p) = K \langle p, GH(p) \rangle$ for all $p \in \text{dom}(f\delta_X)$. Now we describe maps $F_1, F_2 : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ that satisfy the conditions of Definition 7.1 for f . For each $p \in \text{dom}(f\delta_X)$ the function H computes a non-empty set $A_p = \psi_-^A H(p)$ and by evaluation there exists a computable function F_2 such that $\delta_{\mathbb{S}} F_2 \langle p, r \rangle = \chi_{A \setminus A_p}(r)$ for all $p \in \text{dom}(f\delta_X)$ and $r \in A$. That means to choose A_p as the set of successful advices. We can

² The advice space is not made explicit in [38], but we conclude implicitly that the advice space $\mathbb{N}^{\mathbb{N}}$ is meant.

also choose $F_1 := K$ and verify the conditions (1) and (2) of Definition 7.1. Firstly, it is clear that $A_p \neq \emptyset$ for all $p \in \text{dom}(f\delta_X)$ and hence F_2 satisfies condition (1). Secondly, for each $r \in A_p$ there is a realizer G of \mathbf{C}_A such that $GH(p) = r$ and hence we obtain $F_1\langle p, r \rangle = K\langle p, GH(p) \rangle = F(p)$ for a realizer F of f . This implies $\delta_Y F_1\langle p, r \rangle \in f\delta_X(p)$ and hence condition (2) holds as well. Altogether f is non-deterministically computable with advice space A . \square

The main benefit of this characterization of closed choice is that using it we can easily prove the following theorem that shows that the advice for compositions can be determined a priori and independently. We note that due to the fact that Baire space admits a computable and bijective pairing function, we can always consider $A \times B$ as subspace of Baire space for any two subspaces A, B of Baire space.

Theorem 7.3 (Independent Choice) *Let $A, B \subseteq \mathbb{N}^{\mathbb{N}}$ and let f and g be multi-valued functions on represented spaces. If $f \leq_W \mathbf{C}_A$ and $g \leq_W \mathbf{C}_B$, then $f \circ g \leq_W \mathbf{C}_{A \times B}$.*

Proof. We consider represented spaces (X, δ_X) , (Y, δ_Y) and (Z, δ_Z) . Let now $f : \subseteq Y \rightrightarrows Z$ and $g : \subseteq X \rightrightarrows Y$ be non-deterministically computable with advice spaces A and B , respectively. Due to Theorem 7.2 it suffices to show that $f \circ g$ is non-deterministically computable with advice space $A \times B$. Intuitively, we can choose an advice $(r, s) \in A \times B$ and use advice r for f and advice s for g . More precisely, let f and g be non-deterministically computable using computable functions F_1, F_2 and G_1, G_2 according to Definition 7.1, respectively. We define H_1 and H_2 that witness non-deterministic computability of $f \circ g$ with advice space $A \times B$. We can define a computable H_1 by

$$H_1\langle p, \langle r, s \rangle \rangle := F_1\langle G_1\langle p, s \rangle, r \rangle$$

and there exists a computable H_2 such that

$$\delta_{\mathbb{S}} H_2\langle p, \langle r, s \rangle \rangle = \begin{cases} 1 & \text{if } \delta_{\mathbb{S}} G_2\langle p, s \rangle = 1 \\ \delta_{\mathbb{S}} F_2\langle G_1\langle p, s \rangle, r \rangle & \text{otherwise} \end{cases}$$

for all $p \in \text{dom}(fg\delta_X)$ and all $(r, s) \in A \times B$. Such a computable H_2 exists, since $\delta_{\mathbb{S}} G_2\langle p, s \rangle = 0$ implies that $\delta_Y G_1\langle p, s \rangle \in g(\delta_X(p)) \subseteq \text{dom}(f)$. Now we verify that H_1 and H_2 satisfy conditions (1) and (2) of Definition 7.1 for $f \circ g$. To this end, let $p \in \text{dom}(fg\delta_X)$.

By condition (1) for g there is an $s \in B$ such that $\delta_{\mathbb{S}} G_2\langle p, s \rangle = 0$ and hence by condition (2) for g we obtain $\delta_Y G_1\langle p, s \rangle \in \text{dom}(f)$. Hence by condition (1) for

f there is an $r \in A$ such that $\delta_{\mathbb{S}}F_2\langle G_1\langle p, s \rangle, r \rangle = 0$ and thus $\delta_{\mathbb{S}}H_2\langle p, \langle r, s \rangle \rangle = 0$, which shows that condition (1) also holds for fg .

Now let $(r, s) \in A \times B$ be such that $\delta_{\mathbb{S}}H_2\langle p, \langle r, s \rangle \rangle = 0$. Then $\delta_{\mathbb{S}}G_2\langle p, s \rangle = 0$ and $\delta_{\mathbb{S}}F_2\langle G_1\langle p, s \rangle, r \rangle = 0$. Hence by conditions (2) for g and f we obtain $\delta_Y G_1\langle p, s \rangle \in g\delta_X(p)$ and hence $\delta_Z F_1\langle G_1\langle p, s \rangle, r \rangle \in fg\delta_X(p)$, which proves condition (2) for fg . \square

We recall that we call a multi-valued function h on represented spaces *closed under composition* if the principal ideal of h is closed under composition, i.e. if $f \leq_W h$ and $g \leq_W h$ implies $f \circ g \leq_W h$ (for f and g of appropriate type). It is worth pointing out that closure under composition entails idempotency.

Proposition 7.4 *Every multi-valued function f on represented spaces that is closed under composition is also idempotent.*

Proof. Let $f : \subseteq X \rightrightarrows Y$ be a multi-valued function on represented spaces. Then we have $f \times f = (f \times \text{id}_Y) \circ (\text{id}_X \times f)$ and $f \times \text{id}_Y \leq_W f$ and $\text{id}_X \times f \leq_W f$. That is, if f is closed under composition, then $f \times f \leq_W f$. \square

We get the following consequence of Theorem 7.3, which is a strengthening of Corollary 3.8 for $A \subseteq \mathbb{N}^{\mathbb{N}}$.

Corollary 7.5 (Closure under composition) *Let $A \subseteq \mathbb{N}^{\mathbb{N}}$ be a subspace of Baire space. If there is a computable surjection $s : A \rightarrow A^2$, then $\mathbf{C}_{A \times A} \leq_W \mathbf{C}_A$ and hence \mathbf{C}_A is closed under composition and idempotent.*

In particular, we can apply this result in the following cases.

Corollary 7.6 *The choice functions $\mathbf{C}_{\mathbb{N}}, \mathbf{C}_{\{0,1\}^{\mathbb{N}}}, \mathbf{C}_{\mathbb{N}^{\mathbb{N}}}, \mathbf{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}$ and hence $\mathbf{C}_{\mathbb{R}}$ are closed under composition and idempotent.*

For most of these functions this was known. However, the proofs in [14] and [7] for the case $\mathbf{C}_{\{0,1\}^{\mathbb{N}}}$ are considerably more difficult whereas the Independent Choice Theorem 7.3 has a simple proof and covers many cases simultaneously. The results for $\mathbf{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}$ and $\mathbf{C}_{\mathbb{R}}$ seem to be new and are of independent interest. Closure of non-deterministically computable functions for advice space $\mathbb{N}^{\mathbb{N}}$ was observed in [38].

Now we want to prove that the class of (single-valued) functions below choice for Baire space $\mathbf{C}_{\mathbb{N}^{\mathbb{N}}}$ is essentially the class of effectively Borel measurable functions. It is known that there is no complete Borel measurable function, since any particular function has to be Σ_{ξ}^0 -measurable in the Borel hierarchy

for some countable ordinal ξ (see 1G.15 in [22]). Nevertheless, we will see that choice of Baire space $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ is complete for Borel measurable functions in a certain sense. We will say that a function $f : X \rightarrow Y$ on computable Polish spaces X and Y is *effectively Borel measurable*, if its graph is an effective Σ_1^1 -set (see Theorem 3E.5 in [22]). Here a subset $A \subseteq X$ of a computable Polish space X is called *effective Σ_1^1 -set*, if there exists a co-c.e. closed set $B \subseteq X \times \mathbb{N}^{\mathbb{N}}$ such that $x \in A \iff (\exists p \in \mathbb{N}^{\mathbb{N}})(x, p) \in B$. We will use once again Theorem 7.2 for the proof.

Theorem 7.7 (Choice of Baire space) *Let X and Y be computable Polish spaces and let $f : X \rightarrow Y$ be a function. Then the following are equivalent:*

- (1) $f \leq_W \mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$,
- (2) f is effectively Borel measurable.

Proof. By Theorem 7.2 it suffices to show that f is non-deterministically computable with advice space $\mathbb{N}^{\mathbb{N}}$ if and only if it is effectively Borel measurable. Since X and Y are Polish, we can assume that we have total computably admissible representations δ_X and δ_Y for X and Y , respectively (see, for instance, Corollary 4.4.12 in [2]).

If f is non-deterministically computable with advice space $\mathbb{N}^{\mathbb{N}}$, then there are computable functions F_1, F_2 according to Definition 7.1. We obtain for all $(x, y) \in X \times Y$

$$\begin{aligned} f(x) = y \\ \iff (\exists \langle p, r \rangle \in \mathbb{N}^{\mathbb{N}})(\delta_X(p) = x, \delta_S F_2 \langle p, r \rangle = 0 \text{ and } \delta_Y F_1 \langle p, r \rangle = y). \end{aligned}$$

Since all involved functions in the matrix of the formula are computable and total, it follows that the matrix constitutes a co-c.e. closed subset of $X \times Y \times \mathbb{N}^{\mathbb{N}}$ in the parameters $(x, y, \langle p, r \rangle)$. Hence f is effectively Borel measurable.

Let now f be an effectively Borel measurable function. Then $\text{graph}(f)$ is a Σ_1^1 -set in the effective Borel hierarchy and there exists a co-c.e. closed set $A \subseteq X \times Y \times \mathbb{N}^{\mathbb{N}}$ such that

$$f(x) = y \iff (\exists r \in \mathbb{N}^{\mathbb{N}})(x, y, r) \in A.$$

We devise a non-deterministic computation for f , by defining suitable computable functions F_1, F_2 according to Definition 7.1. Firstly, there exists a computable function F_2 with $\delta_S F_2 \langle p, \langle q, r \rangle \rangle = \chi_{A^c}(\delta_X(p), \delta_Y(q), r)$ and we define $F_1 \langle p, \langle q, r \rangle \rangle := q$. Then F_1 is computable too and we obtain

$$(\exists r \in \mathbb{N}^{\mathbb{N}}) \delta_S F_2 \langle p, \langle q, r \rangle \rangle = 0 \iff (\exists r \in \mathbb{N}^{\mathbb{N}})(\delta_X(p), \delta_Y(q), r) \in A$$

$$\iff f\delta_X(p) = \delta_Y(q)$$

and if this condition holds, then we have $\delta_Y F_1 \langle p, \langle q, r \rangle \rangle = \delta_Y(q) = f\delta_X(p)$. Altogether, this shows that F_1, F_2 satisfy the conditions of Definition 7.1. \square

We note that $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ itself is not Borel measurable, which is not a contradiction, since it is not a single-valued function defined on a Polish space. In contrast, the domain of $\mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ corresponds to the set of ill-founded trees (i.e. trees with at least one infinite branch), which is known to be Σ_1^1 -complete (see Theorem 27.1 in [18]). We mention that the relativized version of the above proof leads to the following corollary.

Corollary 7.8 *Let X and Y be Polish spaces represented by their Cauchy representations and let $f : X \rightarrow Y$ be a function. Then the following are equivalent:*

- (1) $f \leq_W \mathbb{C}_{\mathbb{N}^{\mathbb{N}}}$ with respect to some oracle,
- (2) f is Borel measurable.

Here, reducibility “with respect to some oracle” is equivalent to using the continuous version of Weihrauch reducibility. Now we will consider another model of hypercomputation, namely finitely revising computation as considered in [39] and as known as computation with finitely many mind changes in learning theory [11]. A Turing machine that computes with finitely many mind changes or that is finitely revising can erase its output tape at any stage during its computation and start writing anew, however, this can be done only finitely often, ensuring that the output is well-defined. In [39], the power of finite revising was characterized in terms of an operator mapping one representation into another. We will define this concept here using the discrete limit $\lim_{\Delta} : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}, \langle p_0, p_1, \dots \rangle \mapsto \lim_{i \rightarrow \infty} p_i$ where the Δ stands for the discrete topology on $\mathbb{N}^{\mathbb{N}}$ and the limit on the right-hand side is taken with respect to this topology. That is a sequence $(p_i)_{i \in \mathbb{N}}$ converges with respect to Δ if and only if it is eventually constant. Now we use the discrete limit to define a discrete version of the jump of a representation (as equivalently considered in [39]).

Definition 7.9 (Discrete jump) Let (X, δ) be a represented space. Then we define the *discrete jump* of δ by $\delta^{\Delta} := \delta \circ \lim_{\Delta}$.

It is easy to see that the following result holds (cf. Lemma 3.7 in [39]).

Proposition 7.10 (Computability with finitely many mind changes) *Let (X, δ_X) and (Y, δ_Y) be represented spaces and let $f : \subseteq X \rightrightarrows Y$ be a multi-*

valued function. Then the following are equivalent:

- (1) f is (δ_X, δ_Y) -computable with finitely many mind changes,
- (2) f is $(\delta_X, \delta_Y^\Delta)$ -computable,
- (3) f is $(\delta_X^\Delta, \delta_Y^\Delta)$ -computable.

With this proposition we can produce the following characterization of the discrete limit and the power of computations with finitely many mind changes in terms of closed choice, showing that finite revision allows exactly to perform closed choice in \mathbb{N} .

Theorem 7.11 (Choice on natural numbers) *Let f be a multi-valued function on represented spaces. Then the following are equivalent:*

- (1) $f \leq_W \mathbf{C}_{\mathbb{N}}$,
- (2) $f \leq_W \lim_{\Delta}$,
- (3) f is computable with finitely many mind changes.

Proof. It is easy to see that $\mathbf{C}_{\mathbb{N}}$ is computable with finitely many mind changes. Starting with $n = 0$, the machine outputs a $\delta_{\mathbb{N}}$ name for n and searches for n in the input at the same time. If the search is successful, the output is erased, n is increased by 1, and the machine starts again. A valid input never causes the machine to erase its output tape infinitely often, and an output can only avoid erasion, if it is a valid result for $\mathbf{C}_{\mathbb{N}}$. Moreover, being computable with finitely many mind changes is preserved downwards by Weihrauch reducibility (see Lemma 4.4 in [6]) and hence $f \leq_W \mathbf{C}_{\mathbb{N}}$ implies that f is computable with finitely many mind changes. Hence (1) implies (3).

Now we assume that f is of type $f : \subseteq X \rightrightarrows Y$ for represented spaces (X, δ_X) and (Y, δ_Y) . If f is computable with finitely many mind changes, then f has a computable $(\delta_X, \delta_Y^\Delta)$ -realizer F by Proposition 7.10, which means $\delta_Y \circ \lim_{\Delta} F(p) \in f\delta_X(p)$ for all $p \in \text{dom}(\delta_X)$. Hence $f \leq_W \lim_{\Delta}$ and (3) implies (2).

In order to prove that (2) implies (1) it suffices to show $\lim_{\Delta} \leq_W \mathbf{C}_{\mathbb{N}}$. we describe a machine computing a function G in the following: The input for G is a sequence $(p_n)_{n \in \mathbb{N}}$ with $p_n \in \mathbb{N}^{\mathbb{N}}$. Now we start to test simultaneously $p_n = p_{n+j}$ for each $n, j \in \mathbb{N}$. If a contradiction is found, we print n on the output tape. If $\mathbf{C}_{\mathbb{N}}$ is applied to the output of G , the answer is an index n_0 , so that the initial sequence is constant after n_0 . The remaining task is to output the n_0 th entry of the sequence. \square

We get the following corollary that shows that the discrete limit is equivalent to choice on natural numbers.

Corollary 7.12 $\text{lim}_\Delta \equiv_{\text{w}} \mathbb{C}_{\mathbb{N}}$.

We close by mentioning that the class of operations characterized by choice on Cantor space is also of independent interest. These functions have been called *weakly computable* in [7] and basically the equivalence of (1) and (3) below is the definition. With Theorem 7.2 we get a characterization of weakly computable functions as non-deterministically computable ones with advice space $\{0, 1\}^{\mathbb{N}}$.

Corollary 7.13 (Choice on Cantor space) *Let f be a multi-valued function on represented spaces. Then the following are equivalent:*

- (1) $f \leq_{\text{w}} \mathbb{C}_{\{0,1\}^{\mathbb{N}}}$,
- (2) f is non-deterministically computable with advice space $\{0, 1\}^{\mathbb{N}}$,
- (3) f is weakly computable.

A surprising omission in our list of classes of computable functions characterized by closed choice of some space is the class of limit computable functions. In light of Corollary 7.5 it seems that choice for most natural spaces will correspond to classes of functions that are closed under composition, whereas the class of limit computable functions is not closed under composition (see for instance [3]). Thus, the following conjecture is plausible.

Conjecture 7.14 *There is no represented space (X, δ) such that $\mathbb{C}_X \equiv_{\text{w}} \text{lim}$.*

At least for Polish spaces (X, δ) this conjecture follows topologically from Corollary 4.10. The closest we can get to a characterization of limit computable functions by a choice principle of a Polish space is expressed in the following result.

Corollary 7.15 (Parallelized choice on natural numbers) *Let f be a multi-valued function on represented spaces. Then the following are equivalent:*

- (1) $f \leq_{\text{w}} \widehat{\mathbb{C}}_{\mathbb{N}}$,
- (2) f is limit computable.

This follows from $\widehat{\mathbb{C}}_{\mathbb{N}} \equiv \text{lim}$ (see Example 3.10) and the fact that lim is complete for limit computable functions, see for instance [3].

8 A Uniform Low Basis Theorem

The choice of Cantor space $C_{\{0,1\}^{\mathbb{N}}}$ is known to be even not non-uniformly computable, since there is a co-c.e. closed set $A \subseteq \{0,1\}^{\mathbb{N}}$ that has no computable points (this can be seen, for instance, using the Kleene tree [19] or Proposition V.5.25 in [24]). However, by the Low Basis Theorem of Jockusch and Soare (see Theorem 2.1 in [16] or Proposition V.5.27 in [24]) any co-c.e. closed set $A \subseteq \{0,1\}^{\mathbb{N}}$ has a low point, that is for computable w , the set $\psi_{\{0,1\}^{\mathbb{N}}}^{\{0,1\}^{\mathbb{N}}}(w)$ always contains a low point. As shown in [6], this carries over to all problems below $C_{\mathbb{R}}$: For every computable instance, there is a solution that is low. We will demonstrate that this result even holds uniformly, after some necessary definitions have been introduced.

Definition 8.1 (Turing jump operator) Let $(U_n)_{n \in \mathbb{N}}$ be a standard enumeration of the c.e. open subsets of Baire space $\mathbb{N}^{\mathbb{N}}$. Define the *jump operator* $J : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ by:

$$J(p)(n) = \begin{cases} 1 & \text{if } p \in U_n \\ 0 & \text{otherwise} \end{cases}$$

Contrary to its behavior on Turing degrees, as a function on Baire space, the jump is injective. It even admits a computable inverse J^{-1} . In [4], for any representation δ of some set X , a representation $f \delta$ is defined by $(f \delta)(p) = \delta(J^{-1}(p))$. Together with the operator $'$ studied in [39], where a representation δ' is defined by $\delta'(p) = \delta(\lim p)$, f forms a Galois connection, as shown in [4]. We define the *low representation* $\delta^{\vee} := (f \delta)'$ for any represented space (X, δ) and if $f : \subseteq X \rightrightarrows Y$ is a multi-valued map on represented spaces (X, δ_X) and (Y, δ_Y) , then f is called *low computable*, if f is $(\delta_X, \delta_Y^{\vee})$ -computable. In particular, we will be interested in the low representation $\delta_{\{0,1\}^{\mathbb{N}}}^{\vee} = \text{id}_{\{0,1\}^{\mathbb{N}}} \circ J^{-1} \circ \text{lim}$ of Cantor space and the low representation of Baire space $\delta_{\mathbb{N}^{\mathbb{N}}}^{\vee} = J^{-1} \circ \text{lim}$, which we also denote by \mathfrak{L} .

Lemma 8.2 (Low points) *A sequence $p \in \mathbb{N}^{\mathbb{N}}$ is low if and only if it has a computable \mathfrak{L} -name.*

Proof. By definition, a sequence $p \in \mathbb{N}^{\mathbb{N}}$ is called low, if its Turing jump is Turing reducible to the halting problem, which is equivalent to $J(p)$ being in the class Δ_2^0 of the arithmetical hierarchy (see Proposition IV.1.16 in [24]). By Shoenfield's Limit Lemma (see Proposition IV.1.17 in [24]), $J(p) \in \Delta_2^0$ if and only if there exists a computable sequence $q = \langle q_0, q_1, \dots \rangle \in \mathbb{N}^{\mathbb{N}}$ such that $J(p) = \lim_{i \rightarrow \infty} q_i = \text{lim}(q)$, i.e. if and only if $p = \mathfrak{L}(q)$. \square

Analogously, $p \in \{0, 1\}^{\mathbb{N}}$ is low if and only if it has a computable $\delta_{\{0,1\}^{\mathbb{N}}}^{\vee}$ -name. Now we can formulate and prove our uniform low basis theorem, which states that, given an enumeration of the complement of a non-empty compact subset A of $\{0, 1\}^{\mathbb{N}}$, we can compute a sequence converging to the jump of a point $p \in A$.

Theorem 8.3 (Uniform Low Basis Theorem) $C_{\{0,1\}^{\mathbb{N}}}$ is low computable.

Proof. We describe a machine M that given a $\psi_{-}^{\{0,1\}^{\mathbb{N}}}$ -name of a compact set $A \subseteq \{0, 1\}^{\mathbb{N}}$ produces a sequence $(p_m)_{m \in \mathbb{N}}$ converging to a $\int \text{id}_{\{0,1\}^{\mathbb{N}}}$ -name of some element of A . The input of M is a list enumerating basic open sets exhausting $\{0, 1\}^{\mathbb{N}} \setminus A$. The complement of the union of the first m of these subsets shall be denoted A^m . Likewise, for each $n \in \mathbb{N}$, we let U_n^m be the union of the first m basic open subsets exhausting U_n . Here, for simplicity, $(U_n)_{n \in \mathbb{N}}$ is supposed to be a standard enumeration of the c.e. open subsets of Cantor space $\{0, 1\}^{\mathbb{N}}$ and the aforementioned results on the jump and integral are used analogously for Cantor space.

The computation of each p_m can be considered independently, and proceeds as follows. For each $n \in \mathbb{N}$, the machine M performs the following tests³ in the given order:

- (1) Does $A^m \subseteq U_n^m$ hold? If the answer is YES, the n th bit of p_m is 1.
- (2) Let K be the set of indexes $i < n$, so that the i th bit of p_m is 0. Test $A^m \subseteq U_n^m \cup \bigcup_{i \in K} U_i^m$. If the answer is YES, the n th bit of p_m is 1.
- (3) Otherwise, the n th bit of p_m is 0.

All operations are performed on a finite set of basic open sets, either obtained from the input, or computable by definition. Therefore, each test is decidable. We will first prove that the p_m converge as m goes to infinity. This is equivalent to showing that each bit of the p_m changes only finitely many times.

The first test is monotone in m , as we have $A^{m+1} \subseteq A^m$ and $U_n^m \subseteq U_n^{m+1}$. Thus, if for some m the n th bit of p_m was set to 1 due to the first test, the n th bit of all $p_{m'}$ for $m' > m$ is 1, too.

Now consider the second test, and assume that all bits i with $i < n$ remain unchanged. Then, again by the same argument, once the second test yields YES for some m , it will do so for all larger m' as well. The only way for the second test to change the corresponding bit from 1 to 0 is if some smaller bit has been set from 0 to 1 previously.

³ Of course the first test could be subsumed by the second one; however, since their interpretation is different, we prefer to mention the first test separately.

An inductive argument concludes the proof of convergence: The first bit can change at most once, from 0 to 1. All other bits n can change at most once *for each given configuration of the lower bits*. If only finitely many changes of the bits smaller than n are possible, then there will be only finitely many changes of the n th bit.

It remains to show that the p_m actually converge to a correct output w . Basically, the first test ensures that the limit sequence w specifies a point $x \in A$, while the second test ensures that w is a valid $f \text{id}_{\{0,1\}^{\mathbb{N}}}$ -name, i.e. $w \in \text{dom}(J^{-1})$, in the first instance.

To elaborate this, assume $A \subseteq U_n$ for some $n \in \mathbb{N}$. Then for every $f \text{id}_{\{0,1\}^{\mathbb{N}}}$ -name w with $J^{-1}(w) \in A$ obviously $w(n) = 1$ has to be true. On the other hand, for $x \notin A$, there is some neighborhood U of A with $x \notin U$. It is possible to choose U as c.e. open (for instance by choosing the complement of some sufficiently small clopen basic neighborhood of x), thus, there is an $n \in \mathbb{N}$ with $A \subseteq U_n$, but $x \notin U_n$. Thus, having $w(n) = 1$ for each $n \in \mathbb{N}$ with $A \subseteq U_n$ for a $(f \text{id})$ -name w is both necessary and sufficient to ensure $J^{-1}(w) \in A$.

In the next step, we have to show that $A \subseteq U_n$ already guarantees the existence of an $m \in \mathbb{N}$ with $A^m \subseteq U_n^m$. The other direction is trivial. As $A \subseteq U_n$ is equivalent to $A^c \cup U_n = \{0,1\}^{\mathbb{N}}$, the basic open sets exhausting A^c and U_n are an open cover of $\{0,1\}^{\mathbb{N}}$. Since $\{0,1\}^{\mathbb{N}}$ is compact, there has to be a finite subcover. Thus, there is some $m \in \mathbb{N}$, so that the first m basic open sets in the ψ_- -name of A together with the first m basic open sets listed for U_n already cover $\{0,1\}^{\mathbb{N}}$, that is fulfills $A^m \subseteq U_n^m$, which concludes this part of the proof.

Now we have to show that the second test ensures that the limit sequence w is in the domain of $f \text{id}_{\{0,1\}^{\mathbb{N}}}$. This amounts to proving

$$\left(\bigcap_{i \in \mathbb{N}, w(i)=1} U_i \right) \setminus \left(\bigcup_{j \in \mathbb{N}, w(j) \neq 1} U_j \right) \neq \emptyset.$$

We note that this difference is automatically a singleton, if non-empty, since any two distinct points can be separated by two c.e. open sets. We will use the abbreviations $X := \{i \in \mathbb{N} \mid w(i) = 1 \text{ due to the first test}\}$, $Y := \{i \in \mathbb{N} \mid w(i) = 1 \text{ due to the second test}\}$ and $Z := \{i \in \mathbb{N} \mid w(i) = 0\}$, and $U_i^c := \{0,1\}^{\mathbb{N}} \setminus U_i$. With this, we have to show:

$$\left(\bigcap_{i \in X} U_i \right) \cap \left(\bigcap_{j \in Y} U_j \right) \cap \left(\bigcap_{k \in Z} U_k^c \right) \neq \emptyset.$$

Taking into consideration our results on the first test, this simplifies to:

$$A \cap \left(\bigcap_{j \in Y} U_j \right) \cap \left(\bigcap_{k \in Z} U_k^c \right) \neq \emptyset.$$

Assume that already $A \cap \left(\bigcap_{k \in Z} U_k^c \right) = \emptyset$ would hold. By the finite intersection property in compact spaces, this implies the existence of a (smallest) $k_0 \in \mathbb{N}$ with $A \cap \left(\bigcap_{k \in Z, k \leq k_0} U_k^c \right) = \emptyset$. Rearranging the expression yields $A \subseteq U_{k_0} \cup \bigcup_{k \in Z, k < k_0} U_k$, so the second test would have been triggered for k_0 , so $k_0 \notin Z$ follows. This contradicts the assumption, so we have $A \cap \left(\bigcap_{k \in Z} U_k^c \right) \neq \emptyset$.

Now we choose some $x \in A \cap \left(\bigcap_{k \in Z} U_k^c \right)$. Assume $x \notin \bigcap_{j \in Y} U_j$. There has to be some $j_0 \in Y$ with $x \notin U_{j_0}$. Now $j_0 \in Y$ implies $A \subseteq U_{j_0} \cup \bigcup_{k \in Z, k < j_0} U_k$. According to the choice of x , we have $x \in A$, but $x \notin \bigcup_{k \in Z, k < j_0} U_k$. This implies $x \in U_{j_0}$, contradicting the assumption. Thus, we have:

$$A \cap \left(\bigcap_{k \in Z} U_k^c \right) = A \cap \left(\bigcap_{j \in Y} U_j \right) \cap \left(\bigcap_{k \in Z} U_k^c \right).$$

As the set on the left is non-empty, so is the set on the right. With that we know that our Limit-machine always produces a valid output, that is the jump of some element. We have already established that any valid output is necessarily correct, and thereby the proof is complete. \square

As a corollary of this uniform result we get the known version of the Low Basis Theorem.

Corollary 8.4 (Low Basis Theorem of Jockusch and Soare) *Any non-empty co-c.e. closed set $A \subseteq \{0, 1\}^{\mathbb{N}}$ contains a low point.*

The property that computable instances always admit low solutions is preserved under Weihrauch reducibility, as pointed out in [6]. We will now show that this property also holds uniformly. This observation invites the question where $\mathfrak{L} = J^{-1} \circ \text{lim}$ is placed in the Weihrauch lattice. We will start the answer with an obvious corollary to Theorem 8.3, which will then be extended.

Corollary 8.5 $C_{\{0,1\}^{\mathbb{N}}} \leq_{\text{sW}} \mathfrak{L}$.

Now we will lift this observation from compact to locally compact choice. This involves again the same idea as the proof of Proposition 4.8, albeit in a new disguise as the following lemma:

Lemma 8.6 *Let β be a representation of Cantor space $\{0, 1\}^{\mathbb{N}}$, and let $\delta_{\mathbb{N}}$ be the standard representation of \mathbb{N} . If the multi-valued function $\mathbf{C}_{\{0,1\}^{\mathbb{N}}} : \subseteq \mathcal{A}_-(\{0, 1\}^{\mathbb{N}}) \rightrightarrows \{0, 1\}^{\mathbb{N}}$ is $(\psi_-^{\{0,1\}^{\mathbb{N}}}, \beta)$ -computable, then the multi-valued function $\mathbf{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}} : \subseteq \mathcal{A}_-(\mathbb{N} \times \{0, 1\}^{\mathbb{N}}) \rightrightarrows \mathbb{N} \times \{0, 1\}^{\mathbb{N}}$ is $(\psi_-^{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}, (\delta_{\mathbb{N}} \times \beta)^{\Delta})$ -computable.*

Proof. We describe a machine solving the latter task. Given a $\psi_-^{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}$ -name of a closed set $A \subseteq \mathbb{N} \times \{0, 1\}^{\mathbb{N}}$, it produces a sequence $(p_m)_{m \in \mathbb{N}}$. Again we use A^m to denote the complement of the union of the first m basic open sets listed in the input.

As $(\{n\} \times \{0, 1\}^{\mathbb{N}}) \cap A^m = \emptyset?$ is decidable and we have $A^m \neq \emptyset$, we can compute $n_m = \min\{n \in \mathbb{N} \mid (\{n\} \times \{0, 1\}^{\mathbb{N}}) \cap A^m \neq \emptyset\}$. Using these values, the output sequence shall be of the form $p_m = \langle \delta_{\mathbb{N}}^{-1}(n_m), q_{n_m} \rangle$. Note that n_m will be eventually constant as m goes to infinity, hence the same is true for the p_m .

The values q_{n_m} are computed as follows. A machine computing $\mathbf{C}_{\{0,1\}^{\mathbb{N}}} : \subseteq (\mathcal{A}_-(\{0, 1\}^{\mathbb{N}}), \psi_-^{\{0,1\}^{\mathbb{N}}}) \rightrightarrows (\{0, 1\}^{\mathbb{N}}, \beta)$ is simulated on input denoting $\text{pr}_2((\{n_m\} \times \{0, 1\}^{\mathbb{N}}) \cap A)$ for k steps, as long as $(\{n_m\} \times \{0, 1\}^{\mathbb{N}}) \cap A^k \neq \emptyset$ for $k \in \mathbb{N}$. If a k is reached with $(\{n_m\} \times \{0, 1\}^{\mathbb{N}}) \cap A^k = \emptyset$, the sequence q_{n_m} will be continued by 0s.

If n_m has reached its final value for m_0 , then $q_{n_{m_0}}$ will be a β -name for some w with $n_{m_0} \times w \in A$; this is sufficient to ensure that the overall output of the described computation is a $(\delta_{\mathbb{N}} \times \beta)^{\Delta}$ -name of $n_{m_0} \times w \in A$. \square

As a consequence we obtain that \mathfrak{L} is strictly above locally compact choice.

Theorem 8.7 $\mathbf{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}} \leq_{sW} \mathfrak{L}$ and $\mathfrak{L} \not\leq_W \mathbf{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}$.

Proof. To show the reduction, we make use of Theorem 8.3 together with Lemma 8.6 and the observation that $\beta^{\vee\Delta} \equiv \beta^{\vee}$ for any representation β . To see $\mathfrak{L} \not\leq_W \mathbf{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}$, observe that \mathfrak{L} is single-valued. Therefore, the assumption of the contrary together with Corollary 5.3 would imply $\mathfrak{L} \leq_W \mathbf{C}_{\mathbb{N}}$. By transitivity and Corollary 8.5 we get $\mathbf{C}_{\{0,1\}^{\mathbb{N}}} \leq_W \mathbf{C}_{\mathbb{N}}$. As shown in [6], the latter is wrong, providing the sought contradiction. \square

As J^{-1} is computable, the upper bound $\mathfrak{L} = J^{-1} \circ \lim \leq_W \lim$ is obtained directly. As \lim maps some computable inputs to non-low outputs, we even have $\mathfrak{L} <_W \lim$. With this, we have determined precisely the place of $J^{-1} \circ \lim$ in the diagram provided in Figure 1.

A question regarding the Weihrauch degree of \mathfrak{L} that is left open by the results presented so far is its behavior under products. Remarkable consequences of the following answers are that the low real numbers do not form a field, and that the integral does not commute with products.

Theorem 8.8 $\mathfrak{L} <_W \mathfrak{L} \times \mathfrak{L}$.

Proof. By a result of Spector (see [30] or Proposition V.2.26 in [24]) there are sequences $a, b \in \{0, 1\}^{\mathbb{N}}$, so that both a and b are low, but $\langle a, b \rangle$ is not low. Since a and b are low, $J(a)$ and $J(b)$ are Turing reducible to the halting problem, there are computable sequences $\langle a_0, a_1, \dots \rangle$ and $\langle b_0, b_1, \dots \rangle$ with $\lim_{i \rightarrow \infty} a_i = J(a)$ and $\lim_{i \rightarrow \infty} b_i = J(b)$. Then $(\langle a_0, a_1, \dots \rangle, \langle b_0, b_1, \dots \rangle)$ is computable, and we have $(J^{-1} \circ \lim \times J^{-1} \circ \lim)(\langle a_0, a_1, \dots \rangle, \langle b_0, b_1, \dots \rangle) = (a, b)$. Thus, $(J^{-1} \circ \lim) \times (J^{-1} \circ \lim)$ can map a computable input to an output that is not low. \square

In other words, this means that \mathfrak{L} is not idempotent. However, it has a different property. We call a function $T : \subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ a *jump operator*, if for all computable functions $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ there exists a computable function $G : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $F \circ T = T \circ G$. This notion has been introduced in [10] (for single-valued functions) and using this terminology the following has been proved in [4].

Lemma 8.9 *The limit \lim and the inverse of the Turing jump J^{-1} are jump operators and hence \mathfrak{L} is also a jump operator.*

Now we can formulate our main characterization of low computability.

Theorem 8.10 (Low computability) *Let f be a multi-valued function on represented spaces. Then the following are equivalent:*

- (1) $f \leq_{\text{sW}} \mathfrak{L}$,
- (2) f is low computable.

Proof. We consider the represented spaces (X, δ_X) and (Y, δ_Y) . If $f : \subseteq X \rightrightarrows Y$ is low computable, then there is a computable realizer F such that $\delta_Y^v \circ F(p) \in f\delta_X(p)$ for all $p \in \text{dom}(f\delta_X)$. Since $\delta_Y^v \circ F = \delta_Y \circ \mathfrak{L} \circ F$, this means that $\mathfrak{L} \circ F$ is a (δ_X, δ_Y) -realizer of f and hence $f \leq_{\text{sW}} \mathfrak{L}$. If, on the other hand, $f \leq_{\text{sW}} \mathfrak{L}$, then there are computable functions H, K such that $F = H\mathfrak{L}K$

is a (δ_X, δ_Y) -realizer of f . By Lemma 8.9 there is a computable function L such that $H\mathfrak{L} = \mathfrak{L}L$ and hence $F = \mathfrak{L}LK$ and LK is a $(\delta_X, \delta_Y^\vee)$ -realizer of f . \square

Next we want to show that certain choice principles are cylinders. We recall that a multi-valued map f on represented spaces is called a *cylinder*, if $\text{id} \times f \leq_{\text{sW}} f$. For cylinders f we have $g \leq_{\text{sW}} f \iff g \leq_{\text{W}} f$ (see [7]). It has already been proved in [7] that $\mathbf{C}_{\{0,1\}^{\mathbb{N}}}$ is a cylinder, here we present another proof that can be directly transferred to $\mathbf{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}$.

Proposition 8.11 $\mathbf{C}_{\{0,1\}^{\mathbb{N}}}$ and $\mathbf{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}$ are cylinders.

Proof. There is a computable embedding

$$\iota : \mathbb{N}^{\mathbb{N}} \rightarrow \{0, 1\}^{\mathbb{N}}, p \mapsto 01^{p(0)+1}01^{p(1)+1} \dots$$

and using this embedding we get $\text{id}_{\mathbb{N}^{\mathbb{N}}}(p) = \iota^{-1} \circ \mathbf{C}_{\{0,1\}^{\mathbb{N}}} \circ \text{in}_{\{0,1\}^{\mathbb{N}}} \circ \iota(p)$ and hence $\text{id}_{\mathbb{N}^{\mathbb{N}}} \leq_{\text{sW}} \mathbf{C}_{\{0,1\}^{\mathbb{N}}}$. The proofs of Propositions 3.4 and 3.7 even show strong Weihrauch reducibility. Hence, using a computable surjective pairing function $\pi : \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 1\}^{\mathbb{N}} \times \{0, 1\}^{\mathbb{N}}$ one obtains

$$\text{id}_{\mathbb{N}^{\mathbb{N}}} \times \mathbf{C}_{\{0,1\}^{\mathbb{N}}} \leq_{\text{sW}} \mathbf{C}_{\{0,1\}^{\mathbb{N}}} \times \mathbf{C}_{\{0,1\}^{\mathbb{N}}} \leq_{\text{sW}} \mathbf{C}_{\{0,1\}^{\mathbb{N}} \times \{0,1\}^{\mathbb{N}}} \leq_{\text{sW}} \mathbf{C}_{\{0,1\}^{\mathbb{N}}}.$$

Hence $\mathbf{C}_{\{0,1\}^{\mathbb{N}}}$ is a cylinder. The fact that $\mathbf{C}_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}$ is a cylinder can be proved analogously. \square

Together with Propositions 4.8 and 8.11, Corollary 4.9 and Theorems 8.7 and 8.10 we obtain the following corollary.

Corollary 8.12 *If X is a computable K_σ -space, then \mathbf{C}_X is low computable.*

This applies, in particular, to $\mathbf{C}_{\mathbb{N}}$, $\mathbf{C}_{\{0,1\}^{\mathbb{N}}}$ and $\mathbf{C}_{\mathbb{R}}$. We also obtain the following generalization of the non-uniform Low Basis Theorem of Jockusch and Soare. The case $\mathbf{C}_{\mathbb{R}}$ was already treated as Theorem 4.7 in [6].

Corollary 8.13 (Low Basis Theorem) *If X is a computable K_σ -space, then any non-empty co-c.e. closed set $A \subseteq X$ contains a low point.*

Together with Corollary 7.13 and Theorem 7.11 we obtain that the class of low computable functions contains several others.

Corollary 8.14 *Any multi-valued function f on represented spaces that is computable with finitely many mind changes or weakly computable is also low computable.*

We mention that one gets consequences as the following.

Corollary 8.15 *The Brouwer Fixed Point Theorem BFT is low computable.*

Here, $\text{BFT} : \mathcal{C}([0, 1]^n, [0, 1]^n) \rightrightarrows [0, 1]^n$ is the multi-valued map with $\text{BFT}(f) := \{x \in [0, 1]^n : f(x) = x\}$. In [6] it was already proved that any computable function $f : [0, 1]^n \rightarrow [0, 1]^n$ has a low fixed point and that the Brouwer Fixed Point Theorem is weakly computable. The above property is a uniform version of the former fact. The benefit of having uniform results is highlighted by the following result.

An interesting property of the class of low computable functions is that if they are composed with limit computable functions from the left, then one obtains a limit computable function again. This is in contrast to the fact that the limit computable functions themselves are not closed under composition.

Proposition 8.16 (Composition) *Let $f : \subseteq X \rightrightarrows Y$ and $g : \subseteq Y \rightrightarrows Z$ be multi-valued functions on represented spaces. If f is low computable and g is limit computable, then $g \circ f$ is limit computable. If f and g are both low computable, then $g \circ f$ is low computable.*

Proof. We use the represented spaces (X, δ_X) , (Y, δ_Y) and (Z, δ_Z) . We exploit the fact that integral and derivative of representations form a Galois connection (see [4]). That g is limit computable means that it is (δ_Y, δ'_Z) -computable, which is equivalent to g being $(f \delta_Y, \delta_Z)$ -computable and that f is low computable means that it is (δ_X, δ'_Y) -computable, which is equivalent to f being $(f \delta_X, f \delta_Y)$ -computable. It follows that $g \circ f$ is $(f \delta_X, \delta_Z)$ -computable, which is equivalent to $g \circ f$ being limit computable. Analogously, if f and g are both low computable, then it follows that $g \circ f$ is $(f \delta_X, f \delta_Z)$ -computable, which is equivalent to $g \circ f$ being low computable. \square

It can easily be seen that the composition $g \circ f$ of a limit computable f even with a g that is computable with finitely many mind changes is not necessarily limit computable. The class of low computable functions is the largest known class with the stability property expressed in Proposition 8.16.

We note that \mathcal{L} cannot be closed under composition by Theorem 8.8 and Proposition 7.4. Hence strict Weihrauch reducibility cannot be replaced by ordinary Weihrauch reducibility in Theorem 8.10.

9 The Jump Topology

Connecting to the results of Section 7, it seems reasonable to inquire whether other interesting Weihrauch degrees can be characterized by restrictions of the limit operation \lim of Baire space $\mathbb{N}^{\mathbb{N}}$. Since all such restrictions are single-valued, neither $\mathbf{C}_{\{0,1\}^{\mathbb{N}}}$ nor $\mathbf{C}_{\mathbb{R}}$ can be equivalent to such an operation, as their Weihrauch degrees do not contain any single-valued functions by Corollaries 5.2 and 5.3. In the remainder of this section, we will study the limit operator \lim_J with respect to the initial topology of the jump J . Like \lim_{Δ} we will consider this operation as an operation with respect to Baire space (with the identity as standard representation).

Initially, we suspected that \lim_J might be equivalent to $\mathfrak{L} = J^{-1} \circ \lim$, but this is only true topologically, as we will show in Theorem 9.10. Computationally, the contrary result is given below (see Theorem 9.6). It turned out that the initial topology of the jump is identical to the Π -topology studied by Miller [21, Chapter IV].

Theorem 9.1 *The initial topology of J is generated by the co-c.e. closed sets (that is identical to the Π -topology).*

Proof. As every basic set of the form $w\mathbb{N}^{\mathbb{N}}$ for some finite w is co-c.e. closed, every set that is open in the ordinary Baire topology is also open in the Π -topology. Now consider the preimage:

$$J^{-1}(w\mathbb{N}^{\mathbb{N}}) = \left(\bigcap_{i < |w|, w(i)=1} U_i \right) \cap \left(\bigcap_{j < |w|, w(j)=0} U_j^c \right).$$

In the Π -topology, this is an intersection of finitely many open sets, and therefore open. As the Baire topology is generated by sets of the form $w\mathbb{N}^{\mathbb{N}}$, this shows that the jump J is continuous with the Π -topology on its domain and the Baire topology on its codomain. This is equivalent to the inclusion of the initial topology of J in the Π -topology.

For the other inclusion, fix some co-c.e. closed set U_n^c . We have

$$U_n^c = \bigcup_{w \in \mathbb{N}^n} J^{-1}(w0\mathbb{N}^{\mathbb{N}}),$$

so U_n^c is open in the initial topology of the jump. This concludes the proof. \square

A sequence $(p_n)_{n \in \mathbb{N}}$ in $\mathbb{N}^{\mathbb{N}}$ converges to $p \in \mathbb{N}^{\mathbb{N}}$ regarding the Π -topology, if $(J(p_n))_{n \in \mathbb{N}}$ converges to $J(p)$ in Baire space. The limit value p cannot be left out here: There is a sequence $(p_n)_{n \in \mathbb{N}}$, so that $(J(p_n))_{n \in \mathbb{N}}$ converges in Baire

space, but not to some element of the range of J , as the range of J is not closed in Baire space. The above description of the convergence relation of the Π -topology implies

$$\lim_J = J^{-1} \circ \lim \circ J^{\mathbb{N}} = \mathfrak{L} \circ J^{\mathbb{N}},$$

with $J^{\mathbb{N}}\langle p_0, p_1, p_2, \dots \rangle := \langle J(p_0), J(p_1), J(p_2), \dots \rangle$.

In order to understand the computability aspects of \lim_J , we would like to know which points are limits of computable sequences with respect to the Π -topology. We introduce a name for these points.

Definition 9.2 A point $p \in \mathbb{N}^{\mathbb{N}}$ is called *limit computable in the jump*, if there is a computable sequence $(p_n)_{n \in \mathbb{N}}$ in $\mathbb{N}^{\mathbb{N}}$ such that $\lim_{n \rightarrow \infty} J(p_n) = J(p)$.

Here the limit is understood with respect to the ordinary Baire topology and by continuity of J^{-1} we automatically obtain $\lim_{n \rightarrow \infty} p_n = p$. Some necessary properties of points p that are limit computable in the jump are clear. For one, they are limit computable and secondly they are in the closure of the set of computable points with respect to the Π -topology. These points are called *unavoidable* following Kalantari and Welch (see [17] and [21]).

Another observation is that all limit computable 1-generics are limit computable in the jump. We recall that a point $p \in \mathbb{N}^{\mathbb{N}}$ is called *1-generic*, if for all $n \in \mathbb{N}$ there exists a finite word $w \sqsubseteq p$ such that either $w\mathbb{N}^{\mathbb{N}} \subseteq U_n$ or $w\mathbb{N}^{\mathbb{N}} \cap U_n = \emptyset$ (see [23]). Here $(U_n)_{n \in \mathbb{N}}$ denotes the computable standard enumeration of all c.e. open subsets of Baire space that was used to define the Turing jump J . The definition directly implies the following observation.

Lemma 9.3 *The Turing jump operator $J : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ is continuous in $p \in \mathbb{N}^{\mathbb{N}}$ if and only if p is 1-generic.*

Using this lemma, we obtain the following sufficient condition for limit computability in the jump.

Proposition 9.4 *If p is 1-generic and limit computable, then p is limit computable in the jump.*

Proof. If p is limit computable, then there is a computable sequence $(p_n)_{n \in \mathbb{N}}$ that converges to p . If p is 1-generic, then $(J(p_n))_{n \in \mathbb{N}}$ also converges to $J(p)$ according to Lemma 9.3. This means that p is limit computable in the jump. \square

It is known that there is a 1-generic and limit computable $p \in \mathbb{N}^{\mathbb{N}}$ (see Theorem 1.8.52 in [23]). Moreover, a 1-generic cannot be computable (see for instance Proposition XI.2.3 in [25]). Hence, it follows that \lim_J maps some computable input to a non-computable output and hence it is not non-uniformly computable.

It will follow from Proposition 9.11 below that points which are non-computable and limit computable in the jump are not necessarily 1-generic. However, they seem to share a lot of properties with the class of limit computable 1-generics. As one such property we prove that points which are limit computable in the jump do not bound diagonally non-computable functions. A total function $f : \mathbb{N} \rightarrow \mathbb{N}$ is called *diagonally non-computable* if $f(i) \neq \varphi_i(i)$ for all $i \in \mathbb{N}$ (that means either $\varphi_i(i)$ does not exist or otherwise the two values are not equal). Here φ denotes some standard Gödel numbering of the partial computable functions $g : \subseteq \mathbb{N} \rightarrow \mathbb{N}$. Diagonally non-computable functions are, in particular, not computable. As we will show below, our following proposition is related to the known result that 1-generics do not bound diagonally non-computable functions (due to Demuth and Kučera, see Corollary 9 in [13]). The proof is inspired by Nies (see Exercise 4.1.6 in [23]).

Proposition 9.5 *Let f be diagonally non-computable and let p be limit computable in the jump. Then $f \not\leq_T p$.*

Proof. Let f be diagonally non-computable and let p be limit computable in the jump. Let us assume that $f \leq_T p$. Then there is a computable function $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $F(p) = f$ and there is a computable sequence $(p_n)_{n \in \mathbb{N}}$ which converges to p in the Π -topology. Since F is computable, there is a Turing machine M that computes F . Let us denote by $F_M(r)(n)$ the n -th symbol written by this machine M upon input r , irrespectively of whether $r \in \text{dom}(F)$. Then the set

$$U := \{r \in \mathbb{N}^{\mathbb{N}} : (\exists i \in \mathbb{N}) (F_M(r)(i) = \varphi_i(i) \text{ and } i \in \text{dom}(\varphi_i))\}$$

is c.e. open and since $f = F(p)$ is diagonally non-computable, it follows that $p \notin U$. Since $(p_n)_{n \in \mathbb{N}}$ converges to p in the Π -topology and the complement of U is open in the Π -topology by Theorem 9.1, it follows that $p_n \notin U$ for all $n \geq m$ with some fixed $m \in \mathbb{N}$. Since $f = F(p)$ is total and $(p_n)_{n \in \mathbb{N}}$ converges to p , there must be an $n \geq m$ for each $i \in \mathbb{N}$ such that $F_M(p_n)(i)$ exists. Since $(p_n)_{n \in \mathbb{N}}$ is computable, we can even find such an n effectively, i.e. there is a computable function $s : \mathbb{N} \rightarrow \mathbb{N}$ such that $F_M(p_{s(i)})(i)$ exists and $s(i) \geq m$ for all $i \in \mathbb{N}$. Since $p_{s(i)} \notin U$, we obtain $F_M(p_{s(i)})(i) \neq \varphi_i(i)$. But that means that $g(i) := F_M(p_{s(i)})(i)$ defines a total computable function $g : \mathbb{N} \rightarrow \mathbb{N}$ that is diagonally non-computable, which is a contradiction! \square

From this result we can directly conclude that choice on Cantor space $C_{\{0,1\}^{\mathbb{N}}}$ is not reducible to \lim_J . A function f is called *two-valued diagonally non-computable* if it is diagonally non-computable and $\text{range}(f) \subseteq \{0, 1\}$. It is known that the set of all such functions is co-c.e. closed in Cantor space $\{0, 1\}^{\mathbb{N}}$ (see Fact 1.8.31 in [23]).

Theorem 9.6 *We obtain $C_{\{0,1\}^{\mathbb{N}}} \not\leq_W \lim_J$.*

Proof. Let us assume to the contrary that $C_{\{0,1\}^{\mathbb{N}}} \leq_W \lim_J$. Then there are computable functions H, K such that $H\langle p, \lim_J K(p) \rangle \in C_{\{0,1\}^{\mathbb{N}}}\psi_-(p)$ for all p in the domain of the right-hand side. It is known and easy to see that the set

$$A := \{f \in \{0, 1\}^{\mathbb{N}} : f \text{ is two-valued diagonally non-computable}\}$$

is a co-c.e. closed set. Hence, there is a computable p such that $A = \psi_-(p)$ and we obtain that $f := H\langle p, \lim_J K(p) \rangle$ is diagonally non-computable. Hence $K(p)$ is computable and $q := \lim_J K(p)$ is limit computable in the jump. Moreover, $f \leq_T q$, which contradicts Proposition 9.5. \square

Next we prove that \lim_J is low computable.

Theorem 9.7 *We obtain $\lim_J <_{sW} \mathfrak{L}$.*

Proof. We use the computable standard enumeration $(U_n)_{n \in \mathbb{N}}$ of c.e. open subsets $U_n \subseteq \mathbb{N}^{\mathbb{N}}$ that was used to define the Turing jump operator J . By U_n^m we denote the union of the first m basic clopen balls in the union that constitutes U_n . We define a function $F : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ by $F\langle p_0, p_1, p_2, \dots \rangle := \langle q_0, q_1, q_2, \dots \rangle$ with

$$q_{\langle k, m \rangle}(n) := \begin{cases} 1 & \text{if } p_k \in U_n^m \\ 0 & \text{otherwise} \end{cases}$$

Since the property $p_k \in U_n^m$ is decidable in the input sequence and the parameters k, n, m , it follows that F is computable. We claim that $\lim_J = \mathfrak{L} \circ F$. Let $(p_k)_{k \in \mathbb{N}}$ and p be such that $\lim_{k \rightarrow \infty} J(p_k) = J(p)$. Then also $\lim_{k \rightarrow \infty} p_k = p$. Let $(q_i)_{i \in \mathbb{N}}$ be the corresponding output of F . Let us assume that $J(p)(n) = 1$ for some $n \in \mathbb{N}$, i.e. $p \in U_n$. Then $p \in U_n^m$ for all sufficiently large m and hence $p_k \in U_n^m$ for all sufficiently large k, m . This implies that $q_{\langle m, k \rangle}(n) = 1$ for sufficiently large $\langle m, k \rangle$. Let us now assume that $J(p)(n) = 0$, i.e. $p \notin U_n$. Since the complement of U_n is co-c.e. closed and hence open in the Π -topology, this implies that $p_k \notin U_n$ for all sufficiently

large k . In particular, $p_k \notin U_n^m$ for all m and all sufficiently large k . This implies that $q_{\langle m, k \rangle}(n) = 0$ for all sufficiently large $\langle m, k \rangle$. Altogether, this means $\mathfrak{L} \circ F \langle p_0, p_1, p_2, \dots \rangle = J^{-1} \circ \lim \langle q_0, q_1, q_2, \dots \rangle = p$, as desired. By Theorem 9.6 the reduction is strict. \square

As a corollary we obtain the following.

Corollary 9.8 *All $p \in \mathbb{N}^{\mathbb{N}}$ which are limit computable in the jump are also low.*

This is another property that points which are limit computable in the jump share with limit computable 1-generics (see Proposition XI.2.3.2 in [25]). Another straightforward observation is the following.

Corollary 9.9 *We obtain $\lim_{\Delta} <_W \lim_J <_W \lim$.*

Since the corresponding topologies are included in each other in the converse order, each limit operation in this sequence is just a restriction of the next one. This implies the positive part of the reduction chain. The first reduction is strict, since \lim_{Δ} is non-uniformly computable and \lim_J is not (as observed after Proposition 9.4). The second reduction is strict since $C_{\{0,1\}^{\mathbb{N}}}$ is reducible to \lim , but not to \lim_J (by Theorem 9.6).

In light of Theorem 9.7 it might be surprising that topologically \lim_J turns out to be equivalent to \mathfrak{L} .

Theorem 9.10 *We obtain $\lim_J \equiv_{sW} \mathfrak{L}$ with respect to some oracle.*

Proof. By Theorem 9.7 it is clear that $\lim_J \leq_{sW} \mathfrak{L}$. We need to show the reverse reduction with respect to some oracle.

Let $(U_i)_{i \in \mathbb{N}}$ be the standard enumeration of c.e. open sets used to define the jump operator J . Given a finite word $w = w_0 \dots w_n \in \mathbb{N}^*$ we use the sets

$$A_{w,i} := \begin{cases} U_i & \text{if } w_i \neq 0 \\ \mathbb{N}^{\mathbb{N}} \setminus U_i & \text{otherwise} \end{cases}$$

for all $i = 0, \dots, n$. Moreover, we set $A_w := \bigcap_{i=0}^n A_{w,i}$. Now we define inductively a function $f : \mathbb{N}^* \rightarrow \mathbb{N}^{\mathbb{N}}$ by $f(\varepsilon) := \hat{0}$ for the empty word ε and for $w := w_0 \dots w_{n+1}$ we select $f(w) \in A_w$ if $A_w \neq \emptyset$ and $f(w) := f(w_0 \dots w_n)$ otherwise. By the Axiom of Choice such a function f exists and we use it as an oracle in the following. Given a sequence $p = \langle p_0, p_1, p_2, \dots \rangle \in \text{dom}(\mathfrak{L})$ we let

$$F(p) := \langle q_0, q_1, q_2, \dots \rangle \text{ with } q_i := f(p_i[i]),$$

where $p_i[j] = p_i(0) \dots p_i(j)$ denotes the prefix of p_i of length $j + 1$. It is clear that F is computable in the oracle f .

We claim that $\lim_J F(p) = \mathfrak{L}(p)$ for all $p \in \text{dom}(\mathfrak{L})$. Given a sequence $p = \langle p_0, p_1, p_2, \dots \rangle \in \text{dom}(\mathfrak{L})$ it follows that the sequence $(p_i)_{i \in \mathbb{N}}$ converges to $J\mathfrak{L}(p)$ in the usual Baire topology. We consider $(q_i)_{i \in \mathbb{N}}$ with $q_i = f(p_i[i])$ as above. Let $n \in \mathbb{N}$. Then there is an $i \geq n$ such that

$$p_j(m) = 1 \iff \mathfrak{L}(p) \in U_m$$

for all $j \geq i$ and $m \leq n$. In this situation $A_{p_i[n]} \neq \emptyset$ since $\mathfrak{L}(p) \in A_{p_i[n]}$ and hence $q_j \in A_{p_i[n]}$ for all $j \geq i$ by definition of f . In particular,

$$q_j \in U_n \iff \mathfrak{L}(p) \in U_n$$

for all $j \geq i$. This means that $(q_j)_{j \in \mathbb{N}}$ converges to $\mathfrak{L}(p)$ in the Π -topology and hence $\lim_J F(p) = \mathfrak{L}(p)$. \square

As a last result on limit computability in the limit we prove that this class of points is closed under total computable functions.

Proposition 9.11 *Let $p, q \in \mathbb{N}^{\mathbb{N}}$ be such that $F(p) = q$ for some total computable function $F : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$. If p is limit computable in the jump, then q is limit computable in the jump too.*

Proof. Let $F : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ be some total computable function such that $F(p) = q$. Hence JF is limit computable and hence there exists a computable $G : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $JF = GJ$ by Lemma 8.9. If p is limit computable in the jump, then there is a computable sequence $(p_n)_{n \in \mathbb{N}}$ such that $(J(p_n))_{n \in \mathbb{N}}$ converges to $J(p)$. Since G is continuous, we obtain that $(GJ(p_n))_{n \in \mathbb{N}}$ converges to $GJ(p)$, which implies that $(JF(p_n))_{n \in \mathbb{N}}$ converges to $JF(p)$. Since F is computable, it follows that $(F(p_n))_{n \in \mathbb{N}}$ is computable and this means that $q = F(p)$ is limit computable in the jump. \square

From this result it follows that p which are non-computable and limit computable in the jump are not necessarily 1-generic. For instance, for each limit computable 1-generic p we have that $\langle \widehat{0}, p \rangle$ is limit computable in the jump and non-computable, but it is not 1-generic, since no finite prefix proves that it does belong to the co-c.e. closed set $\{\langle \widehat{0}, q \rangle : q \in \mathbb{N}^{\mathbb{N}}\}$. So far, we have no example of a point that is limit-computable in the jump and not below a 1-generic with respect to truth-table reducibility. It would be useful to clarify the relation between 1-generics and points that are limit computable in the jump somewhat further.

The Π -topology shows further interesting behavior. If p is computable in Baire space, then it is isolated regarding the Π -topology, that is the singletons $\{p\}$ with computable p are clopen. We characterize the singletons $\{p\}$ that are clopen in the Π -topology.

Lemma 9.12 *Let $p \in \mathbb{N}^{\mathbb{N}}$. Then $\{p\}$ is clopen in the Π -topology if and only if $\{p\}$ is co-c.e. closed in Baire space.*

Proof. Since the Π -topology includes the ordinary Baire topology, it is clear that all singletons $\{p\}$ are closed in the Π -topology. If $\{p\}$ is co-c.e. closed in Baire space, then $\{p\}$ is also open in the Π -topology (since this topology is generated by the co-c.e. closed sets). Let now $\{p\}$ be open in the Π -topology. Then there is a finite prefix $w \sqsubseteq J(p)$ such that $\{p\} = J^{-1}(w\mathbb{N}^{\mathbb{N}})$. Similarly to the proof of Theorem 9.1 we obtain

$$\{p\} = J^{-1}(w\mathbb{N}^{\mathbb{N}}) = \left(\bigcap_{i < |w|, w(i)=1} U_i \right) \cap \left(\bigcap_{j < |w|, w(j)=0} U_j^c \right).$$

However, in this case the open sets U_i with $w(i) = 1$ can be replaced by clopen balls, since $\{p\}$ is a singleton. Altogether, this implies that $\{p\}$ can be written as a finite intersection of co-c.e. closed sets and hence it is co-c.e. closed. \square

It is easy to see that there are co-c.e. closed singletons $\{p\}$ with non-computable $p \in \mathbb{N}^{\mathbb{N}}$. Co-c.e. closed singletons $\{p\}$ can even be such that p is not arithmetical (see Propositions 1.8.62 and 1.8.70 in [23]). Lemma 9.12 implies that the set of computable points is open in the Π -topology, although not effectively so. In general, a set O is c.e. open in the Π -topology, if and only if it is effectively F_σ in Baire space, in turn, a set A is co-c.e. closed in the Π -topology, if and only if it is effectively G_δ in Baire space. In particular, the Martin-Löf random points form a c.e. open set (and a proper subset of the open set of avoidable points). While the Π -topology *makes everything easier* when considering sets, it *makes everything more complicated* when considering points: a point is Turing reducible to the n -th jump of the empty set in Baire space if and only if it has a name in the Π -space that is Turing reducible to the $n + 1$ -st jump of the empty set. This contrary behavior of points and sets is based on the fact that points are mapped forwards and sets are mapped backwards.

We close by mentioning another property of the Turing jump J . The Galois connection between the Turing jump J and the limit \lim cannot be extended to the continuous category. That is, we get the following counterexample, which shows that the inverse J^{-1} is not a “topological jump operator” (see Lemma 8.9).

Proposition 9.13 *There exists a total continuous function $F : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$*

such that there is no continuous function $G : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ with $FJ^{-1} = J^{-1}G$.

Proof. Let $c, d \in \mathbb{N}^{\mathbb{N}}$ be such that c is computable and $\{d\}$ is not co-c.e. closed. Then according to Lemma 9.12 $\{c\}$ is clopen and $\{d\}$ is not clopen in the Π -topology. Since by Theorem 9.1 the Π -topology is just the initial topology of the jump J , which is injective, it follows that $\{J(c)\}$ is clopen and $\{J(d)\}$ is not clopen in $\text{range}(J)$ with respect to the ordinary Baire topology. Now we define a continuous map $F : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ by

$$F(p) := c + |d - p|$$

for all $p \in \mathbb{N}^{\mathbb{N}}$, where all arithmetic operations are meant pointwise. It is clear that F is continuous with $F^{-1}\{c\} = \{d\}$. Let us now assume that some map $G : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ has the property $FJ^{-1} = J^{-1}G$. In particular $\text{dom}(G) = \text{range}(J)$ in this situation. Then we obtain

$$G^{-1}\{J(c)\} = (J^{-1}G)^{-1}\{c\} = (FJ^{-1})^{-1}\{c\} = \{J(d)\}.$$

That is, although the set $\{J(c)\}$ is clopen in $\text{range}(J) = \text{dom}(J^{-1})$, its preimage under G is not clopen in $\text{dom}(G) = \text{range}(J)$ and hence G cannot be continuous. \square

10 Conclusions

We summarize some of the results that we have obtained in tables and figures. Figure 1 extends the results provided in [6, Figure 6]. Here $\mathbf{0}$ denotes the Weihrauch degree of the nowhere defined functions and one obtains as $\mathbf{0}^*$ the degree of all pointed computable multi-valued functions on represented spaces. The table below gives a list of some classes of multi-valued functions on represented spaces that can be characterized by choice for certain spaces. The given topological counterparts are at least correct for computable Polish spaces and in some cases they have only been proved for single-valued functions.

The notion “weakly computable with finitely many mind changes” has not been used before and is an ad hoc creation just for the purposes of this table.




Fig. 1. Closed choice in the Weihrauch lattice

Choice	Class of functions (topologically)
$C_{\{0\}}$	computable (continuous)
$C_{\mathbb{N}}$	computable with finitely many mind changes (piecewise continuous)
$C_{\{0,1\}^{\mathbb{N}}}$	weakly computable (upper semi-continuous compact-valued selectors)
$C_{\mathbb{N} \times \{0,1\}^{\mathbb{N}}}$	weakly computable with finitely many mind changes
$\widehat{C}_{\mathbb{N}}$	limit computable (Σ_2^0 -measurable)
$C_{\mathbb{N}^{\mathbb{N}}}$	effectively Borel measurable (Borel measurable)

Acknowledgement

We thank the anonymous referees for several valuable corrections, comments and remarks that helped to improve the final version of this paper.

References

- [1] V. Bosserhoff, Computable functional analysis and probabilistic computability, Ph.D. thesis, University of the Armed Forces, Munich (2008).
- [2] V. Brattka, Recursive and computable operations over topological structures, Informatik Berichte 255, University of Hagen, Hagen, PhD Thesis (Jul. 1999).
- [3] V. Brattka, Effective Borel measurability and reducibility of functions, *Mathematical Logic Quarterly* 51 (1) (2005) 19–44.
URL <http://dx.doi.org/10.1002/ma1q.200310125>
- [4] V. Brattka, Limit computable functions and subsets, unpublished notes (2007).
- [5] V. Brattka, Plottable real number functions and the computable graph theorem, *SIAM Journal on Computing* 38 (1) (2008) 303–328.
URL <http://dx.doi.org/10.1137/060658023>
- [6] V. Brattka, G. Gherardi, Effective choice and boundedness principles in computable analysis, *The Bulletin of Symbolic Logic* (to appear), preliminary version: <http://arxiv.org/abs/0905.4685>.
- [7] V. Brattka, G. Gherardi, Weihrauch degrees, omniscience principles and weak computability, *The Journal of Symbolic Logic* (to appear), preliminary version: <http://arxiv.org/abs/0905.4679>.
- [8] V. Brattka, G. Gherardi, Borel complexity of topological operations on computable metric spaces, *Journal of Logic and Computation* 19 (1) (2009) 45–76.
URL <http://dx.doi.org/10.1093/logcom/exn027>

- [9] V. Brattka, G. Presser, Computability on subsets of metric spaces, *Theoretical Computer Science* 305 (2003) 43–76.
URL [http://dx.doi.org/10.1016/S0304-3975\(02\)00693-X](http://dx.doi.org/10.1016/S0304-3975(02)00693-X)
- [10] M. de Brecht, Jump operators, unpublished notes (2009).
- [11] M. de Brecht, A. Yamamoto, Mind change complexity of inferring unbounded unions of pattern languages from positive data, *Theoretical Computer Science* 411 (2010) 976–985.
- [12] M. de Brecht, A. Yamamoto, Topological properties of concept space, *Information and Computation* 208 (4) (2010) 327–340.
- [13] O. Demuth, A. Kučera, Remarks on 1-genericity, semigenericity and related concepts, *Commentationes Mathematicae Universitatis Carolinae* 28 (1) (1987) 85–94.
- [14] G. Gherardi, A. Marcone, How incomputable is the separable Hahn-Banach theorem?, *Notre Dame Journal of Formal Logic* 50 (2009) 393–425.
- [15] P. Hertling, Unstetigkeitsgrade von Funktionen in der effektiven Analysis, *Informatik Berichte* 208, FernUniversität Hagen, Hagen, dissertation (Nov. 1996).
- [16] C. G. Jockusch, Jr., R. I. Soare, Degrees of members of Π_1^0 classes, *Pacific J. Math.* 40 (1972) 605–616.
- [17] I. Kalantari, L. Welch, A blend of methods of recursion theory and topology, *Annals of Pure and Applied Logic* 124 (1–3) (2003) 141–178.
- [18] A. S. Kechris, *Classical Descriptive Set Theory*, vol. 156 of Graduate Texts in Mathematics, Springer, Berlin, 1995.
- [19] S. C. Kleene, *Introduction to Metamathematics*, vol. 1 of *Bibliotheca Mathematica*, North-Holland, Amsterdam, 1952.
- [20] O. V. Kudinov, V. L. Selivanov, A. V. Zhukov, Undecidability in Weihrauch degrees, in: F. Ferreira, B. Löwe, E. Mayordomo, L. Mendes Gomes (eds.), *Programs, Proofs, Processes*, vol. 6158 of *Lecture Notes in Computer Science*, Springer, Berlin, 2010, 6th Conference on Computability in Europe, CiE 2010, Ponta Delgada, Azores, Portugal, June/July 2010.
URL <http://dx.doi.org/10.1007/978-3-642-13962-8>
- [21] J. S. Miller, Π_1^0 -1 classes in computable analysis and topology, Ph.D. thesis, Cornell University, Ithaca, USA (2002).
- [22] Y. N. Moschovakis, *Descriptive Set Theory*, vol. 100 of *Studies in Logic and the Foundations of Mathematics*, North-Holland, Amsterdam, 1980.
- [23] A. Nies, *Computability and Randomness*, vol. 51 of *Oxford Logic Guides*, Oxford University Press, New York, 2009.
- [24] P. Odifreddi, *Classical Recursion Theory*, vol. 125 of *Studies in Logic and the Foundations of Mathematics*, North-Holland, Amsterdam, 1989.

- [25] P. Odifreddi, Classical Recursion Theory - Volume II, vol. 143 of Studies in Logic and the Foundations of Mathematics, North-Holland, Amsterdam, 1999.
- [26] A. Pauly, How discontinuous is computing Nash equilibria? preliminary version: <http://arxiv.org/abs/0907.1482>.
- [27] A. Pauly, Infinite oracle queries in type-2 machines (extended abstract) preliminary version: <http://arxiv.org/abs/0907.3230>.
- [28] A. Pauly, On the (semi)lattices induced by continuous reducibilities, *Mathematical Logic Quarterly* 56 (5) (2010) 488–502.
- [29] V. L. Selivanov, Hierarchies of Δ_2^0 -measurable k -partitions, *Mathematical Logic Quarterly* 53 (4–5) (2007) 446–461.
URL <http://dx.doi.org/10.1002/malq.200710011>
- [30] C. Spector, On degrees of recursive unsolvability, *Annals of Mathematics* (2) 64 (1956) 581–592.
- [31] T. v. Stein, Vergleich nicht konstruktiv lösbarer Probleme in der Analysis, Diplomarbeit, Fachbereich Informatik, FernUniversität Hagen (1989).
- [32] W. Wadge, Degrees of complexity of subsets of the Baire space, *Notices of the Amer. Math. Soc. A* (1972) 714–715.
- [33] W. Wadge, Reducibility and determinateness on the Baire space, Thesis, University of California, Berkeley (1983).
- [34] K. Weihrauch, The degrees of discontinuity of some translators between representations of the real numbers, Technical Report TR-92-050, International Computer Science Institute, Berkeley (Jul. 1992).
- [35] K. Weihrauch, The TTE-interpretation of three hierarchies of omniscience principles, *Informatik Berichte* 130, FernUniversität Hagen, Hagen (Sep. 1992).
- [36] K. Weihrauch, *Computable Analysis*, Springer, Berlin, 2000.
- [37] K. Weihrauch, Computational complexity on computable metric spaces, *Mathematical Logic Quarterly* 49 (1) (2003) 3–21.
- [38] M. Ziegler, Real hypercomputation and continuity, *Theory of Computing Systems* 41 (1) (2007) 177–206.
URL <http://dx.doi.org/10.1007/s00224-006-1343-6>
- [39] M. Ziegler, Revising type-2 computation and degrees of discontinuity, in: D. Cenzer, R. Dillhage, T. Grubba, K. Weihrauch (eds.), *Proceedings of the Third International Conference on Computability and Complexity in Analysis*, vol. 167 of *Electronic Notes in Theoretical Computer Science*, Elsevier, Amsterdam, 2007, CCA 2006, Gainesville, Florida, USA, November 1–5, 2006.
URL <http://dx.doi.org/10.1016/j.entcs.2006.08.015>