



Swansea University  
Prifysgol Abertawe



## Cronfa - Swansea University Open Access Repository

---

This is an author produced version of a paper published in:  
*International Journal of Population Data Science*

Cronfa URL for this paper:  
<http://cronfa.swan.ac.uk/Record/cronfa44400>

---

### Paper:

North, L., Davies, L., Berridge, D., Walters, A., Akbari, A. & Lyons, R. (2018). Osteoporosis and fracture risk - a linked data study in Wales. *International Journal of Population Data Science*, 3(4)  
<http://dx.doi.org/10.23889/ijpds.v3i4.684>

This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.

---

This item is brought to you by Swansea University. Any person downloading material is agreeing to abide by the terms of the repository licence. Copies of full text items may be used or reproduced in any format or medium, without prior permission for personal research or study, educational or non-commercial purposes only. The copyright for any work remains with the original author unless otherwise specified. The full-text must not be sold in any format or medium without the formal permission of the copyright holder.

Permission for multiple reproductions should be obtained from the original author.

Authors are personally responsible for adhering to copyright and publisher restrictions when uploading content to the repository.

<http://www.swansea.ac.uk/library/researchsupport/ris-support/>

## Osteoporosis and fracture risk - a linked data study in Wales

North, L<sup>1</sup>, Davies, L<sup>2</sup>, Berridge, D<sup>1</sup>, Walters, A<sup>1</sup>, Akbari, A<sup>3</sup>, and Lyons, R<sup>1</sup>

<sup>1</sup>Farr Institute, Swansea University Medical School

<sup>2</sup>Public Health Wales

<sup>3</sup>Health Data Research UK - Wales and Northern Ireland, Swansea University Medical School

### Introduction

Osteoporosis is a global disease with a 30-40% lifetime fracture risk according to the World Health Organisation. Over half a million people receive treatment for fragility fractures annually in the UK. Osteoporosis incidence is rising with aging populations; however, medical secondary prevention treatment may reduce fracture risk.

### Conclusion/Implications

A population based e-cohort was successfully created by linking data across multiple datasets. Preliminary findings identified that <50% of eligible patients receive secondary medical prevention treatment after an index fragility fracture. These findings may help inform and unify treatment pathways for those at risk of fragility fractures.

### Objectives and Approach

Primary aims were to investigate if secondary medical prevention treatment following an index fracture was associated with survival and subsequent fracture risk, evaluated using a pseudonymised population based e-cohort study design. Patients aged  $\geq 60$  years with an index fragility fracture at any anatomical location were identified from the Secure Anonymised Information Linkage (SAIL) databank. Fracture data were identified from secondary care datasets (emergency department and inpatient) and the National Hip Fracture Register data. In addition linkages were made to primary care datasets for medical prescription and Office for National Statistics records for mortality, supplementing data on demographic characteristics and co-morbidity.

### Results

The cohort comprised 81,252 cases between April 2009 and December 2016 of median age 78 years (range 60-109) and 22,896 (28%) males. Medical secondary prevention treatment was received by 29,393 cases (36%). Subsequent fractures were reported for 10,907 cases (14%) and 29,026 cases (36%) died during the study period. For those that received medical prevention, the subsequent fracture and mortality rates were 15% and 28% respectively compared to 12% and 31% for those that did not receive the prevention treatment. Further analyses will include a discrete time competing risks model.

