




جامعة نايف العربية للعلوم الأمنية  
Naif Arab University for Security Sciences

Naif Arab University for Security Sciences  
Arab Journal of Forensic Sciences & Forensic Medicine

www.nauss.edu.sa  
http://ajfsfm.nauss.edu.sa



الجمعية العربية للعلوم الجنائية والطب الشرعي  
Arab Society for forensic Sciences and forensic Medicine

**Commentary on: Attafi IM, Albeishy MY, Oraiby ME, Khardali IA, Shaikhain GA, Fageeh MM. Postmortem Distribution of Cathinone and Cathine in Human Biological Specimens in a Case of Death Associated with Khat Chewing. Arab J Forensic Sci Forensic Med. 2018 Jun 7;1(7).**

مستويات الكاثين والكاثينون في الوفيات المرتبطة بالقات

John M. Corkery <sup>1,\*</sup>, Fabrizio Schifano <sup>1</sup>, Amira Guirguis <sup>1</sup>

<sup>1,2</sup> Psychopharmacology, Drug Misuse and Novel Psychoactive Substances Research Unit, Department of Clinical and Pharmaceutical Sciences, School of Life and Medical Sciences, University of Hertfordshire, College Lane Campus, Hatfield, Hertfordshire, AL10 9AB, United Kingdom.

Received 16 Apr. 2019; Accepted 02 May. 2019; Available Online xxxx

## Abstract

The interpretation of post-mortem human tissue toxicology levels may be affected by the sampling site chosen. It is important to bear this in mind when looking at the psychoactive constituents of khat (*Catha edulis* Forsk) that have been consumed and have contributed to or caused death. The post-mortem levels of cathine, cathinone and norephedrine/ norpseudoephedrine are very rarely reported, thereby making it impossible for toxicologists, pathologists and others investigating khat-related fatalities to decide if a level is toxic or fatal. This paper presents all the published data that exists to help start documenting this neglected area. Such information should be collected and reported on a systematic basis to facilitate correct interpretations in the future.

## المستخلص

من الممكن أن تتأثر مستويات السموم الدوائية في الأنسجة البشرية بعد الوفاة من خلال الموقع الجسدي المختار لأخذ العينة. من المهم أخذ ذلك في الاعتبار عند تحليل المكونات ذات التأثير النفسي في القات (*Catha edulis* Forsk). التي تم استهلاكها وساهمت في أو تسببت في الوفاة. نادرا ما يتم الإبلاغ عن مستويات الكاثين والكاثينون والنورفيدرين / نوريسودوايفيدرين بعد الوفاة، مما يجعل من المستحيل على علماء السموم وعلماء الأمراض وغيرهم من الأشخاص الذين يحققون في الوفيات المرتبطة بالقات أن يقرروا إذا كان المستوى ساما أو قاتلا. تعرض هذه مقالة جميع الأبحاث المنشورة المتاحة حالياً للمساعدة في البدء في توثيق هذا النطاق المهم. ينبغي جمع هذه المعلومات والإبلاغ عنها على أساس منهجي لتسهيل التفسيرات الصحيحة في المستقبل.

**Keywords:** Forensic Science, Khat, Deaths, Fatalities, Toxicology, Interpretation.

الكلمات المفتاحية: القات، الوفيات، علم السموم، التفسير



Production and hosting by NAUSS

\* Corresponding Author: John M. Corkery

Email: [j.corkery@herts.ac.uk](mailto:j.corkery@herts.ac.uk)

doi: xxxxxxxxxxxx

## 1. Introduction

The recent case report by Attafi et al. [1] raises the important issue of the need to take into account the post-mortem (re)-distribution of the key psychoactive constituents of khat (*Catha edulis* Forsk), i.e. cathine and cathinone and their precursors, when interpreting their levels in various human body tissues.

The sampling site may be affected by metabolism, especially if there is any hepatic or renal impairment. The levels of cathine, cathinone and related metabolites may also be affected by deterioration during the period between death and post-mortem sample collection, as well as storage methods, etc. Another important consideration in determining the toxic/lethal levels of these molecules is whether the decedent(s) had co-ingested any other drugs which may have given rise to drug-drug interactions and/or synergistic effects.

It is important that portmortem toxicology levels in 'deaths associated with khat consumption' are not only quantified, but also recorded and published. This will greatly assist in establishing what levels of these two molecules are toxic

and which are fatal. It is likely that there are many deaths which come under this heading, but they are not properly identified, recorded, analysed and published [2]. At present, the only information collated on such fatalities is presented in a book chapter [3] and two papers by the correspondent [2, 4], and the aforementioned paper [1]. This is summarised in Table-1.

This shows that the cathinone levels reported in the most recent case for both blood and vitreous humour are towards the low end of those previously reported by the correspondent. Perhaps, more importantly, the results presented show just how scant the published information is on this issue and, therefore, how difficult – if not impossible – it is to state with any confidence what constitutes a toxic or fatal level. One can only do that if there are sufficient cases reported with toxicological levels and where no other substances were found and/or implicated.

It would have been of interest if Attafi et al. [1] had provided the autopsy findings (including any information on hepatic or renal impairment and co-ingestion of other drugs), and the findings reached by the investigating officials in

**Table 1-** Summary of measured toxicological findings for the psychoactive constituents of khat (*Catha edulis* Forsk) in deceased humans.

Reference	Case number	Tissue levels ( $\mu\text{g/L}$ )		
		Norephedrine/ norpseudoephedrine	Cathine	Cathinone
1			Kidney - 310 Stomach - 3280	Brain - 30 Liver - 30 Stomach - 1850 Blood - 60 Vitreous humour - 70
3	2	Blood - 154		Blood - 104
3	5	Blood - 1000 Urine - detected		Blood - <50
3	9		Vitreous humour - 310	Vitreous humour - 110
3	10	Blood - <50 Urine - detected		Blood - <25 Urine - detected
3	13			Blood 19
3	15		Peripheral blood - 1447	Peripheral blood - 122



respect of the cause and manner of death, for example, accidental asphyxiation caused by the khat content in the decedent's mouth.

It is recommended that toxicologists and pathologists should bear in mind the need to build up the very limited published evidence-base we have on post-mortem cathine, cathinone and norephedrine/norpseudoephedrine tissue levels for deaths involving khat consumption.

### Acknowledgements

Thanks are due to Dr Eman Al-Saeed and Dr Zamzam Ahmed for technical assistance in translation.

### References

1. Attafi IM, Albeishy MY, Oraiby ME, Khardali IA, Shaikhain GA, Fageeh MM. Postmortem Distribution of Cathinone and Cathine in Human Biological Specimens in a Case of Death Associated with Khat Chewing. *Arab Journal of Forensic Sciences & Forensic Medicine (AJFSFM)*. 2018;1(7). <https://doi.org/10.26735/16586794.2018.013>
2. Corkery JM, Schifano F, Oyefeso A, Ghodse AH, Tonia T, Naidoo V, Button J. 'Bundle of fun' or 'bunch of problems'? Case series of khat-related deaths in the UK. *Drugs: education, prevention and policy*. 2011;18(6):408-25. <https://doi.org/10.3109/09687637.2010.504200>
3. Corkery JM. Khat—Chewing it over: Continuing 'cultural cement', cardiac challenge or catalyst for change?. *Forensic toxicology—Drug use and misuse*. 2016:165-207.
4. Corkery JM, Schifano F, Oyefeso A, Ghodse AH, Tonia T, Naidoo V, Button J. Overview of literature and information on "khat-related" mortality: a call for recognition of the issue and further research. *Annali dell'Istituto superiore di sanità*. 2011;47:445-64.

