

RAMSEY'S THEOREM AND PRODUCTS IN THE WEIHRAUCH DEGREES

DAMIR D. DZHAFAROV, JUN LE GOH, DENIS R. HIRSCHFELDT, LUDOVIC PATEY,
AND ARNO PAULY

ABSTRACT. We study the positions in the Weihrauch lattice of parallel products of various combinatorial principles related to Ramsey's theorem. Among other results, we obtain an answer to a question of Brattka, by showing that Ramsey's theorem for pairs (RT_2^2) is Weihrauch-incomparable to the parallel product of the stable Ramsey's theorem for pairs and the cohesive principle ($\text{SRT}_2^2 \times \text{COH}$).

1. INTRODUCTION

Reverse mathematics is a foundational area of logic devoted to calibrating the precise axioms needed to prove a given theorem of ordinary mathematics. For a standard reference, see Simpson [34]. A particularly fruitful line of research in this endeavor has been looking at theorems from combinatorics, particularly Ramsey's theorem and its many variants. See Hirschfeldt [20] for an introduction to the area. One recent way of extending the scope of this analysis is to replace the traditional framework of reverse mathematics, which is provability in fragments of second-order arithmetic, by Weihrauch reducibility. The latter is a tool that has been widely deployed in computable analysis and complexity theory; see the recent survey article by Brattka, Gherardi, and Pauly [4]. Recently it has gained prominence also in the study of computable combinatorics, and it is currently seeing a surge of activity; see, e.g., [1, 10, 13, 15, 18, 22, 24, 25, 28, 30, 31, 32]. See also Brattka [2] for an updated bibliography.

In this paper, we turn the lens of Weihrauch reducibility on various results concerning Ramsey's theorem and its products with other mathematical principles. We begin with some background on Weihrauch reducibility and Ramsey's theorem.

Definition 1.1. A *problem* P is a partial multifunction from 2^ω to 2^ω , written $P : \subseteq 2^\omega \rightrightarrows 2^\omega$. We call each $X \in \text{dom}(P)$ an *instance of* P , or P -*instance* for short, and each $Y \in P(X)$ a *solution to* X as an *instance of* P , or just a P -*solution to* X .

In general, a problem may be a partial multifunction between other kinds of *represented spaces*. We shall consider such problems in Section 4, and refer the reader

Dzhafarov was supported by grants DMS-1400267 and DMS-1854355 from the National Science Foundation of the United States and a Collaboration Grant for Mathematicians from the Simons Foundation. Goh was supported by NSF grant DMS-1161175. Hirschfeldt was supported by grants DMS-1101458, DMS-1600543, and DMS-1854279 from the National Science Foundation of the United States and a Collaboration Grant for Mathematicians from the Simons Foundation. All authors thank the Leibniz-Zentrum für Informatik at Schloss Dagstuhl, where the initial work for this project was conducted. We also thank Vasco Brattka for asking questions that motivated much of this work, and the anonymous referees for several highly useful comments.

to [4, Section 2] for definitions. Elsewhere in this paper, the above definition will be sufficient. (To be precise, we do work with represented spaces, since we code objects such as colorings of n -tuples of natural numbers as elements of Cantor space, but our codings are transparent enough that we can safely ignore this distinction, which we believe will improve clarity for most readers.)

We assume familiarity with standard computability-theoretic notation. For a partial function ψ , we write $\psi(x) \simeq y$ to mean that $\psi(x)$ is equal to y if defined.

A broad class of problems comes from reverse mathematics, where a typical object of study is a mathematical principle of the syntactic form

$$(\forall X)[\varphi(X) \rightarrow (\exists Y)[\theta(X, Y)]],$$

where φ and θ are arithmetical formulas of second-order arithmetic. Such a principle gives rise to the problem whose instances are the sets X such that $\varphi(X)$ holds, and where the solutions to any such X are the Y such that $\theta(X, Y)$ holds. In general, the formulas φ and θ above need not be unique for a given principle, but in practice, each principle one studies has a natural such pair of formulas associated to it. We adopt this terminology for specifying problems in this paper.

Definition 1.2. Let P and Q be problems.

- (1) Q is *computably reducible* to P , written $Q \leq_c P$, if every instance X of Q computes an instance \widehat{X} of P , such that for every solution \widehat{Y} to \widehat{X} , we have that $X \oplus \widehat{Y}$ computes a solution Y to X .
- (2) Q is *strongly computably reducible* to P , written $Q \leq_{sc} P$, if every instance X of Q computes an instance \widehat{X} of P , such that every solution \widehat{Y} to \widehat{X} computes a solution Y to X .
- (3) Q is *Weihrauch reducible* to P , written $Q \leq_w P$, if there exist Turing functionals Φ and Ψ such that for every instance X of Q , we have that Φ^X is an instance of P , and for every solution \widehat{Y} to Φ^X we have that $\Psi^{X \oplus \widehat{Y}}$ is a solution to X .
- (4) Q is *strongly Weihrauch reducible* to P , written $Q \leq_{sw} P$, if there exist Turing functionals Φ and Ψ such that for every instance X of Q , we have that Φ^X is an instance of P , and for every solution \widehat{Y} to Φ^X we have that $\Psi^{\widehat{Y}}$ is a solution to X .

We write $P \equiv_c Q$ if $P \leq_c Q$ and $Q \leq_c P$, and similarly for the other reducibilities above. All of these reducibilities are transitive, so the resulting notions of equivalence are in fact equivalence relations, which yield degree structures in the usual way. Figure 1 summarizes the relationships that hold between these reducibilities. We refer the reader to Hirschfeldt and Jockusch [22, Section 4.1] for a more thorough discussion of these reducibilities, and for various generalizations of them with applications to reverse mathematics.

The following two definitions list several important operations one can perform on problems.

Definition 1.3. Let P_0 and P_1 be problems.

- (1) The (*parallel*) *product* of P_0 and P_1 , written $P_0 \times P_1$, is the problem whose instances are pairs $\langle X_0, X_1 \rangle$ with X_i a P_i -instance, and where the solutions to $\langle X_0, X_1 \rangle$ are all pairs $\langle Y_0, Y_1 \rangle$ with Y_i a P_i -solution to X_i .




FIGURE 1. Relations between notions of reduction. An arrow from one reducibility to another means that whenever Q is reducible to P according to the first then it is also reducible according to the second. In general, no relations hold other than the ones shown.

- (2) The *coproduct* of P_0 with P_1 , written $P_0 \sqcup P_1$, is the problem whose instances are all pairs $\langle i, X \rangle$ for $i < 2$ such that X is a P_i -instance, and where the solutions to $\langle i, X \rangle$ are just the P_i -solutions to X .
- (3) The *meet* of P_0 with P_1 , written $P_0 \sqcap P_1$, is the problem whose instances are all pairs $\langle X_0, X_1 \rangle$ such that for each $i < 2$, X_i is a P_i -instance, and the solutions to $\langle X_0, X_1 \rangle$ are all pairs $\langle i, Y \rangle$ for $i < 2$ such that Y is a P_i -solution to X_i .

It is easy to see that the above operations lift to the \equiv_{w^-} , \equiv_{sW^-} , \equiv_{c^-} , and \equiv_{sc^-} -degrees. Furthermore, it is easy to see that the \equiv_{w^-} , \equiv_{c^-} , and \equiv_{sc^-} -degrees form a lattice with \sqcup as join and \sqcap as meet. Recently, Dzhafarov [16] has shown that the \equiv_{sW^-} -degrees also form a lattice, with \sqcap as meet but using a different operation for the join than \sqcup .

In this paper, all problems we consider will have some computable instance. It is easy to see that the coproduct of such problems is Weihrauch reducible to their parallel product.

Definition 1.4. Let P_0 and P_1 be problems.

- (1) The *composition* of P_1 with P_0 , written $P_1 \circ P_0$, is the problem whose instances are all the P_0 -instances X such that every solution to X is a P_1 -instance, and whose solutions to such an instance X are all the P_1 -solutions to the P_0 -solutions to X .
- (2) The *compositional product* of P_1 with P_0 , written $P_1 \star P_0$, is defined as $\max_{\leq_W} \{Q_1 \circ Q_0 : Q_i \leq_W P_i\}$.

The compositional product $P_1 \star P_0$, first defined by Brattka, Gherardi, and Marcone [3, Definition 4.1], captures exactly what can be achieved by applying P_0 and P_1 consecutively in series (possibly with some intermediate computation). Brattka and Pauly [9] showed that the compositional product is always defined. The definition of $P_1 \star P_0$ above does not yield a specific problem, of course, but only a Weihrauch degree. We will not use this notion except in the context of Weihrauch reducibility, however, so this fact will pose no problems.

The following proposition summarizes the relationships between the above operations on problems.

Proposition 1.5. *If P_0 and P_1 are problems with some computable instance, then*

$$P_0 \sqcap P_1 \leq_w P_0 \sqcup P_1 \leq_w P_0 \times P_1 \leq_w P_0 \star P_1.$$

To illustrate the definition of \star , we provide a proof of the last reduction. Observe that $P_0 \times P_1$ is the same problem as $(P_0 \times \text{id}) \circ (\text{id} \times P_1)$. Since $P_0 \times \text{id} \leq_w P_0$ and $P_1 \times \text{id} \leq_w P_1$, we have that $(P_0 \times \text{id}) \circ (\text{id} \times P_1) \leq_w P_0 \star P_1$, completing the proof.

Our focus here will be on Ramsey's theorem and its various combinatorial relatives. We begin with some definitions.

Definition 1.6. Let X be a subset of ω and k a positive number.

- (1) $[X]^2 = \{\langle x, y \rangle \in X \times X : x < y\}$.
- (2) A k -coloring of pairs (frequently just *coloring*) is a function $c : [\omega]^2 \rightarrow k$. We write $c(x, y)$ instead of $c(\langle x, y \rangle)$ for $\langle x, y \rangle \in [X]^2$. The coloring is *stable* if for every x there is an $i < k$ such that $c(x, y) = i$ for all sufficiently large y , in which case we write $\lim_y c(x, y) = i$.
- (3) A set $H \subseteq X$ is *homogeneous* for such a c if $c \upharpoonright [H]^2$ is constant. A set $Y \subseteq X$ is *almost homogeneous* for c if there is a finite set F such that $Y - F$ is homogeneous for c .
- (4) A set $L \subseteq X$ is *limit-homogeneous* for c if there is an $i < k$ such that $c(x, y) = i$ for all $x \in L$ and all sufficiently large $y \in L$, in which case we write $\lim_{y \in L} c(x, y) = i$. A set $Y \subseteq X$ is *almost limit-homogeneous* for c if there is a finite set F such that $Y - F$ is limit-homogeneous for c .

If $i < k$ is the color witnessing that some set is homogeneous or limit-homogeneous then we say the set is homogeneous/limit-homogeneous *with color i* . Note that if c is stable and L is limit-homogeneous for c with color i then also $\lim_y c(x, y) = i$ for all $x \in L$.

The following mathematical principles are well-known, and have been studied extensively in computability theory, reverse mathematics, and more recently, in the context of Weihrauch reducibility.

Ramsey's theorem for k -colorings of pairs (RT_k^2). *For every coloring $c : [\omega]^2 \rightarrow k$, there is an infinite homogeneous set for c .*

Stable Ramsey's theorem for k -colorings of pairs (SRT_k^2). *For every stable coloring $c : [\omega]^2 \rightarrow k$, there is an infinite homogeneous set for c .*

Δ_2^0 k -partition subset principle (D_k^2). *For every stable coloring $c : [\omega]^2 \rightarrow k$, there is an infinite limit-homogeneous set for c .*

(So, for concreteness, the instances of RT_k^2 are all colorings $c : [\omega]^2 \rightarrow k$, and the solutions to a given such c are its infinite homogeneous sets. Similarly for the other problems.) One additional principle that has been studied extensively alongside RT_k^2 and SRT_k^2 is the following:

Cohesive principle (COH). *For every sequence $\langle c_0, c_1, \dots \rangle$, where $c_i : \omega \rightarrow 2$ for each $i \in \omega$, there exists an infinite set X that is almost homogeneous for each c_i .*

It is an easy exercise to see that D_k^2 is strongly Weihrauch equivalent to the problem asserting that for every Δ_2^0 k -partition $\langle A_0, \dots, A_{k-1} \rangle$ of ω , there exists an infinite subset X of some A_i , and in the sequel, we will use whichever formulation is more convenient. It is obvious that $\text{D}_k^2 \leq_{\text{sw}} \text{SRT}_k^2 \leq_{\text{sw}} \text{RT}_k^2$. While every

computable instance of SRT_k^2 has a Δ_2^0 solution, Jockusch [26, Theorem 3.1] constructed a computable instance of RT_k^2 with no Δ_2^0 solution. Thus, $\text{RT}_k^2 \not\leq_W \text{SRT}_k^2$. Dzhafarov [15, Corollary 3.3] showed that $\text{SRT}_k^2 \not\leq_W \text{D}_k^2$. An independent proof can be found in [10, Corollary 6.12]. Note that if $j < k$ then the version of each of the above principles for j -colorings is strongly Weihrauch reducible to the version for k -colorings. Patey [32] showed that the converse is false; in fact, if $j < k$, then even $\text{D}_k^2 \not\leq_c \text{RT}_j^2$. Further relationships between SRT_2^2 , D_2^2 , and related principles under the various reductions from Definition 1.2 have been investigated by Nichols [30].

Definition 1.7. For a problem P , let P^{FE} be the problem whose instances are the same as those of P , but such that Y is a P^{FE} -solution to X if there is a P -solution Z to X such that $Y =^* Z$ (i.e., such that Y and Z agree on a cofinite domain).

Thus, for instance, $(\text{D}_k^2)^{\text{FE}}$ asserts that every stable coloring $c : [\omega]^2 \rightarrow k$ has an infinite almost limit-homogeneous set. For some well-behaved principles P , we can express P^{FE} in terms of the *implication operation* introduced by Brattka and Pauly in [9, Section 3.3]. In lieu of a definition, we use the following property (see [9, Theorem 3.13]): for problems P and Q , the infimum $\inf_{\leq_W} \{R : P \leq_W Q \star R\}$ exists and is Weihrauch equivalent to $Q \rightarrow P$. We also recall the following *choice principle* (see [4, Section 7]).

Definition 1.8. $C_{\mathbb{N}}$ is the problem whose instances are functions $e : \omega^2 \rightarrow 2$ such that

- for all x , $e(x, 0) = 0$ and there is at most one s with $e(x, s) \neq e(x, s + 1)$;
- there is at least one x with $e(x, s) = 0$ for all s .

A solution to such an e is any $x \in \omega$ such that $e(x, s) = 0$ for all s .

Thus the instances of $C_{\mathbb{N}}$ are enumerations of sets with nonempty complements, and the solutions are the elements of these complements.

Proposition 1.9. Let $P \in \{\text{RT}_k^2, \text{SRT}_k^2\}$. Then $P^{\text{FE}} \equiv_W C_{\mathbb{N}} \rightarrow P$.

Proof. To show that $P^{\text{FE}} \leq_W C_{\mathbb{N}} \rightarrow P$, it suffices to show that for any R such that $P \leq_W C_{\mathbb{N}} \star R$, we have that $P^{\text{FE}} \leq_W R$. Equivalently, we will show that for any $Q \leq_W C_{\mathbb{N}}$ and R such that $P \leq_W Q \circ R$, we have that $P^{\text{FE}} \leq_W R$. Let Φ and Ψ witness that $P \leq_W Q \circ R$. Let Γ and Δ witness that $Q \leq_W C_{\mathbb{N}}$. We describe a uniform procedure for reducing P^{FE} to R . Given a P -instance c , we use Φ to convert this to an instance X of $Q \circ R$. Any R -solution Y to X is also a Q -instance, so we can use Γ to convert it into an instance Z of $C_{\mathbb{N}}$. More precisely, Γ^Y enumerates a set \bar{Z} such that $Z \neq \emptyset$. And given any $C_{\mathbb{N}}$ -solution to Z , i.e., a point $x \in Z$, $\Delta(\langle Y, x \rangle)$ is a Q -solution to Y . Hence $\Psi(\langle c, \Delta(\langle Y, x \rangle) \rangle)$ must be a solution to P . Thus, to uniformly compute a P^{FE} -solution H to c from a given R -solution Y to X , we proceed as follows. To determine $H(n)$, we choose the least x not yet enumerated by Γ^Y at stage n , and wait for x either to be enumerated, in which case we let $H(n) = 0$, or for $\Psi(\langle c, \Delta(\langle Y, x \rangle) \rangle)(n)$ to converge, in which case we let $H(n) = \Psi(\langle c, \Delta(\langle Y, x \rangle) \rangle)(n)$. It is easy to see that H is then an infinite set and is almost homogeneous for c .

In the other direction, it suffices to show that $P \leq_W C_{\mathbb{N}} \star P^{\text{FE}}$. Consider the following uniform procedure. Given an instance $c : [\omega]^2 \rightarrow k$ of P , we regard it also as an instance of P^{FE} . Now, given any P^{FE} -solution Y to c , i.e., an infinite almost homogeneous set, define

$$Z = \{x \in Y : (\exists i < k)(\forall y \geq x)(\forall z > y)[y, z \in Y \rightarrow c(y, z) = i]\}.$$

Note that Z agrees with Y on all but finitely many elements, and so is in particular nonempty. Moreover, Z is a $\Pi_1^{0,Y}$ subset of \mathbb{N} , and hence can be passed as an instance to $\mathbb{C}_{\mathbb{N}}$. Let x be any $\mathbb{C}_{\mathbb{N}}$ -solution to this instance. Then $\{y \in Y : y \geq x\}$ is a \mathbb{P} -solution to c .

We use the above uniform procedure to show that $\mathbb{P} \leq_w \mathbb{C}_{\mathbb{N}} \star \mathbb{P}^{\text{FE}}$. Let G be a computable function that takes in a pair $\langle c, Y \rangle$ and produces an enumeration of the complement of Z , as defined above. Then the above procedure shows that $\mathbb{P} \leq_w \mathbb{C}_{\mathbb{N}} \circ G \circ (\text{id} \times \mathbb{P}^{\text{FE}})$. Since $G \circ (\text{id} \times \mathbb{P}^{\text{FE}}) \leq_w \mathbb{P}^{\text{FE}}$, this proves the desired result. \square

2. RAMSEY'S THEOREM FOR PAIRS

Our starting point is the following summary of known facts concerning relationships between RT_2^2 , SRT_2^2 , and COH under Weihrauch reducibility.

Theorem 2.1.

- (1) $\text{SRT}_2^2 \sqcup \text{COH} \leq_w \text{RT}_2^2 \leq_w \text{SRT}_2^2 \star \text{COH}$;
- (2) $\text{SRT}_2^2 \sqcup \text{COH} \leq_w \text{SRT}_2^2 \times \text{COH} \leq_w \text{SRT}_2^2 \star \text{COH}$.

Proof. Part (1) follows by the proof of Cholak, Jockusch, and Slaman [11, Lemma 7.11] that $\text{SRT}_2^2 \wedge \text{COH}$ and RT_2^2 are equivalent in the formal system RCA_0 , together with the proof of their Theorem 12.5, which is needed because the argument that RT_2^2 implies COH in the proof of Lemma 7.11 was not correct, as noted in [12] (see also [13, Section 5.2]). Part (2) follows from Proposition 1.5. \square

Our main motivation for this section is the following question, asked during the workshop ‘‘Measuring the Complexity of Computational Content: Weihrauch Reducibility and Reverse Analysis’’, at the Leibniz-Zentrum für Informatik at Schloss Dagstuhl in September, 2015.

Question 2.2 (Brattka, see [7]). What additional reductions hold between the problems $\text{SRT}_2^2 \sqcup \text{COH}$, $\text{SRT}_2^2 \star \text{COH}$, $\text{SRT}_2^2 \times \text{COH}$, and RT_2^2 in Theorem 2.1?

We will answer this question by showing that RT_2^2 and $\text{SRT}_2^2 \times \text{COH}$ are Weihrauch-incomparable, and hence there are no Weihrauch reductions between the above principles other than the ones given in Theorem 2.1.

We begin by recalling some ancillary notions.

Definition 2.3. Let $c : [\omega]^2 \rightarrow k$ be a coloring, and let X be a set.

- (1) The coloring c is *unbalanced on X* if for some $i < k$, every infinite homogeneous set for c contained in X has color i . If c is not unbalanced on X , it is *balanced on X* .
- (2) The coloring c *avoids the color $i < k$ on X* if $c(x, y) \neq i$ for all $x, y \in X$.

If, in the definition above, $X = \omega$, we shall say simply that c is unbalanced / balanced / avoids the color i , without further qualification.

The following lemma will allow us to prove our main result, from which we will derive a number of consequences, including an answer to Question 2.2.

Lemma 2.4. *Let $c : [\omega]^2 \rightarrow k$ be a computable coloring, A an infinite computable set, and $\mathcal{C} \subseteq 2^\omega$ a nonempty Π_1^0 class of k -partitions of A . If, for every $\langle P_0, \dots, P_{k-1} \rangle \in \mathcal{C}$, c is unbalanced on P_j for every $j < k$, then c has a computable infinite homogeneous set.*

Proof. Fix c and \mathcal{C} , and suppose that c has no computable infinite homogeneous set. We construct a set $G = \{G_{i,j} : i, j < k\}$, and exhibit a $\langle P_0, \dots, P_{k-1} \rangle \in \mathcal{C}$, such that $G_{i,j} \subseteq P_j$ for all $i, j < k$, and c avoids the color i on $G_{i,j}$. We will furthermore satisfy the following requirement for each $n \in \omega$ and all $\alpha \in k^k$:

$$\mathcal{R}_{n,\alpha} : (\exists j < k)(\exists x \geq n)[x \in G_{\alpha(j),j}].$$

The claim is that c is then balanced on some P_j . For if not, define $\alpha \in k^k$ by letting $\alpha(j)$ be the color $i < k$ such that every infinite homogeneous set for c contained in P_j has color i . Since G satisfies $\mathcal{R}_{n,\alpha}$ for all n , there must be a $j < k$ such that $G_{\alpha(j),j}$ is infinite. Let H be any infinite homogeneous set for c contained in $G_{\alpha(j),j}$. As c avoids the color $\alpha(j)$ on $G_{\alpha(j),j}$, it follows that H has some other color than $\alpha(j)$, which is a contradiction since $G_{\alpha(j),j} \subseteq P_j$.

The construction of G is by a forcing notion whose conditions are tuples

$$p = (\{E_{i,j} : i, j < k\}, X, \mathcal{D}),$$

such that for all $i, j < k$:

- $E_{i,j}$ is a finite subset of A ;
- X is a computable infinite subset of A such that $\max E_{i,j} < \min X$;
- for every $x \in X$, c avoids the color i on $E_{i,j} \cup \{x\}$;
- \mathcal{D} is a nonempty Π_1^0 subclass of \mathcal{C} such that for every $\langle P_0, \dots, P_{k-1} \rangle \in \mathcal{D}$, $E_{i,j} \subseteq P_j$.

A condition $q = (\{F_{i,j} : i, j < k\}, Y, \mathcal{E})$ extends p if $Y \subseteq X$, $\mathcal{E} \subseteq \mathcal{D}$, and $E_{i,j} \subseteq F_{i,j} \subseteq E_{i,j} \cup X$ for all $i, j < k$.

Say a condition p as above *satisfies* $\mathcal{R}_{n,\alpha}$ if there are some $j < k$ and some $x \geq n$ such that $x \in E_{\alpha(j),j}$. We claim that the set of conditions satisfying $\mathcal{R}_{n,\alpha}$ is dense. Fix $p = (\{E_{i,j} : i, j < k\}, X, \mathcal{D})$. First, suppose there are some $\langle Q_0, \dots, Q_{k-1} \rangle \in \mathcal{D}$, some $\ell < k$, and some $x \in X \cap Q_\ell$ such that $Y = \{y \in X : c(x, y) \neq \alpha(\ell)\}$ is infinite. Let $q = (\{F_{i,j} : i, j < k\}, Y, \mathcal{E})$, where $F_{\alpha(\ell),\ell} = E_{\alpha(\ell),\ell} \cup \{x\}$, $F_{i,j} = E_{i,j}$ for all $i, j < k$ with $i \neq \alpha(\ell)$ or $j \neq \ell$, and $\mathcal{E} = \{\langle P_0, \dots, P_{k-1} \rangle \in \mathcal{D} : x \in P_\ell\}$. Then q is an extension of p satisfying $\mathcal{R}_{n,\alpha}$. So suppose now that there are no such $\langle Q_0, \dots, Q_{k-1} \rangle$, ℓ , and x . We derive a contradiction. The assumption implies that for every $x \in X$, $\lim_{y \in X} c(x, y)$ exists, since given any $\langle Q_0, \dots, Q_{k-1} \rangle \in \mathcal{D}$, we have that $\lim_{y \in X} c(x, y) = \alpha(\ell)$ for the unique ℓ with $x \in Q_\ell$. So the map $g : X \rightarrow k$ defined by $g(x) = \lim_{y \in X} c(x, y)$ for all $x \in X$ is computable from every member of \mathcal{D} . By the cone-avoidance basis theorem (see, e.g., [17, Theorem 2.1]), this implies that g is computable. But then c has a computable infinite homogeneous set, which we assumed it did not.

To complete the proof, let $\mathcal{G} = \langle q_0, q_1, \dots \rangle$ be a sufficiently generic sequence on our forcing poset, where for each s ,

$$q_s = (\{E_{i,j}^s : i, j < k\}, X^s, \mathcal{D}^s),$$

and q_s is extended by q_{s+1} . Define

$$G_{i,j} = \bigcup_{s \in \omega} E_{i,j}^s$$

for all $i, j < k$. Let $\langle P_0, \dots, P_{k-1} \rangle$ be any element of $\bigcap_{s \in \omega} \mathcal{D}_s$, which is an intersection of a nested sequence of Π_1^0 classes and hence is nonempty. Then $G = \{G_{i,j} : i, j < k\}$ and $\langle P_0, \dots, P_{k-1} \rangle$ have the desired properties. \square

The following important problems arise frequently in the study of Weihrauch degrees.

Definition 2.5.

- (1) LPO is the principle whose instances are all infinite binary sequences of the form 0^ω or $0^n 1^\omega$ for some $n \in \omega$, and the solutions are either the singleton $\{0\}$ if the instance is 0^ω , or $\{1\}$ if the instance is $0^n 1^\omega$ for some n .
- (2) NON is the principle whose instances are all sets, and the solutions to an instance X are all sets $Y \not\leq_T X$.

Viewed as a Π_2^1 principle, NON is thus equivalent over RCA_0 to the principle AST considered by Hirschfeldt, Shore, and Slaman [23, Section 6]. (See specifically [23, Theorem 6.3].)

Theorem 2.6. $\text{LPO} \times \text{NON} \not\leq_W \text{RT}_2^2$.

Proof. Assume otherwise, and fix functionals Φ and Ψ witnessing the reduction. We build an instance S of LPO such that the pair $\langle S, \emptyset \rangle$ contradicts this assumption. We have that $\Phi^{S \oplus \emptyset}$ is a coloring $[\omega]^2 \rightarrow 2$, and for every infinite homogeneous set H for this coloring, $\Psi^{S \oplus \emptyset \oplus H} = \langle \{b\}, Y \rangle$, where b is 0 or 1 depending on whether $S = 0^\omega$ or $S = 0^n 1^\omega$ for some n , and $Y \not\leq_T \emptyset$. (We think of $\Psi^{S \oplus \emptyset \oplus H}$ as the characteristic function of $\{b\} \oplus Y$, so that $\Psi^{S \oplus \emptyset \oplus H}(0) \downarrow = 1$ if and only if $b = 0$.) We show that the coloring $\Phi^{S \oplus \emptyset}$ necessarily has an infinite homogeneous set H satisfying one of the following properties:

- (1) H is computable;
- (2) $\Psi^{S \oplus \emptyset \oplus H}(0) \simeq 0$ and $S = 0^\omega$;
- (3) $\Psi^{S \oplus \emptyset \oplus H}(0) \downarrow = 1$ and $S = 0^n 1^\omega$ for some n .

In the first case, $S \oplus \emptyset \oplus H$ obviously cannot compute a solution to our NON-instance. And in the remaining cases, we have a contradiction to $\Psi^{S \oplus \emptyset \oplus H}$ giving us a solution to our LPO-instance.

Let c be the coloring $\Phi^{0^\omega \oplus \emptyset} : [\omega]^2 \rightarrow 2$. Define \mathcal{C} to be the Π_1^0 class consisting of all 2-partitions $\langle P_0, P_1 \rangle$ of ω such that

$$(\forall i < 2)(\forall \text{ finite } F \subseteq P_i)[(\forall x, y \in F)[c(x, y) = i] \rightarrow \Psi^{0^\omega \oplus \emptyset \oplus F}(0) \simeq 0].$$

We consider two cases. First, suppose \mathcal{C} is nonempty. By Lemma 2.4 with $k = 2$ and $A = \omega$, if c is unbalanced on P_0 and P_1 for every $\langle P_0, P_1 \rangle \in \mathcal{C}$, then c has a computable infinite homogeneous set. We can then take this to be H , set $S = 0^\omega$, and satisfy Property (1) above. So assume not. Fix $\langle P_0, P_1 \rangle \in \mathcal{C}$ and $i < 2$ such that c is balanced on P_i , so that in particular, P_i is infinite. Let $H \subseteq P_i$ be any infinite homogeneous set for c with color i . If we then take $S = 0^\omega$, it follows by the definition of \mathcal{C} that $\Psi^{S \oplus \emptyset \oplus H}(0) \simeq 0$, so we satisfy Property (2).

So now, suppose $\mathcal{C} = \emptyset$. By compactness, choose m so that for every partition $\langle P_0, P_1 \rangle$ of ω , there are an $i < 2$ and a finite $F \subseteq P_i \upharpoonright m$ such that $c(x, y) = i$ for all $x, y \in F$ and $\Psi^{0^\omega \oplus \emptyset \oplus F}(0) \downarrow = 1$. Note that there are only finitely many such F across all possible partitions, so there is a global bound u on the uses of all these computations. Without loss of generality, $u \geq m$. Choose $n > u$ large enough so that $\Phi^{0^n 1^\omega \oplus \emptyset}$ agrees with $c = \Phi^{0^\omega \oplus \emptyset}$ below u . Let $S = 0^n 1^\omega$, and let $d = \Phi^{S \oplus \emptyset}$. By repeatedly taking subsets, we see that there is a computable infinite set Y such that $\min Y > m$ and for each $x < m$, $\lim_{y \in Y} d(x, y)$ exists. For each $i < 2$, let $Q_i = \{x < m : \lim_{y \in Y} d(x, y) = i\}$, so that for some partition $\langle P_0, P_1 \rangle$ of ω , we have

$Q_0 = P_0 \upharpoonright m$ and $Q_1 = P_1 \upharpoonright m$. Choose $i < 2$ and $F \subseteq Q_i$ as above. If Y contains no infinite homogeneous set for d with color i , then by [26, Theorem 5.11], Y contains a computable infinite homogeneous set with color $1 - i$, and we satisfy Property (1) again. Otherwise, we can take an infinite homogeneous set H for d having F as an initial segment, and by construction, this set satisfies $\Psi^{S \oplus \emptyset \oplus H}(0) \downarrow = 1$ even though $S \neq 0^\omega$. Thus, we satisfy Property (3). \square

Trivially, $\text{LPO} \leq_w \text{RT}_2^1$, since every instance of LPO can be regarded as an instance of RT_2^1 , so we have the following.

Corollary 2.7. $\text{RT}_2^1 \times \text{NON} \not\leq_w \text{RT}_2^2$.

Clearly $\text{RT}_2^1 \leq_w \text{SRT}_2^2$, and there is a uniform construction of an X -computable instance of COH with no X -computable solution, so we also have the following.

Corollary 2.8. $\text{SRT}_2^2 \times \text{COH} \not\leq_w \text{RT}_2^2$.

Corollary 2.9. *No additional relations hold between the problems in Theorem 2.1.*

Proof. Since every computable instance of each of COH and SRT_2^2 admits a Δ_2^0 solution, so does every instance of $\text{COH} \times \text{SRT}_2^2$. By contrast, it is known that RT_2^2 has a computable instance with no Δ_2^0 solution. Thus, $\text{RT}_2^2 \not\leq_w \text{COH} \times \text{SRT}_2^2$. The remaining non-reductions follow from this fact and Corollary 2.8 by transitivity. \square

Let lim be the problem where an instance is a convergent sequence of elements of $\mathbb{N}^{\mathbb{N}}$, and the unique solution to this problem is the limit of this sequence. The following fact about lim is well-known, but it is worth mentioning that it follows directly from Theorem 2.6, since $\text{LPO} \times \text{NON} \leq_w \text{lim}$.

Corollary 2.10. $\text{lim} \not\leq_w \text{RT}_2^2$.

It is an interesting open question whether LPO can be replaced by an even weaker combinatorial principle in Theorem 2.6. A first candidate would be $\text{LLPO} \equiv_w \text{C}_2$ (see [5, §3]). Already $\text{LLPO} \times \text{NON} \not\leq_w \text{RT}_2^2$ would imply $\text{WKL} \not\leq_w \text{RT}_2^2$, which is known by Liu's celebrated result [27]. An even further improvement to $\text{ACC}_n \times \text{NON} \not\leq_w \text{RT}_2^2$ would have as a consequence that $\text{DNC}_n \not\leq_w \text{RT}_2^2$ (see [5, §3, §5], which also has definitions of ACC_X and DNC_X). On the other hand, it is the case that $\text{ACC}_{\mathbb{N}} \times \text{NON} \leq_w \text{DNC}_{\mathbb{N}} \leq_w \text{RT}_2^2$ (the latter reduction having been shown in [21, Theorem 2.3]).¹

Note that Theorem 2.6 also cannot be improved to show that $\text{LPO} \times \text{NON} \not\leq_w \text{RT}_k^2$ for arbitrary $k \geq 2$. Indeed, each of LPO and NON is Weihrauch reducible to RT_2^2 , and so $\text{LPO} \times \text{NON} \leq_w \text{RT}_2^2 \times \text{RT}_2^2 \leq_w \text{RT}_4^2$.

We can improve on this reduction with the following strong counterpoint to Theorem 2.6, which shows that the theorem fails as soon as the number of colors is allowed to increase from two, even via a stable coloring. It is easy to see that $\text{LPO} \leq_w \text{SRT}_2^1$ and $\text{NON} \leq_w \text{SRT}_2^2$, so the following result also follows from [10, Theorem 3.24], which has as a special case that $\text{SRT}_2^1 \times \text{SRT}_2^2 \leq_w \text{SRT}_3^2$.

Proposition 2.11. $\text{LPO} \times \text{NON} \leq_w \text{SRT}_3^2$.

¹The remarks in this paragraph were kindly provided by an anonymous referee.

Proof. Let S be an arbitrary instance of LPO, and let X be any set. Let $c : [\omega]^2 \rightarrow 2$ be the result of applying a standard uniform construction of an X -computable stable coloring $c : [\omega]^2 \rightarrow 2$ with no X -computable infinite homogeneous set (e.g., as in [26, Theorem 2.1]). Define $d : [\omega]^2 \rightarrow 3$ by

$$d(x, y) = \begin{cases} c(x, y) & \text{if } S(x) = S(y), \\ 2 & \text{otherwise.} \end{cases}$$

Clearly, d is uniformly computable from $S \oplus X$. If $S = 0^\omega$ then $d = c$, while if $S = 0^n 1^\omega$ for some n then $\lim_y d(x, y) = 2$ for all $x < n$, and $d(x, y) = c(x, y)$ for all $x \geq n$. Hence, every infinite homogeneous set for d has color 0 or 1, and is also homogeneous for c . In particular, no infinite homogeneous set for d is X -computable. Moreover, for any such infinite homogeneous set H , we have that $S = 0^n 1^\omega$ if and only if $(\exists x \leq \min H)[S(x) = 1]$. Hence, $\langle \{b\}, H \rangle$, where b is 0 or 1 depending as $S = 0^\omega$ or $S = 0^n 1^\omega$ for some n , is a uniformly $(S \oplus X \oplus H)$ -computable solution to the LPO \times NON-instance $\langle S, X \rangle$. \square

We do not know whether SRT_3^2 above can be replaced by D_3^2 . However, we have the following related result, which does work for D_3^2 . The proof uses a novel coding mechanism.

Theorem 2.12. *For every $k \geq 1$, $(\text{RT}_k^1)^{\text{FE}} \times \text{NON} \leq_w \text{D}_{k+1}^2$.*

Proof. Let $c : \omega \rightarrow k$ be a coloring and X a set. We describe a uniform procedure to define an X -computable stable coloring $d : [\omega]^2 \rightarrow k + 1$ with no X -computable solution (i.e., no X -computable infinite limit-homogeneous set), and a uniform procedure for turning any such solution into an almost limit-homogeneous set for c . Fix a canonical $(c \oplus X)$ -computable enumeration of $(c \oplus X)'$, and let

$$s_0 < s_1 < \dots$$

be a $(c \oplus X)'$ -computable sequence such that for all e , we have that $(c \oplus X)'[s_e]$ and $(c \oplus X)'$ agree on all $x \leq e$. Using $(c \oplus X)'$, choose

$$x_{0,0} < x_{0,1} < x_{1,0} < x_{1,1} < x_{2,0} < x_{2,1} < \dots$$

with $x_{e+1,0} - x_{e,1} \geq s_e$ for all e , and such that either $\Phi_e^X(x_{e,0}) \downarrow = \Phi_e^X(x_{e,1}) \downarrow = 1$, or $\Phi_e^X(x) \simeq 0$ for all sufficiently large x .

Now define a $(c \oplus X)'$ -computable $(k + 1)$ -partition $P_0 \cup \dots \cup P_k$ of ω as follows. For each e , put $x_{e,0}$ into P_k , and put every other x into $P_{c(x)}$. Thus, for all e , we have that $x_{e,0}$ and $x_{e,1}$ belong to different parts of the partition, so by construction, if Φ_e^X defines an infinite set, this set cannot be an infinite subset of any P_i . We also have that if $z_0 < \dots < z_{n-1} \in P_k$ then $z_{e+1} - z_e \geq s_e$ for all $e < n$, so any infinite subset of P_k computes $(c \oplus X)'$. We can regard $P_0 \cup \dots \cup P_k$ as a $(c \oplus X)$ -computable stable coloring d . Clearly, d is defined uniformly from X and c , and no infinite limit-homogeneous set for d is X -computable.

Consider any D_{k+1}^2 -solution to d , i.e., any infinite set $Z = \{z_0 < z_1 < \dots\}$ contained in one of P_0, \dots, P_k . We construct a set $Y = \{y_0 < y_1 < \dots\}$ inductively by stages, defining y_n at stage n . At any stage, we may choose to *exit the construction*, which simply means to let m be the maximum of all y_n defined thus far, and let the rest of our set be $\{z_n \in Z : z_n > m\}$. At stage $n = 0$, let $y_0 = 0$, and declare no color $i < k$ *forbidden*. If we have not exited the construction by stage $n + 1$, assume we have defined y_n and there is at least one $i < k$ that is still not forbidden.

For the least such i , we compute a number e such that $e \in (c \oplus X)'$ if and only if $(\exists y > y_n)[c(y) = i]$, which we can do uniformly from y_n , the color i , and an index for c as a $(c \oplus X)$ -computable coloring. If $e \in (c \oplus X)'[z_{e+1} - z_e]$ then certainly $e \in (c \oplus X)'$, so we can find a $y > y_n$ with $c(y) = i$, and we let y_{n+1} be the least such y . If $e \notin (c \oplus X)'[z_{e+1} - z_e]$, we declare i forbidden and restart the process with the next smallest non-forbidden color. In this case, we promise that if at any future stage we see a $y > y_n$ with $c(y) = i$, we exit the construction. Note that this can happen only if $z_{e+1} - z_e < s_e$. Note also that it must happen if all $i < k$ become forbidden.

It is easy to see that Y is uniformly computable from $c \oplus X \oplus Z$. We claim that Y is almost limit-homogeneous for c . This is clear if we never exit the construction, because in that case there must be some least i that is never declared forbidden, and then $c(y_n) = i$ for almost all n . If, on the other hand, we do exit the construction, then as noted above we must have $z_{e+1} - z_e < s_e$ for some e , and hence Z cannot be a subset of P_k . In this case, Z is therefore a subset of P_i for some $i < k$, and by construction, if $x \in P_i$ for such an i then $c(x) = i$. As $Y =^* Z$, it follows that Y is almost limit-homogeneous for c . \square

3. STABLE RAMSEY'S THEOREM FOR PAIRS

As mentioned above, every instance of LPO can be regarded as an instance of RT_2^1 . The latter instance, however, is consequently unbalanced. It is interesting to ask whether this is the only possible reduction, or whether LPO can in fact be reduced to RT_2^1 via a balanced coloring. The following proposition shows that the answer is no. It also points to an additional point of disagreement in the uniform strengths of SRT_2^2 and D_2^2 , to complement the aforementioned result that $\text{SRT}_2^2 \not\leq_{\text{W}} \text{D}_k^2$ for all k .

We first give a definition.

Definition 3.1. For $P \in \{\text{RT}_k^1, \text{SRT}_k^2, \text{D}_k^2\}$, let $\mathbf{b}\text{-}P$ be the restriction of P to balanced colorings (on ω), and $\mathbf{u}\text{-}P$ the restriction to unbalanced colorings.

Proposition 3.2. $\text{LPO} \leq_{\text{W}} \mathbf{b}\text{-SRT}_2^2$, but $\text{LPO} \not\leq_{\text{W}} \mathbf{b}\text{-D}_k^2$ for all k .

Proof. For the positive reduction, let S be any instance of LPO. Let $\text{par}(x)$ be 0 or 1 depending on whether x is even or odd, and define $c : [\omega]^2 \rightarrow 2$ by

$$c(x, y) = \begin{cases} \text{par}(x) & \text{if } (\forall z < y)[S(z) = 0], \\ 1 - \text{par}(x) & \text{otherwise.} \end{cases}$$

Thus, if $S = 0^\omega$ then $\lim_y c(x, y) = \text{par}(x)$ for all x , and if $S = 0^n 1^\omega$ for some n then $\lim_y c(x, y) = 1 - \text{par}(x)$ for all x . In either case, for each $i < 2$, there are infinitely many x with $\lim_y c(x, y) = i$, so c is balanced. Now, every element in an infinite homogeneous set for c has the same parity. So if H is any such homogeneous set, and if x_0 and x_1 are its least two elements, then $S = 0^\omega$ if and only if $c(x_0, x_1) = \text{par}(x_0)$. Thus, we have the desired uniform reduction.

For the negative reduction, assume towards a contradiction that $\text{LPO} \leq_{\text{W}} \mathbf{b}\text{-D}_k^2$ via some Φ and Ψ . Then $c = \Phi^{0^\omega}$ is a computable balanced stable coloring $[\omega]^2 \rightarrow k$. Define \mathcal{C} to be the Π_1^0 class of all partitions $\langle P_0, \dots, P_{k-1} \rangle$ of ω such that

$$(\forall i < k)(\forall \text{ finite } F \subseteq P_i)[\Psi^{0^\omega \oplus F}(0) \simeq 0].$$

First, we claim that $\mathcal{C} \neq \emptyset$. Otherwise, by compactness, there is an m such that for every partition $\langle P_0, \dots, P_{k-1} \rangle$ of ω , there are an $i < k$ and a finite $F \subseteq P_i \upharpoonright m$ such that $\Psi^{0^\omega \oplus F}(0) \downarrow = 1$. Let $u \geq m$ be a bound on the uses of all these computations, for all possible such F . Choose $n > u$ large enough so that $\Phi^{0^n 1^\omega}$ and $c = \Phi^{0^\omega}$ agree below u . Let $S = 0^n 1^\omega$ and $d = \Phi^S$, which is another balanced stable coloring. For the partition $\langle P_0, \dots, P_{k-1} \rangle$ of ω given by $P_i = \{x : \lim_y d(x, y) = i\}$, fix $i < k$ and $F \subseteq P_i \upharpoonright m$ as above. As d is balanced, there is an infinite homogeneous set H for d with color i that has F as an initial segment. But then we have $\Psi^{S \oplus H}(0) \downarrow = 1$ even though $S \neq 0^\omega$, a contradiction. So $\mathcal{C} \neq \emptyset$. Choose any $\langle P_0, \dots, P_{k-1} \rangle \in \mathcal{C}$. Since $P_0 \cup \dots \cup P_{k-1} = \omega$, there is an $i < k$ such that P_i is infinite. Let L be any infinite limit-homogeneous set for c contained in P_i . Then $\Psi^{0^\omega \oplus L}(0) \simeq 0$, which contradicts the choice of Ψ . \square

Corollary 3.3. $\text{LPO} \not\leq_{\text{w}} \text{b-RT}_k^1$ for all k .

Proof. The usual proof that $\text{RT}_k^1 \leq_{\text{w}} \text{D}_k^2$ shows that $\text{b-RT}_k^1 \leq_{\text{w}} \text{b-D}_k^2$. \square

One generalization of the notion of unbalanced coloring is the following, in which merely one of the possible colors of homogeneous sets—rather than, all but one—is omitted.

Definition 3.4. Let $c : [\omega]^2 \rightarrow k$ be a coloring and X a set. The coloring c is *thin-unbalanced on X* if for some $i < k$, there is no infinite homogeneous set for c contained in X with color i . The color i is called a *witness of thin-unbalancing for c on X* . If c is not thin-unbalanced on X , it is *thin-balanced on X* .

When $X = \omega$, we shall simply say c is thin-unbalanced / thin-balanced. Note that if $k = 2$, then c is thin-unbalanced on a set if and only if it is unbalanced on that set in the sense of Definition 2.3, which in turn holds if and only if c avoids one of its two colors on that set.

Definition 3.5. For $\text{P} \in \{\text{SRT}_k^2, \text{D}_k^2\}$, we define the following variations on P :

- $\Delta_2^0\text{-wtu-P}$ is the problem whose instances are pairs $\langle c, \ell \rangle$ where c is a thin-unbalanced instance of P and $\ell : \omega \rightarrow k$ is a function such that $\lim_y \ell(y)$ exists and is a witness of thin-unbalancing for c , and the solutions to such a pair are the P -solutions to c .
- wtu-P is the problem whose instances are pairs $\langle c, i \rangle$ where c is a thin-unbalanced instance of P and i is a witness of thin-unbalancing for c , and the solutions to such a pair are the P -solutions to c .

The above are arguably not natural problems from a combinatorial point of view, and we will not study them in their own right. Rather, our interest is in what they can reveal about SRT_k^2 and D_k^2 . As we will see, the above restrictions capture various elements of standard proofs of the latter principles.

Proposition 3.6.

- (1) For $\text{P} \in \{\text{SRT}_k^2, \text{D}_k^2\}$, we have $\text{LPO} \leq_{\text{w}} \text{wtu-P} \leq_{\text{sW}} \Delta_2^0\text{-wtu-P}$.
- (2) $\text{wtu-SRT}_k^2 \times \text{wtu-SRT}_2^2 \leq_{\text{sW}} \text{wtu-SRT}_k^2$ and $\text{wtu-D}_k^2 \times \text{wtu-D}_2^2 \leq_{\text{sW}} \text{wtu-D}_k^2$.

Proof. For part (1), it is enough to show that $\text{LPO} \leq_{\text{w}} \text{wtu-D}_2^2$, the rest of the reductions being obvious. This is proved much like Proposition 2.11. Given an

instance S of LPO, define $c : [\omega]^2 \rightarrow 2$ by

$$c(x, y) = \begin{cases} 1 & \text{if } S(x) = S(y), \\ 0 & \text{otherwise.} \end{cases}$$

If $S = 0^\omega$ then $c(x, y) = 1$ for all $x < y$, and if $S = 0^n 1^\omega$ then $\lim_y c(x, y) = 1$ for all $x > n$ and $\lim_y c(x, y) = 0$ for all $x \leq n$. Either way, c is an instance of wtu-D_2^2 with witness of thin-unbalancing 0. Now if L is any limit-homogeneous set for c then $S = 0^\omega$ if and only if there is an $x \leq \min L$ such that $S(x) = 1$.

For part (2), we prove the result for SRT_k^2 , the proof for D_k^2 being similar. Let $c : [\omega]^2 \rightarrow k$ and $d : [\omega]^2 \rightarrow 2$ be instances of wtu-SRT_k^2 and wtu-SRT_2^2 , respectively. Say the witnesses of thin-unbalancing for c and d are $i_c < k$ and $i_d < 2$, respectively. Define $e : [\omega]^2 \rightarrow k$ by

$$e(x, y) = \begin{cases} i_c & \text{if } c(x, y) = i_c \text{ or } d(x, y) = i_d, \\ c(x, y) & \text{otherwise.} \end{cases}$$

Notice that e is stable. We claim that e is thin-unbalanced as witnessed by i_c . Indeed, if H were infinite and homogeneous for e with color i_c then we could define $f : [H]^2 \rightarrow 2$ by

$$f(x, y) = \begin{cases} 0 & \text{if } c(x, y) = i_c, \\ 1 & \text{otherwise.} \end{cases}$$

Any infinite homogeneous set for f contained in H with color 0 would be homogeneous for c with color i_c , and any infinite homogeneous set for f contained in H with color 1 would be homogeneous for d with color i_d . Neither of these is possible by assumption, so the claim holds. Hence, e is an instance of wtu-SRT_k^2 , and it is clear that any infinite homogeneous set for e is homogeneous for both c and d . \square

As we will see in Proposition 4.3, $\text{wtu-P} \equiv_{\text{w}} \Delta_2^0\text{-wtu-P}$ for $\text{P} \in \{\text{SRT}_2^2, \text{D}_2^2\}$. Note that in part (1) above, the reduction from LPO to wtu-P cannot be improved from \leq_{w} to \leq_{sw} . Indeed, it follows from a result of Brattka and Rakotoniaina [10, Corollary 3.15] that $\text{LPO} \not\leq_{\text{sw}} \text{RT}_k^n$ for all $n, k \geq 1$.

Theorem 3.7. *Let P be a problem. Then $\text{P} \leq_{\text{w}} \Delta_2^0\text{-wtu-D}_k^2$ if and only if $\text{LPO} \times \text{P} \leq_{\text{w}} \text{D}_k^2$.*

Proof. For the forward direction, we prove that $\text{LPO} \times \Delta_2^0\text{-wtu-D}_k^2 \leq_{\text{w}} \text{D}_k^2$. Again, we emulate the proof of Proposition 2.11. Let $S \in 2^\omega$ be an instance of LPO. Let $\langle c, \ell \rangle$ be an instance of $\Delta_2^0\text{-wtu-D}_k^2$, so that c is a stable coloring $[\omega]^2 \rightarrow k$, and $\ell : \omega \rightarrow k$ is a function with $\lim_y \ell(y) = i < k$ a witness of thin-unbalancing for c . Define $d : [\omega]^2 \rightarrow k$ by

$$d(x, y) = \begin{cases} c(x, y) & \text{if } S(x) = S(y), \\ \ell(y) & \text{otherwise.} \end{cases}$$

Thus, if $S = 0^\omega$ then $d = c$, and if $S = 0^n 1^\omega$ for some n then $c(x, y) = d(x, y)$ for all $x \geq n$ and $\lim_y d(x, y) = \lim_y \ell(y) = i$ for all $x < n$. Since c has no infinite limit-homogeneous set with color i , it follows that every infinite limit-homogeneous set L for d is also limit-homogeneous for c . Moreover, we have that $S = 0^n 1^\omega$ if and only if $(\exists x \leq \min L)[S(x) = 1]$. Hence, $\langle \{b\}, L \rangle$, where b is 0 or 1 depending on whether $S = 0^\omega$ or $S = 0^n 1^\omega$ for some $n < L$, is a uniformly $S \oplus L$ -computable solution to the $\text{LPO} \times \Delta_2^0\text{-wtu-D}_k^2$ -instance $\langle S, \langle c, \ell \rangle \rangle$.

For the reverse direction, fix a problem P such that $LPO \times P \leq_w D_k^2$, say via functionals Φ and Ψ . Fix an instance X of P . We describe a uniform procedure to define an X -computable thin-unbalanced stable coloring $d : [\omega]^2 \rightarrow k$ with a witness given in a Δ_2^0 way, and a uniform procedure for turning any infinite limit-homogeneous set for d into a P -solution for X . To begin, let $c = \Phi^{0^\omega \oplus X}$, which is a stable coloring $[\omega]^2 \rightarrow k$ by assumption. Define \mathcal{C} to be the $\Pi_1^{0,X}$ class consisting of all partitions $\langle P_0, \dots, P_{k-1} \rangle$ of ω such that

$$(\forall i < k)(\forall \text{ finite } F \subseteq P_i)[\Psi^{0^\omega \oplus X \oplus F}(0) \simeq 0].$$

(As in the proof of Theorem 2.6, we think of $\Psi^{0^\omega \oplus X \oplus F}$ as the characteristic function of a join.)

It must be that $\mathcal{C} = \emptyset$. For suppose otherwise, and choose any $\langle P_0, \dots, P_{k-1} \rangle \in \mathcal{C}$ and an $i < k$ such that P_i is infinite. Let $L \subseteq P_i$ be an infinite limit-homogeneous set for c . Then by the definition of \mathcal{C} , we have $\Psi^{0^\omega \oplus X \oplus L}(0) \simeq 0$, which is a contradiction because $\langle 0^\omega, X \rangle$ is an instance of $LPO \times P$, and we should thus have $\Psi^{0^\omega \oplus X \oplus L}(0) \downarrow = 1$. So \mathcal{C} is empty, as claimed. By compactness, we can uniformly X -computably find an m such that for every partition $\langle P_0, \dots, P_{k-1} \rangle$ of ω , there are an $i < k$ and a finite $F \subseteq P_i \upharpoonright m$ such that $\Psi^{0^\omega \oplus X \oplus F}(0) \downarrow = 1$. Let $u \geq m$ be a bound on the uses of all these computations, for all possible such F . Choose $n > u$ large enough so that $\Phi^{0^n 1^\omega \oplus X}$ and $c = \Phi^{0^\omega \oplus X}$ agree below u . Let $S = 0^n 1^\omega$, and let $d = \Phi^{S \oplus X}$. Note that d is uniformly X -computable.

We claim that d is thin-unbalanced. To see this, let $\langle P_0, \dots, P_{k-1} \rangle$ be the partition of ω given by $P_i = \{x \in \omega : \lim_y d(x, y) = i\}$. Let $i < k$ and $F \subseteq P_i \upharpoonright m$ be as above. If P_i were infinite, then there would be an infinite limit-homogeneous set L for d having F as an initial segment, and by construction, this set would satisfy $\Psi^{S \oplus X \oplus L}(0) \downarrow = 1$ even though $S \neq 0^\omega$. Thus, P_i is finite, so i is a witness to thin-unbalancing for d . Moreover, since i depends only on $\lim_y d(x, y)$ for $x < m$, it follows that i can be approximated from d , and hence from X , in a uniform Δ_2^0 way. So d is an instance of $\Delta_2^0\text{-wtu-D}_k^2$. Now if L is any infinite limit-homogeneous set for d , we must have $\Psi^{S \oplus X \oplus L} = \langle \{1\}, Y \rangle$, where Y is a P -solution to X . Hence, there is a uniform way to convert $X \oplus L$ into a P -solution for X , as desired. \square

A succinct way to express the characterization given by the preceding theorem is that $\Delta_2^0\text{-wtu-D}_k^2 \equiv_w \sup_{\leq_w} \{P : P \times LPO \leq_w D_k^2\}$. We can obtain several other results of this sort, the proofs of which are similar to the preceding theorem.

Proposition 3.8. *The following all exist and are all Weihrauch equivalent:*

- (1) $\Delta_2^0\text{-wtu-D}_k^2$;
- (2) $\sup_{\leq_w} \{P : P \times LPO \leq_w D_k^2\}$;
- (3) $\sup_{\leq_w} \{P : P \times LPO \leq_w \Delta_2^0\text{-wtu-D}_k^2\}$;
- (4) $\sup_{\leq_w} \{P : P \times \Delta_2^0\text{-wtu-D}_k^2 \leq_w \Delta_2^0\text{-wtu-D}_k^2\}$;
- (5) $\sup_{\leq_w} \{P : P \times \Delta_2^0\text{-wtu-D}_k^2 \leq_w D_k^2\}$.

We do not know a similar characterization for SRT_k^2 , nor even an answer to the following question. (It is worth noting that the Weihrauch lattice is not complete. Indeed, by [19, Proposition 3.15], it does not have any nontrivial infinite suprema.)

Question 3.9. Does $\sup_{\leq_w} \{P : P \times LPO \leq_w \text{SRT}_k^2\}$ exist?

As a partial step, we have the following:

Proposition 3.10. *Let P be a problem.*

- (1) *If $P \leq_w \Delta_2^0\text{-wtu-SRT}_k^2$ then $\text{LPO} \times P \leq_w \text{SRT}_k^2$.*
- (2) *If $\text{LPO} \times P \leq_w \text{SRT}_k^2$ then P is Weihrauch reducible to the problem whose instances are pairs $\langle c, \ell \rangle$ where c is an instance of SRT_k^2 and $\ell : \omega \rightarrow k$ is a function such that $\lim_y \ell(y)$ exists and is a witness to thin-unbalancing for c on some set that is low relative to c , and the solutions to such a pair are the SRT_k^2 -solutions to c .*

Proof. Part (1) is proved just like the forward direction of Theorem 3.7. For part (2), we proceed as in the proof of the reverse direction of Theorem 3.7, only the $\Pi_1^{0,X}$ class \mathcal{C} now consists of all partitions $\langle P_0, \dots, P_{k-1} \rangle$ of ω such that

$$(\forall i < k)(\forall \text{ finite } F \subseteq P_i)[(\forall x, y \in F)[c(x, y) = i] \rightarrow \Psi^{0^\omega \oplus X \oplus F}(0) \simeq 0].$$

We can assume that $c \equiv_{\text{T}} X$, because we can replace it by the coloring c' obtained by letting $c'(n, n+1) = X(n)$ and $c'(x, y) = c(x, y)$ for all other pairs. An infinite solution to c' can be uniformly transformed into one to c by thinning.

Now, if $\mathcal{C} = \emptyset$, let n be as in the corresponding case in the proof of Theorem 3.7. For each $i < k$, let $P_i = \{x : \lim_y \Phi^{0^n 1^\omega \oplus X}(x, y) = i\}$. Then for some $i < k$, there exists a finite $F \subseteq P_i$ such that F is homogeneous for $\Phi^{0^n 1^\omega \oplus X}$ with color i and $\Psi^{0^n 1^\omega \oplus X \oplus F}(0) \downarrow = 1$. Moreover, this F and i can be approximated in a $\Delta_2^{0,X}$ way (i.e., Δ_2^0 relative to the instance $\langle 0^n 1^\omega, X \rangle$). Now if $\Phi^{0^n 1^\omega \oplus X}$ had any infinite homogeneous set with color i , then c would have such a set extending F , which would produce the same contradiction as in Theorem 3.7. Thus, it must be that $\Phi^{0^n 1^\omega \oplus X}$ has no homogeneous set with color i , so in particular, it is thin-unbalanced (on the low set ω).

If, on the other hand, $\mathcal{C} \neq \emptyset$, then let $\langle P_0, \dots, P_{k-1} \rangle$ be the canonical low-over- X element of it (given by the proof of the low basis theorem). Clearly, we can approximate in a $\Delta_2^{0,X}$ way (in fact, in a $\Pi_1^{0,X}$ way), the least i such that P_i is infinite. Then $c = \Phi^{0^\omega \oplus X}$ must be thin-unbalanced on P_i with witness i . Otherwise, we could take a homogeneous set H for c with color i contained in P_i and have, by the definition of \mathcal{C} , that $\Psi^{0^\omega \oplus X \oplus H}(0) \simeq 0$, which is a contradiction because $\langle 0^\omega, X \rangle$ is an instance of $\text{LPO} \times P$, and we should thus have $\Psi^{0^\omega \oplus X \oplus H}(0) \downarrow = 1$. \square

Remark 3.11. With a view to some of the recent work on the algebraic structure of the Weihrauch degrees ([9, 16]), Theorem 3.7 suggests a natural *parallel quotient* operator on problems, given by $P/Q = \sup_{\leq_w} \{R : R \times Q \leq_w P\}$. We have no reason to think this operator is total, but studying the kinds of problems for which it is defined ought to be interesting in its own right.

4. THE COFINITE-TO-INFINITE PRINCIPLE

In this section, we briefly depart from studying products, to investigate wtu-D_2^2 (in the guise of a Weihrauch-equivalent principle introduced below) in the context of other weak Weihrauch degrees. Some of our terminology will be specific to the Weihrauch literature, and we refer the reader to [4] for any definitions we omit.

We begin by introducing the following ‘‘cofinite set to infinite set’’ principle.

Definition 4.1. $\text{CFI}_{\Delta_2^0}$ is the restriction of D_2^2 to colorings $c : [\omega]^2 \rightarrow 2$ such that $\lim_y c(x, y) = 1$ for almost all x .

Thus, informally, $\text{CFI}_{\Delta_2^0}$ is the problem of finding an infinite subset of a cofinite set given by a Δ_2^0 approximation. This leads to the following initial observation:

Proposition 4.2. $\text{CFI}_{\Delta_2^0} \leq_w \text{lim}$.

Proof. Fix an instance $c : [\omega]^2 \rightarrow 2$ of $\text{CFI}_{\Delta_2^0}$. For each x , we have that $\lim_y c(x, y)$ exists, and hence the sequence $(c_y)_{y \in \omega}$, where for each y the function $c_y : \omega \rightarrow \omega$ is given by $c_y(x) = c(x, y)$, is an instance of lim . Apply lim to find a solution to $(c_y)_{y \in \omega}$, i.e., the function $d : \omega \rightarrow \omega$ defined by $d(x) = \lim_y c(x, y)$ for all x . By assumption, the d -computable set $\{x \in \omega : d(x) = 1\}$ is infinite, and so is a $\text{CFI}_{\Delta_2^0}$ -solution to c . \square

The connection to the previous section is provided by the following result.

Proposition 4.3. $\text{CFI}_{\Delta_2^0} \equiv_w \text{wtu-D}_2^2 \equiv_w \Delta_2^0\text{-wtu-D}_2^2$.

Proof. It is clear that $\text{CFI}_{\Delta_2^0} \leq_w \text{wtu-D}_2^2 \leq_w \Delta_2^0\text{-wtu-D}_2^2$. In the other direction, suppose we are given an instance $\langle c, \ell \rangle$ of $\Delta_2^0\text{-wtu-D}_2^2$. Define $d : [\omega]^2 \rightarrow 2$ by

$$d(x, y) = \begin{cases} 1 & \text{if } c(x, y) = 1 - \ell(y), \\ 0 & \text{otherwise.} \end{cases}$$

Now for all x , we have that $\lim_y c(x, y) = 1 - \lim_y \ell(y)$ if and only if $\lim_y d(x, y) = 1$. In particular, since $\lim_y c(x, y) = 1 - \lim_y \ell(y)$ for almost all x , we have $\lim_y d(x, y) = 1$ for almost all x . Clearly, every limit-homogeneous set for d is also limit-homogeneous for c . \square

Notice that a similar proof shows that $\text{wtu-SRT}_2^2 \equiv_w \Delta_2^0\text{-wtu-SRT}_2^2$.

We now compare $\text{CFI}_{\Delta_2^0}$ with the choice principle $\mathbf{C}_{\mathbb{N}}$ defined in Section 1.

Proposition 4.4. $\text{CFI}_{\Delta_2^0} \star \mathbf{C}_{\mathbb{N}} \equiv_w \text{CFI}_{\Delta_2^0}$.

Proof. First we show that $\text{CFI}_{\Delta_2^0} \star \mathbf{C}_{\mathbb{N}} \leq_w \text{CFI}_{\Delta_2^0} \times \mathbf{C}_{\mathbb{N}}$. Note that $\mathbf{C}_{\mathbb{N}}$ is computable with finitely many mindchanges, and these mindchanges can be incorporated into the Δ_2^0 instances of $\text{CFI}_{\Delta_2^0}$. Thus, we can compute directly the impact $\mathbf{C}_{\mathbb{N}}$ has on the $\text{CFI}_{\Delta_2^0}$ -instance, and do not need to use them sequentially.

Then we argue that $\text{CFI}_{\Delta_2^0} \times \mathbf{C}_{\mathbb{N}} \leq_w \text{CFI}_{\Delta_2^0}$. We identify an instance e of $\mathbf{C}_{\mathbb{N}}$ with the complement of the set enumerated by e , and an instance c of $\text{CFI}_{\Delta_2^0}$ with the corresponding $\Delta_2^{0,c}$ set. As shown in [33, Lemma 2.3], we may assume without loss of generality that the instances of $\mathbf{C}_{\mathbb{N}}$ are of the form $\{n \mid n > k\}$ for some $k \in \omega$. Given instances of $\mathbf{C}_{\mathbb{N}}$ and of $\text{CFI}_{\Delta_2^0}$, we can compute the intersection of these instances, and think of it as an instance of $\text{CFI}_{\Delta_2^0}$. Any infinite subset of this intersection is a solution to the original $\text{CFI}_{\Delta_2^0}$ instance, and any element a solution to the $\mathbf{C}_{\mathbb{N}}$ instance. \square

We can think of the instances of $\text{CFI}_{\Delta_2^0}$ as being functions $p : \omega \rightarrow \omega$ such that $|\{i \in \omega : p(i) = n + 1\}| < \infty$ for all n , and such that $|\{i \in \omega : p(i) = n + 1\}|$ is even for cofinitely many n . Then, a solution is any infinite set Y such that if $n \in Y$ then $|\{i \in \omega : p(i) = n + 1\}|$ is even. It is easy to see that this formulation is Weihrauch equivalent to the one given in Definition 4.1. However, we shall find this version more convenient for our results below.

Given $p \in \omega^\omega$ as above, let $\psi(p) = \{n : |\{i \in \omega : p(i) = n + 1\}| \text{ is even}\}$. For each $p \in \omega^{<\omega} \cup \omega^\omega$, let $[p] = \{n : \exists i p(i) = n + 1\}$. For $\sigma \in \omega^{<\omega}$, let $\hat{\sigma}$ be the

length-lexicographically least extension of σ such that $|\{i \in \omega : \widehat{\sigma}(i) = n + 1\}|$ is even for all $n \in [\sigma]$.

Definition 4.5. For $\sigma \in \omega^{<\omega}$, let $\text{CFI}_{\Delta_2^0}^\sigma$ denote the restriction of $\text{CFI}_{\Delta_2^0}$ to instances $\widehat{\sigma}p$ such that $[\sigma] \cap [p] = \emptyset$.

Proposition 4.6. For every $\sigma \in \omega^{<\omega}$, we have $\text{CFI}_{\Delta_2^0}^\sigma \equiv_{\text{w}} \text{CFI}_{\Delta_2^0}$.

Proof. Let $h : \omega \rightarrow \{x \in \omega : x > \max[\sigma]\}$ be a computable bijection. Let $\bar{h} : \omega^\omega \rightarrow \omega^\omega$ be defined pointwise via $\bar{h}(p)(n) = h(p(n))$. Then $\text{CFI}_{\Delta_2^0}(p) = h^{-1} \circ \text{CFI}_{\Delta_2^0}^\sigma(\widehat{\sigma}\bar{h}(p))$. \square

We can now prove that $\text{CFI}_{\Delta_2^0}$ has properties very similar to being a *total fractal* (see Brattka, Gherardi, and Pauly [4, Section 4]; see also Theorem 7.15 in that paper). In the context of Weihrauch degrees, a fractal may be thought of as a problem that retains its full power on arbitrarily small (clopen) restrictions of its domain. Since $\text{CFI}_{\Delta_2^0}$ is defined on Δ_2^0 -approximations, it is clear that it is a fractal.

Proposition 4.7. Let P be any problem. If $\text{CFI}_{\Delta_2^0} \leq_{\text{w}} P \star \text{C}_{\mathbb{N}}$, then $\text{CFI}_{\Delta_2^0} \leq_{\text{w}} P$.

Proof. Let Φ map instances of $\text{CFI}_{\Delta_2^0}$ to instances of $\text{C}_{\mathbb{N}}$. If there is no string $\sigma \in \omega^{<\omega}$ such that $\Phi^{\widehat{\sigma}}(0, s) \downarrow = 1$ for some s , then 0 is always a valid answer to the $\text{C}_{\mathbb{N}}$ -instances used in the reduction, and the $\text{C}_{\mathbb{N}}$ -call is useless. So suppose otherwise, and let σ_0 be such a string.

Assume now that we have defined $\sigma_0, \dots, \sigma_{k-1} \in \omega^{<\omega}$, and choose the least $n \notin \bigcup_{i < k} [\sigma_i]$. Suppose there is no string σ with $n \in [\sigma]$ and $[\sigma] \cap [\sigma_i] = \emptyset$ for all $i < k$, and such that $\Phi^{\widehat{\sigma_0 \dots \widehat{\sigma}_{k-1} \widehat{\sigma}}}(k, s) \downarrow = 1$ for some s . Then choose any σ with $n \in [\sigma]$ and $[\sigma] \cap [\sigma_i] = \emptyset$ for all $i < k$. Now $\text{CFI}_{\Delta_2^0}^{\sigma_0 \dots \sigma_{k-1} \sigma} \leq_{\text{w}} P$ because we can replace the output of $\text{C}_{\mathbb{N}}$ by k . By Proposition 4.6, this fact implies the claim. So suppose otherwise, and let σ_k be a string with the desired properties.

If this procedure never stops, then we construct some $p = \widehat{\sigma_0} \widehat{\sigma_1} \widehat{\sigma_2} \dots$. By induction, all the $[\sigma_i]$ are mutually disjoint, and every $n \in \omega$ appears some even number of times in some $[\sigma_i]$, so certainly p is an instance of $\text{CFI}_{\Delta_2^0}$. However, by construction we also find that for each k there is an s such that $\Phi^p(k, s) = 1$, so Φ^p is not an instance of $\text{C}_{\mathbb{N}}$, which is a contradiction. \square

This proposition allows us to deduce a number of non-reduction facts about $\text{CFI}_{\Delta_2^0}$, which point to its strength. We begin with the following. Neumann and Pauly [29] introduced the *sorting principle*, Sort , whose instances are all elements of 2^ω , such that the instance $p \in 2^\omega$ has the unique solution $0^n 1^\omega$ if p contains exactly n many 0's, and 0^ω if p contains infinitely many 0's. We refer to [4, Definition 1.2] for the definitions of the k -fold product and the star operation, $*$.

Theorem 4.8. $\text{CFI}_{\Delta_2^0} \not\leq_{\text{w}} \text{Sort}^*$.

Proof. Assume that $\text{CFI}_{\Delta_2^0} \leq_{\text{w}} \text{Sort}^*$. Then there is a Turing functional mapping instances of $\text{CFI}_{\Delta_2^0}$ to instances of Sort^* , and hence there are a σ and a k such that $\text{CFI}_{\Delta_2^0}^\sigma \leq_{\text{w}} \text{Sort}^k$. Choose k minimal for which there is such a σ . By Proposition 4.6 we also have $\text{CFI}_{\Delta_2^0} \leq_{\text{w}} \text{Sort}^k$. Let Φ and Ψ witness the reduction.

Suppose there is a τ such that Φ^τ outputs at least n many 0's for each input to Sort and the output of Ψ on input $\langle \tau, \langle 0^n, \dots, 0^n \rangle \rangle$ contains some $l \in \mathbb{N}$. Then there

is some p such that $l \notin \psi(\tau p)$, which is a contradiction. Thus for each p , there is some $d \leq k$ such that the d th input to Sort given by Φ^p has finitely many 0's. The set of pairs $\langle d, n \rangle$ such that the d th input has some 0 in a position greater than n is c.e. in p , so from p we can obtain an instance of $\mathbf{C}_{\mathbb{N}}$ whose solutions are pairs $\langle d, n \rangle$ such that the d th input has no 0's at positions greater than n . It follows that $\text{CFI}_{\Delta_2^0} \leq_w \text{Sort}^{k-1} \star \mathbf{C}_{\mathbb{N}}$. By Proposition 4.7, this in turn implies $\text{CFI}_{\Delta_2^0} \leq_w \text{Sort}^{k-1}$, contradicting the minimality of k . \square

In the next proposition, $\mathbf{K}_{\mathbb{N}}$ denotes the choice problem for *compact* subsets of \mathbb{N} (see [4]). We refer the reader to [4, Section 6] for the definition of the jump operator, $'$, on Weihrauch degrees. Definitions of the countable coproduct \coprod and the problems $\mathbf{C}_{\{0, \dots, n\}}$ used in the proof below can also be found in that paper, in Sections 4 and 7, respectively.

Proposition 4.9. $\text{CFI}_{\Delta_2^0} \leq_w \mathbf{C}'_{\mathbb{N}}$ but $\text{CFI}_{\Delta_2^0} \not\leq_w \mathbf{K}'_{\mathbb{N}}$.

Proof. For the reduction, fix some enumeration $(\sigma_i)_{i \in \omega}$ of $\omega^{<\omega}$. Given some input d to $\text{CFI}_{\Delta_2^0}$ we define a sequence $(e_n)_{n \in \omega}$ with $e_n : \omega^2 \rightarrow 2$ by $e_n(x, s) = 0$ iff for all $y < n$ we have that $d(y, s) = 0$ iff y occurs in σ_x . The sequence $(e_n)_{n \in \omega}$ converges to some $e : \omega^2 \rightarrow 2$ with the property that $e(x, s) = 0$ for all $s \in \omega$ precisely when σ_x lists exactly those k with $\lim_y d(k, y) = 0$. Clearly, from such a finite tuple we can compute an infinite subset of its complement.

For the non-reduction, note that $\mathbf{K}'_{\mathbb{N}} \leq_w \left(\coprod_{n \in \omega} \mathbf{C}'_{\{0, \dots, n\}} \right) \star \mathbf{C}_{\mathbb{N}}$, so if $\text{CFI}_{\Delta_2^0} \leq_w \mathbf{K}'_{\mathbb{N}}$, then by Proposition 4.7, we have $\text{CFI}_{\Delta_2^0} \leq_w \left(\coprod_{n \in \omega} \mathbf{C}'_{\{0, \dots, n\}} \right)$. As $\text{CFI}_{\Delta_2^0}$ is a fractal (as discussed above), then there is some $k \in \omega$ with $\text{CFI}_{\Delta_2^0} \leq_w \mathbf{C}'_{\{0, \dots, k\}}$. But this is impossible for reasons of cardinality. \square

The *connected choice problem* of the next theorem was introduced by Brattka, Le Roux, Miller, and Pauly [8]. The instances of \mathbf{CC}_1 are nonempty closed subintervals of the real unit interval (see [8] for details on how the elements of the collection $\mathcal{A}([0, 1])$ of such subintervals are represented), and the solutions to any such instance are the points inside it.

Theorem 4.10. $\mathbf{CC}_1 \not\leq_w \text{CFI}_{\Delta_2^0}$.

Proof. Assume that $\mathbf{CC}_1 \leq_w \text{CFI}_{\Delta_2^0}$ via Φ and Ψ . Let $p_0 \in \omega^\omega$ be a name for $[0, 1] \in \mathcal{A}([0, 1])$. There have to be some finite set $B_0 \subset \omega$ and a prefix σ_0 of p_0 such that upon reading σ_0 and B_0 , the functional Ψ outputs a 2^{-2} -approximation of some $x_0 \in [0, 1]$. We can find some $\tau_0 p_1$ such that $\sigma_0 \tau_0 p_1$ is a name for some interval I_1 with $|I_1| \geq 2^{-2}$ and such that for any q extending $\sigma_0 \tau_0$ and representing some $A \in \mathcal{A}([0, 1])$, we have that $A \cap B(x_0, 2^{-2}) = \emptyset$ (where $B(x_0, 2^{-2})$ is the ball of radius 2^{-2} around x_0). It follows that for any q extending $\sigma_0 \tau_0$, the set $\psi(\Phi^q)$ must not contain B_0 , for if it did, there would be a solution to Φ^q containing B_0 that would trick Ψ into outputting a 2^{-2} -approximation of x_0 , which cannot be correct.

In the next step, Ψ has to output some 2^{-4} -approximation of some x_1 upon reading some prefix $\sigma_0 \tau_0 \sigma_1$ of $\sigma_0 \tau_0 p_1$ and a finite set B_1 with $\max B_1 > \min B_0$. We pick τ_1 to exclude $B(x_1, 2^{-4})$ from the solution set, and thus conclude that for any q extending $\sigma_0 \tau_0 \sigma_1 \tau_1$, the set $\psi(\Phi^q)$ must not contain B_1 (nor B_0).

By iterating the procedure, we obtain some input $\sigma_0\tau_0\sigma_1\tau_1\sigma_2\tau_2\cdots \in \omega^\omega$, which is in the domain of CC_1 (as this has a total domain if represented in a suitable way), but such that $\psi(\Phi^{\sigma_0\tau_0\sigma_1\tau_1\sigma_2\tau_2\cdots})$ excludes countably many disjoint finite sets B_0, B_1, \dots . Hence, $\psi(\Phi^{\sigma_0\tau_0\sigma_1\tau_1\sigma_2\tau_2\cdots}) \notin \text{dom}(\text{CFI}_{\Delta_2^0})$, and we have derived a contradiction. \square

Brattka, Hölzl, and Kuiper [6, Proposition 16] showed that $\text{CC}_1 \leq_w \text{Sort}$, so it follows that $\text{Sort} \not\leq_w \text{CFI}_{\Delta_2^0}$. An alternate proof of this fact can be given by using the following technical notion.

Definition 4.11. Suppose $G : \subseteq \mathbf{X} \rightrightarrows \mathbf{Z}$ is a partial multifunction of represented spaces. Then G is *low for functions* if, for every $f : \mathbf{Y} \rightarrow \omega^\omega$ that satisfies $f \leq_w \lim \star G$, we have $f \leq_w \lim$.

Proposition 4.12. Let $G : \mathbf{X} \rightrightarrows \mathcal{O}(\omega)$ (where $\mathcal{O}(\omega)$ consists of the subsets of ω represented by enumerations of their elements) be such that

$$(\forall x \in \mathbf{X})(\exists k_0 \in \omega)(\forall k \geq k_0)[\{n : n \geq k\} \in G(x)].$$

Then G is low for functions.

Proof. Let $f : \mathbf{Y} \rightarrow \omega^\omega$ be such that $f \leq_w \lim \star G$. Without loss of generality, we may assume that $\mathbf{X}, \mathbf{Y} \subseteq \omega^\omega$. As \lim is transparent (see Brattka, Gherardi, and Marcone [3, Fact 5.5]), we can obtain $f(x) = \lim_{i \rightarrow \infty} \Psi_i(x, G(\Phi(x)))$ for some functionals Φ and Ψ_i . Let $\omega_{\geq k} = \{n \in \omega : n \geq k\} \in \mathcal{O}(\omega)$. Now for any $x \in \mathbf{X}$, we have that $\Psi_k(x, \omega_{\geq k})$ is defined and is an element of $\Psi_k(x, G(\Phi(x)))$ for almost all k . As f is a function, in $\lim_{i \rightarrow \infty} \Psi_i(x, G(\Phi(x)))$ it does not matter whether we choose from $G(\Phi(x))$ once for the entire expression, or separately for each i . Thus, we can compute $f(x)$ as $\lim_{k \rightarrow \infty} \Psi_k(x, \omega_{\geq k})$. (While finitely many of these values may be undefined, this problem can be resolved with a standard argument.) \square

Lemma 4.13. Let G be low for functions, and $f : \mathbf{X} \rightarrow \omega^\omega$ with $f \leq_w G$. Then $\lim \star f \leq_w \lim$.

Proof. As f is a function, so is $\lim \star f$. Moreover, $f \leq_w G$ implies $\lim \star f \leq_w \lim \star G$, so $\lim \star f \leq_w \lim$. \square

Corollary 4.14. $\text{Sort} \not\leq_w \text{CFI}_{\Delta_2^0}$.

Proof. By Proposition 4.12, $\text{CFI}_{\Delta_2^0}$ is low for functions. As Sort is a function, Lemma 4.13 shows that if we had $\text{Sort} \leq_w \text{CFI}_{\Delta_2^0}$ we would also have $\lim \star \text{Sort} \leq_w \lim$. However, it is not difficult to check that $\text{LPO}' \leq_w \text{LPO} \star \text{Sort} \leq_w \lim \star \text{Sort}$, but $\text{LPO}' \not\leq_w \lim$. (That $\lim \star \text{Sort} \not\leq_w \lim$ also follows from [6, Proposition 21].) To see that this non-reduction holds, first note that there is a uniformly computable sequence S_0, S_1, \dots of instances of LPO' such that for each e , the e th Turing functional Φ_e is total if and only if $S_e = 0^\omega$. Thus, for each e , to determine whether S_e has solution 0 is Π_2^0 -hard. On the other hand, every computable \lim -instance has a uniformly Δ_2^0 solution. \square

Note that while the proof above shows that $\lim \star \text{Sort} \not\leq_w \lim$, it was shown by Neumann and Pauly [29, Corollary 32] that $\lim \star \lim \star \text{Sort} \leq_w \lim \star \lim$.

We conclude with one final reduction. Recall that WWKL is the problem whose instances are closed subsets of 2^ω of positive measure, with solutions being the members of the given set. We refer the reader to Downey and Hirschfeldt [14, Chapter 6] for background on Martin-Löf randomness.




FIGURE 2. Location of $\text{CFI}_{\Delta_2^0}$ in the Weihrauch lattice. An arrow from P to Q represents the reduction $Q \leq_W P$. No additional arrows can be added other than those that follow by transitivity. For (non-)reductions not explicitly mentioned above, see [6, Sections 4 and 5 and Figure 5].

Proposition 4.15. $\text{CFI}_{\Delta_2^0} \leq_W \text{WWKL}'$.

Proof. Let \mathcal{C}_0 be a fixed $\Pi_1^{0,\emptyset'}$ class all of whose members are 2-random. Given an instance $c : [\omega]^2 \rightarrow 2$ of $\text{CFI}_{\Delta_2^0}$, let \mathcal{C} consist of all $X \in \mathcal{C}_0$ such that for all i , if $X(i) = 1$ then $\lim_s c(i, s) = 1$. Then \mathcal{C} is a $\Pi_1^{0,c'}$ subclass of \mathcal{C}_0 , and it still has positive measure since, e.g., if X is any 2-random real and n is least such that $c(i, s) = 1$ for all $i \geq n$, then \mathcal{C} contains the 2-random real $0^{n-1}X(n)X(n+1)\cdots$. Thus, \mathcal{C} may be regarded as an instance of WWKL' , and if X is any element of \mathcal{C} then $\{i : X(i) = 1\}$ is infinite and is therefore a $\text{CFI}_{\Delta_2^0}$ -solution to c . \square

We summarize the results of this section in Figure 2.

5. RAMSEY'S THEOREM FOR SINGLETONS

In this section, we investigate Ramsey's theorem for singletons and different numbers of colors, and how these problems behave under Weihrauch reducibility with respect to products. A motivating toy example is the fact that $\text{RT}_2^1 \times \text{RT}_2^1 \leq_W \text{RT}_4^1$, and in fact, it is easy to see that for all $n \geq 1$ and $k_0, \dots, k_n \geq 2$,

$$\prod_{m=0}^n \text{RT}_{k_m}^1 \leq_{\text{sW}} \text{RT}_{\prod_{m=0}^n k_m}^1.$$

We show below that the right-hand side is optimal. Our results extend a number of similar investigations, including by Dorais, Dzhafarov, Hirst, Miletì, and Shafer [13], Hirschfeldt and Jockusch [22], Patey [32], and Brattka and Rakotoniaina [10].

In the sequel, we will regard RT_k^1 as the problem whose instances are colorings $c : \omega \rightarrow k$ and whose solutions are colors that appear infinitely often in c . Note that this formulation of RT_k^1 is Weihrauch equivalent to the more usual one given in Definition 1.6, so we will not distinguish these versions when discussing Weihrauch

reducibility. In the context of strong Weihrauch reducibility, we will refer to the new version as rt_k^1 . The principle rt_k^1 can be understood as the Bolzano-Weierstrass theorem for the discrete space k , and was indeed studied as BWT_k by Brattka, Gherardi and Marcone [3]. A central result there is that $\text{rt}_k^1 \equiv_{\text{sW}} C'_k$, which tells us that we could alternatively strive to understand the principles rt_k^1 by studying the finite choice principles C_k , and transferring the results using the jump of strong Weihrauch degrees.²

Given this formulation, the backward functionals of our strong Weihrauch reductions will have single numbers or tuples of numbers as oracles, and hence can be regarded as partial functions. For such a functional Ψ , we write $\Psi(n)$ instead of Ψ^n .

We begin with the following lemma:

Lemma 5.1. *Suppose that $P \leq_{\text{W}} Q$ and these problems satisfy the following properties:*

- *P has finite tolerance, i.e., there is some Θ such that if C_0 and C_1 are P-instances, $C_0(x) = C_1(x)$ for all x above some m , and S_0 is a P-solution to C_0 , then $\Theta^{S_0 \oplus m}$ is a P-solution to C_1 ;*
- *any finite modification of a P-instance is still a P-instance;*
- *solutions to all instances of P and Q lie in some fixed finite set.*

Then $P \leq_{\text{sW}} Q$.

Proof. Fix functionals Φ and Ψ witnessing that $P \leq_{\text{W}} Q$. Since solutions to all instances of P lie in some fixed finite set, we may assume that for each P-instance C and each s that is a Q-solution to Φ^C , we have that $\Psi^{C \oplus s}$ outputs a number that codes a P-solution to C . Fix a functional Θ witnessing that P has finite tolerance. Fix a finite solution set S for Q. We define functionals that witness that $P \leq_{\text{sW}} Q$.

First, we construct a τ that is a finite initial segment of some P-instance, such that τ decides (in the sense of Cohen 1-genericity) for each $s \in S$ whether $\Psi^{C \oplus s}$ converges for P-instances C extending τ . Since S is finite, such a τ exists.

We define $\widehat{\Phi}$ by $\widehat{\Phi}^C = \Phi^{C'}$, where C' is obtained from C by replacing its initial segment of length $|\tau|$ by τ itself. By our assumption on P, this C' is still a P-instance.

We define $\widehat{\Psi}$ by $\widehat{\Psi}(s) = \Theta^{\Psi^{\tau \oplus s} \oplus |\tau|}$. We show that $\widehat{\Phi}$ and $\widehat{\Psi}$ witness that $P \leq_{\text{sW}} Q$.

Take any P-instance C . Since C' is a P-instance, $\widehat{\Phi}^C = \Phi^{C'}$ is a Q-instance. Let s be any Q-solution to $\widehat{\Phi}^C$. Then $\Psi^{C' \oplus s}$ is a P-solution to C' . In particular, $\Psi^{C' \oplus s}$ converges. Since C' extends τ , by our construction of τ , we have that $\Psi^{\tau \oplus s} \downarrow = \Psi^{C' \oplus s} \downarrow$. Hence $\Psi^{\tau \oplus s}$ is a P-solution to C' . We conclude that $\widehat{\Psi}(s) = \Theta^{\Psi^{\tau \oplus s} \oplus |\tau|}$ is a P-solution to C . \square

It is easy to see that rt_k^1 (and finite parallel products of rt_k^1) satisfy the properties of P and Q in Lemma 5.1. Therefore we have the following.

Corollary 5.2. *If $\prod_{m=0}^n \text{RT}_{k_m}^1 \leq_{\text{W}} \text{RT}_N^1$, then $\prod_{m=0}^n \text{rt}_{k_m}^1 \leq_{\text{sW}} \text{rt}_N^1$.*

Optimality then follows from a counting argument:

Proposition 5.3. *If $\prod_{m=0}^n \text{rt}_{k_m}^1 \leq_{\text{sW}} \text{rt}_N^1$, then $N \geq \prod_{m=0}^n k_m$.*

²This approach seems very promising, but is left to future work.

Proof. Fix Φ and Ψ witnessing that $\prod_{m=0}^n \text{rt}_{k_m}^1 \leq_{\text{sw}} \text{rt}_N^1$. We show that for each $(a_0, \dots, a_n) \in \prod_{m=0}^n k_m$, there is some $i < N$ such that $\Psi(i) = (a_0, \dots, a_n)$.

Consider the tuple of constant colorings $(a_0^\omega, \dots, a_n^\omega)$. This is a $\prod_{m=0}^n \text{rt}_{k_m}^1$ -instance, so $\Phi(a_0^\omega, \dots, a_n^\omega)$ is an rt_N^1 -instance with some solution i . Then $\Psi(i)$ must be a solution to $(a_0^\omega, \dots, a_n^\omega)$, so $\Psi(i) = (a_0, \dots, a_n)$. \square

Corollary 5.4. *If $\prod_{m=0}^n \text{RT}_{k_m}^1 \leq_{\text{w}} \text{RT}_N^1$, then $N \geq \prod_{m=0}^n k_m$.*

Therefore the right-hand side of $\prod_{m=0}^n \text{rt}_{k_m}^1 \leq_{\text{sw}} \text{rt}_{\prod_{m=0}^n k_m}^1$ is optimal, with regards to both \leq_{w} and \leq_{sw} . However, we will see that $\text{RT}_{\prod_{m=0}^n k_m}^1 \not\leq_{\text{w}} \prod_{m=0}^n \text{RT}_{k_m}^1$ for all $n \geq 1$ and $k_0, \dots, k_n \geq 2$ (Proposition 5.13). In the rest of this section, we attempt to find the smallest N such that

$$\text{RT}_N^1 \not\leq_{\text{w}} \prod_{m=0}^n \text{RT}_{k_m}^1.$$

We start by giving a lower bound for N .

Proposition 5.5. *For all $n \geq 1$ and $k_0, \dots, k_n \geq 2$,*

$$\text{rt}_{1+\sum_{m=0}^n (k_m-1)}^1 \leq_{\text{sw}} \prod_{m=0}^n \text{rt}_{k_m}^1.$$

Proof. Suppose we are given an instance c of $\text{rt}_{1+\sum_{m=0}^n (k_m-1)}^1$. For $0 \leq m \leq n$, we define colorings

$$d_m : \omega \rightarrow \left\{ \sum_{i=0}^{m-1} (k_i - 1), \dots, \sum_{i=0}^m (k_i - 1) \right\}$$

as follows. Note that for each m , d_m will be a k_m -coloring.

For each m and x , we define $d_m(x)$ as follows. First check which color among $0, \dots, \sum_{i=0}^m (k_i - 1)$ appears most often among $c(0), \dots, c(x)$. (Resolve ties by picking the smallest color.) If this color is among $0, \dots, \sum_{i=0}^{m-1} (k_i - 1)$, let $d_m(x) = \sum_{i=0}^{m-1} (k_i - 1)$. Otherwise, let $d_m(x)$ be this color.

Now, given (a_0, \dots, a_n) such that, for each m , the color a_m appears infinitely often in d_m , we want to compute a color that appears infinitely often in c . Start by considering a_n . If $a_n \neq \sum_{i=0}^{n-1} (k_i - 1)$, then for infinitely many x , the color a_n appears most often among $c(0), \dots, c(x)$. In particular, a_n appears infinitely often in c .

On the other hand, if $a_n = \sum_{i=0}^{n-1} (k_i - 1)$, then for infinitely many x , some color among $0, \dots, \sum_{i=0}^{n-1} (k_i - 1)$ appears most often among $c(0), \dots, c(x)$. By the pigeonhole principle, some color among $0, \dots, \sum_{i=0}^{n-1} (k_i - 1)$ appears infinitely often in c . We then proceed to consider a_{n-1} and repeat the above case division. Eventually we either reach some a_m that is not equal to $\sum_{i=0}^{m-1} (k_i - 1)$, in which case a_m appears infinitely often in c , or we reach $a_0 = 0$, in which case 0 appears infinitely often in c . \square

In order to obtain upper bounds for N , we begin by restricting the reductions that we need to diagonalize against. Firstly, by Lemma 5.1, we need only handle strong Weihrauch reductions:

Proposition 5.6. *If $\text{RT}_N^1 \leq_{\text{w}} \prod_{m=0}^n \text{RT}_{k_m}^1$, then $\text{rt}_N^1 \leq_{\text{sw}} \prod_{m=0}^n \text{rt}_{k_m}^1$.*

We can impose a further restriction:

Lemma 5.7. *Suppose $\text{rt}_N^1 \leq_{\text{sw}} \prod_{m=0}^n \text{rt}_{k_m}^1$ via some forward functionals Φ_m , $0 \leq m \leq n$, where Φ_m computes the m^{th} coloring in the $\prod_{m=0}^n \text{rt}_{k_m}^1$ -instance, and a backward functional Ψ . Then for any $i < N$, there exists (a_0, \dots, a_n) where each $a_m < k_m$ and $\Psi(a_0, \dots, a_n) = i$.*

Proof. Given $i < N$, consider the coloring c that is constantly i . Then the tuple $(\Phi_0^c, \dots, \Phi_n^c)$ is a $\prod_{m=0}^n \text{rt}_{k_m}^1$ -instance. Hence it has some solution (a_0, \dots, a_n) . The only solution to c is i , so $\Psi(a_0, \dots, a_n)$ must be i . \square

Combining the previous two facts, we obtain:

Corollary 5.8. *Suppose $\text{RT}_N^1 \leq_{\text{w}} \prod_{m=0}^n \text{RT}_{k_m}^1$. Then $\text{rt}_N^1 \leq_{\text{sw}} \prod_{m=0}^n \text{rt}_{k_m}^1$, as witnessed by some Φ_m , $0 \leq m \leq n$, and Ψ where $\Psi : \prod_{m=0}^n k_m \rightarrow N$ is a surjective partial function.*

Henceforth, we will always assume that our reductions of rt_N^1 to $\prod_{m=0}^n \text{rt}_{k_m}^1$ have the above special form. In order to diagonalize against such reductions, it will be convenient to have the following notion of covering a tuple of colors using a set of tuples of colors.

Definition 5.9. If $X \subseteq \prod_{m=0}^n k_m$ and $(i_0, \dots, i_n) \in \prod_{m=0}^n k_m$, we say that X covers (i_0, \dots, i_n) if for each $0 \leq m \leq n$, there is an $(a_0, \dots, a_n) \in X$ such that $a_m = i_m$.

Observe that if c is a $\prod_{m=0}^n \text{rt}_{k_m}^1$ -instance whose solution set contains X , and X covers (i_0, \dots, i_n) , then (i_0, \dots, i_n) is also a solution to c .

The following terminology will also be useful.

Definition 5.10. For a surjective partial function $\Psi : \prod_{m=0}^n k_m \rightarrow N$, we refer to each $\Psi^{-1}(i)$ as a *fiber*. We call a fiber of size one a *singleton*.

We now work towards an upper bound ($\approx \prod_{m=0}^n k_m$) for N . Suppose we want to show that $\text{RT}_N^1 \not\leq_{\text{w}} \prod_{m=0}^n \text{RT}_{k_m}^1$ for some N . Towards a contradiction, we may (by Corollary 5.8) fix Φ_m , $0 \leq m \leq n$, and Ψ witnessing that $\text{rt}_N^1 \leq_{\text{sw}} \prod_{m=0}^n \text{rt}_{k_m}^1$ such that Ψ is a surjective partial function from $\prod_{m=0}^n k_m$ to N . We aim to construct $c : \omega \rightarrow N$ and some (a_0, \dots, a_n) such that (a_0, \dots, a_n) is a solution to $\Phi_0^c, \dots, \Phi_n^c$, yet $\Psi(a_0, \dots, a_n)$ is not a solution to c .

Our basic strategy is to choose N large enough so that the following combinatorial property holds for all surjective partial functions $\Psi : \prod_{m=0}^n k_m \rightarrow N$:

There is some nonempty $S \subsetneq N$ such that for any set
of (a_0, \dots, a_n) 's whose image under Ψ is exactly S , the
 (a_0, \dots, a_n) 's cover some (b_0, \dots, b_n) that maps outside S under Ψ . (*)

Assuming (*), we may construct c by repeatedly looping through colors in S : for each $i \in S$, extend constantly by i until there is some (a_0, \dots, a_n) that maps to i under Ψ , such that for all $0 \leq m \leq n$, we have that Φ_m^c has some new element of color a_m . (This must happen eventually: if c is the rt_N^1 -instance produced by extending the current finite coloring by i forever, then $\Phi_0^c, \dots, \Phi_n^c$ is a $\prod_{m=0}^n \text{rt}_{k_m}^1$ -instance with some solution (a_0, \dots, a_n) . Then $\Psi(a_0, \dots, a_n) = i$, and for each $0 \leq m \leq n$, some new element of color a_m must appear at some finite stage of Φ_m^c .)

Then for each $i \in S$, there is some (a_0, \dots, a_n) such that $\Psi(a_0, \dots, a_n) = i$ and (a_0, \dots, a_n) is a solution to $\Phi_0^c, \dots, \Phi_n^c$. But then the (a_0, \dots, a_n) 's cover some (b_0, \dots, b_n) that maps outside S under Ψ . It follows that (b_0, \dots, b_n) is also a solution to $\Phi_0^c, \dots, \Phi_n^c$. But $\Psi(b_0, \dots, b_n) \notin S$ and is hence not a solution to c , which is a contradiction. Thus $\text{rt}_N^1 \not\leq_{\text{sw}} \prod_{m=0}^n \text{rt}_{k_m}^1$, and hence $\text{RT}_N^1 \not\leq_{\text{w}} \prod_{m=0}^n \text{RT}_{k_m}^1$.

The above strategy may be applied as follows:

Proposition 5.11. *If $N > \max\{\frac{k_0 \cdot k_1}{2}, k_0 + k_1 - 1\}$, then $\text{RT}_N^1 \not\leq_{\text{w}} \text{RT}_{k_0}^1 \times \text{RT}_{k_1}^1$.*

Proof. By the previous discussion, it suffices to show that $(*)$ holds. Since $N > \frac{k_0 \cdot k_1}{2}$, by a counting argument, Ψ must have at least one singleton (a_0, a_1) . Note that there are $1 + (k_0 - 1) + (k_1 - 1) = k_0 + k_1 - 1$ many pairs in $k_0 \times k_1$ that share some color with (a_0, a_1) . But $N > k_0 + k_1 - 1$, so there is some fiber G such that none of its pairs share any colors with (a_0, a_1) . In other words, for every pair in G , the set containing it and (a_0, a_1) covers a pair outside G . Let S be the image of (a_0, a_1) and G under Ψ . Then S witnesses that $(*)$ holds. \square

Corollary 5.12. *We have that*

$$\begin{aligned} \text{RT}_4^1 &\not\leq_{\text{w}} \text{RT}_2^1 \times \text{RT}_2^1, & \text{RT}_5^1 &\not\leq_{\text{w}} \text{RT}_2^1 \times \text{RT}_3^1, \\ \text{RT}_6^1 &\not\leq_{\text{w}} \text{RT}_2^1 \times \text{RT}_4^1, & \text{RT}_6^1 &\not\leq_{\text{w}} \text{RT}_3^1 \times \text{RT}_3^1, \\ \text{RT}_7^1 &\not\leq_{\text{w}} \text{RT}_2^1 \times \text{RT}_5^1, & \text{RT}_7^1 &\not\leq_{\text{w}} \text{RT}_3^1 \times \text{RT}_4^1, \\ \text{RT}_8^1 &\not\leq_{\text{w}} \text{RT}_2^1 \times \text{RT}_6^1, & \text{RT}_8^1 &\not\leq_{\text{w}} \text{RT}_3^1 \times \text{RT}_5^1. \end{aligned}$$

Note that Proposition 5.5 implies that $\text{RT}_{k_0+k_1-1}^1 \leq_{\text{w}} \text{RT}_{k_0}^1 \times \text{RT}_{k_1}^1$. Hence all of the non-reductions in Corollary 5.12 are sharp. We will address the missing case of RT_8^1 and $\text{RT}_4^1 \times \text{RT}_4^1$ in Proposition 5.16.

We can derive more results using variations of the argument in Proposition 5.11.

Proposition 5.13. *If*

$$N > \frac{\max k_m + \prod_{m=0}^n k_m}{2},$$

then $\text{RT}_N^1 \not\leq_{\text{w}} \prod_{m=0}^n \text{RT}_{k_m}^1$.

Proof. As before, we show that $(*)$ holds. By a counting argument, Ψ must have at least $1 + \max k_m$ many (a_0, \dots, a_n) that are singletons. Among these singletons, there must be two of them that differ in at least two entries, i.e., the set consisting of these two singletons covers a new tuple of colors. We can then take S to be the image of two such singletons under Ψ . \square

We can improve on this bound asymptotically, but even then this result seems to be far from optimal.

Proposition 5.14. *If*

$$N > \max \left\{ \frac{2 + \prod_{m=0}^n k_m}{2}, \max k_m - 1 + \frac{\prod_{m=0}^n k_m}{3} \right\},$$

then $\text{RT}_N^1 \not\leq_{\text{w}} \prod_{m=0}^n \text{RT}_{k_m}^1$.

Proof. As before, we show that $(*)$ holds. Since $N > \frac{2 + \prod_{m=0}^n k_m}{2}$, the reduction Ψ must have at least three singletons.

Case 1. If there are two singletons that differ in at least two entries, then we may take S to be the image of two such singletons under Ψ , as in Proposition 5.13.

Case 2. Otherwise, all of the singletons share exactly one common entry. So there are some $0 \leq m \leq n$ and $3 \leq l \leq k_m$ such that there are exactly l many singletons and all of them are of the form $(a_0, \dots, a_{m-1}, b, a_{m+1}, \dots, a_n)$, where $b < k_m$.

We claim that there are at least $k_m + 1$ many fibers of size $< l$. If not, by a counting argument, there are at least

$$\begin{aligned}
& 1 \cdot l + 2 \cdot (k_m - l) + l \cdot (N - k_m) \\
&= l + 2k_m - 2l + lN - lk_m \\
&> l \left(\max k_m - 1 + \frac{\prod_{m=0}^n k_m}{3} \right) + 2k_m - l - lk_m \\
&\geq lk_m - l + \prod_{m=0}^n k_m + 2k_m - l - lk_m \\
&\geq \prod_{m=0}^n k_m
\end{aligned}$$

many tuples, which is a contradiction.

By the claim, there is a fiber U of size $< l$ that does not contain any tuple of the form $(a_0, \dots, a_{m-1}, b, a_{m+1}, \dots, a_n)$. Since $|U| < l$, there is a singleton $(a_0, \dots, a_{m-1}, b, a_{m+1}, \dots, a_n)$ such that b does not appear in any tuple in U . Then for any tuple in U , the set containing it and $(a_0, \dots, a_{m-1}, b, a_{m+1}, \dots, a_n)$ covers some tuple outside U , so we can take S to be the image of U and said singleton. \square

The lower bound in Proposition 5.5 is, in general, much smaller than the upper bounds in Propositions 5.11, 5.13, and 5.14. Observe that in all of our proofs, the sets S consist of two elements, at least one of which is the image of a singleton under Ψ . However, Ψ may not have any singletons, for example in a hypothetical reduction witnessing that $\text{RT}_8^1 \leq_{\text{w}} \text{RT}_4^1 \times \text{RT}_4^1$. Also, there may not be any S that has exactly two elements and satisfies (*), e.g., consider $\Psi : 4 \times 4 \rightarrow 8$ as represented in the grid below. Here Ψ maps $(i, j) \in 4 \times 4$ to the number in the $(i, j)^{\text{th}}$ position.

$$\begin{array}{cccc}
0 & 3 & 2 & 6 \\
0 & 4 & 5 & 7 \\
1 & 2 & 3 & 7 \\
1 & 4 & 5 & 6
\end{array}$$

One can check that for any $c, d < 8$, there is a point labeled c that shares a row or column with a point labeled d . That means that $S = \{c, d\}$ fails to satisfy (*).

Therefore, new techniques will be required to close the gap between our lower and upper bounds. We conclude this section by giving an ad hoc proof that $\text{RT}_8^1 \not\leq_{\text{w}} \text{RT}_4^1 \times \text{RT}_4^1$, which is the smallest case not resolved by Corollary 5.12. In order to do so, we will show that there exists some S that satisfies (*) and has exactly *three* elements.

Before specializing to the case of $\text{RT}_8^1 \not\leq_{\text{w}} \text{RT}_4^1 \times \text{RT}_4^1$, we consider a more general context: let $k_0, k_1 \geq 2$ and fix a surjective partial function $\Psi : k_0 \times k_1 \rightarrow N$ (i.e., a potential backward reduction for $\text{rt}_N^1 \leq_{\text{sw}} \text{rt}_{k_0}^1 \times \text{rt}_{k_1}^1$). We say that a collection of

three fibers is *bad* if its image under Ψ does not satisfy (*). We can characterize the bad collections of three fibers:

Lemma 5.15. *Let $k_0, k_1 \geq 2$ and let $\Psi : k_0 \times k_1 \rightarrow N$ be a surjective partial function. A collection of three fibers is bad if and only if their union contains either:*

- (1) *three pairs in a row/column (e.g., $(a, b_0), (a, b_1), (a, b_2)$), with one pair from each of the three fibers;*
- (2) *four pairs that form a rectangle (i.e., $(a_0, b_0), (a_0, b_1), (a_1, b_0), (a_1, b_1)$), with at least one pair from each of the three fibers.*

Proof. (\Leftarrow). If (1) holds, the three pairs in question do not cover any new pair. If (2) holds, pick three out of the four pairs such that one pair from each of the three fibers is picked. Then these three pairs cover exactly one other pair (the fourth). But the fourth pair is already contained in the union of the three fibers.

(\Rightarrow). Suppose that we have a bad collection of three fibers. Without loss of generality, we may pick one pair (a_i, b_i) from each fiber such that the three pairs (a_0, b_0) , (a_1, b_1) , and (a_2, b_2) witness badness.

Case 1. (a_0, b_0) , (a_1, b_1) , and (a_2, b_2) lie in the same row or column. Then they satisfy (1).

Case 2. Two out of the three pairs, say (a_0, b_0) and (a_1, b_1) , lie in the same row or column (i.e., $a_0 = a_1$ or $b_0 = b_1$). Without loss of generality, suppose that $b_0 = b_1$. Note that (a_0, b_0) , (a_0, b_1) , and (a_2, b_2) cover (a_2, b_0) , (a_0, b_2) , and (a_2, b_1) . Therefore by badness, the latter three pairs lie in the union of the three fibers.

If (a_0, b_0) , (a_1, b_1) , and (a_2, b_2) are vertices of a rectangle (i.e., $b_2 = b_0$ or $b_2 = b_1$), then we satisfy (2). Otherwise, we consider cases depending on which fiber contains (a_2, b_0) . In all cases, we satisfy either (1) or (2). See Figure 3 for an illustration.

Case 3. None of the three pairs lie in the same row or column. Note that by badness, (a_0, b_1) , (a_1, b_0) , (a_0, b_2) , (a_2, b_0) , (a_1, b_2) , and (a_2, b_1) all lie in the union of the three fibers. We consider cases depending on which fiber contains (a_2, b_1) . See Figure 4 for an illustration.

Case 3a. (a_2, b_1) and (a_0, b_0) lie in the same fiber. Then we satisfy (2): (a_1, b_2) , (a_1, b_1) , (a_2, b_1) , and (a_2, b_2) form a rectangle with at least one pair from each of the three fibers.

Case 3b. (a_2, b_1) and (a_1, b_1) lie in the same fiber. Then we consider cases depending on which fiber contains (a_2, b_0) . In all cases, we satisfy either (1) or (2).

Case 3c. (a_2, b_1) and (a_2, b_2) lie in the same fiber. We consider cases depending on which fiber contains (a_0, b_1) . The argument is symmetric to Case 3b. \square

Proposition 5.16. $\text{RT}_8^1 \not\leq_{\text{w}} \text{RT}_4^1 \times \text{RT}_4^1$.

Proof. Towards a contradiction, fix forward functionals Φ_0, Φ_1 and a surjective partial function $\Psi : 4 \times 4 \rightarrow 8$ witnessing that $\text{rt}_8^1 \leq_{\text{sw}} \text{rt}_4^1 \times \text{rt}_4^1$. If Ψ has any singletons, we can derive a contradiction using the proof of Proposition 5.11. Hence we assume that Ψ has no singletons. There are sixteen pairs in 4×4 , so Ψ must be total, and all of the eight fibers in Ψ must contain exactly two pairs each.

As discussed previously, we derive a contradiction by producing a set S that satisfies (*) and consists of three elements. In other words, we show that there is a collection of three fibers that is not bad. To that end, we give an upper bound for the number of bad collections of three fibers. Since each fiber contains exactly two




FIGURE 3. Case 2 in Lemma 5.15, assuming that $b_0 = b_1$. In the array on the top level, 0 lies in position (a_0, b_0) and 1 lies in position (a_1, b_0) , meaning that $\Psi(a_0, b_0) = 0$ and $\Psi(a_1, b_0) = 1$. We have yet to label position (a_2, b_2) . The middle level represents cases depending on whether a_2 equals some a_i , or not. If a star lies in position (a, b) , then (a, b) is known (by badness) to lie in the union of the bad collection of three fibers. Sets of pairs that satisfy (1) or (2) are underlined. The bottom level represents cases depending on which of the three fibers contains (a_2, b_0) . For example, in the array on the bottom right, 2 lies in positions (a_2, b_0) and (a_2, b_2) , meaning that $\Psi(a_2, b_0) = \Psi(a_2, b_2) = 2$ and hence (a_2, b_0) and (a_2, b_2) lie in the same fiber. Then (a_0, b_0) , (a_1, b_0) , and (a_2, b_0) lie in a column, satisfying (1).

pairs, it is either contained in a row or column, or lies in diagonal position. Let k be the number of fibers that are contained in some row or column.

First, we give an upper bound for the number of collections that satisfy (2) in Lemma 5.15. It suffices to give an upper bound for the number of rectangles that intersect at most three fibers. Such rectangles have two possible forms, and we count those cases separately.

Case 1. The rectangle contains at least one of those k fibers. There are at most $(4 - 1)k = 3k$ many such rectangles.

Case 2. The rectangle contains at least one fiber in diagonal position. There are at most $8 - k$ many such rectangles.

Therefore, there are at most $3k + (8 - k) = 2k + 8$ many rectangles that intersect at most three fibers. So there are at most $2k + 8$ many collections that satisfy (2).

Next, we give an upper bound for the number of collections that satisfy (1) in Lemma 5.15.

Case 1. If a row/column contains two fibers (and hence nothing else), then said row/column does not contribute to our upper bound. Let l be the number of such rows and columns. Note that $2l \leq k$.




FIGURE 4. Case 3 in Lemma 5.15. In the array on the top level, for each $i < 3$, the number i lies in position (a_i, b_i) , meaning that $\Psi(a_i, b_i) = i$. On the middle level, we have Case 3a on the left, followed by Cases 3b and 3c. On the bottom level, we have various subcases. For example, in the array on the bottom right, 0 lies in position (a_0, b_0) , 2 lies in position (a_0, b_1) , and 1 lies in position (a_1, b_1) . Together with (a_1, b_0) , they form a rectangle satisfying (2).

Case 2. If a row/column contains one fiber, as well as two other vertices from two different fibers, then said row/column contributes one collection to our upper bound. There are $k - 2l$ many such rows/columns.

Case 3. Finally, the remaining $8 + l - k$ many rows or columns contribute $\binom{4}{3} = 4$ collections each.

Therefore, there are at most

$$l \cdot 0 + (k - 2l) \cdot 1 + (8 + l - k) \cdot 4 = 32 - 3k + 2l \leq 32 - 2k$$

many collections that satisfy (1).

We conclude that there are at most $(2k + 8) + (32 - 2k) = 40$ bad collections of three fibers. There are $\binom{8}{3} = 56 > 40$ collections of three fibers in total, so we can define S to be the image under Ψ of any collection that is not bad. Then S satisfies (*), which is a contradiction. \square

REFERENCES

- [1] Eric P. Astor, Damir D. Dzhafarov, Reed Solomon, and Jacob Suggs. The uniform content of partial and linear orders. *Ann. Pure Appl. Logic*, 168(6):1153–1171, 2017.
- [2] Vasco Brattka. Bibliography on Weihrauch complexity, <http://cca-net.de/publications/weibib.php>.
- [3] Vasco Brattka, Guido Gherardi, and Alberto Marcone. The Bolzano-Weierstrass Theorem is the jump of Weak König’s Lemma. *Ann. Pure Appl. Logic*, 163(6):623–655, 2012.
- [4] Vasco Brattka, Guido Gherardi, and Arno Pauly. Weihrauch complexity in computable analysis. To appear, arXiv:1707.03202.

- [5] Vasco Brattka, Matthew Hendtlass, and Alexander Kreuzer. On the uniform computational content of computability theory. *Theory of Computing Systems*, 61(4), 2017.
- [6] Vasco Brattka, Rupert Hölzl, and Rutger Kuiper. Monte Carlo computability. In Heribert Vollmer and Brigitte Vallée, editors, *34th Symposium on Theoretical Aspects of Computer Science (STACS 2017)*, volume 66 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 17:1–17:14, Dagstuhl, Germany, 2017. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik.
- [7] Vasco Brattka, Akitoshi Kawamura, Alberto Marcone, and Arno Pauly. Measuring the complexity of computational content (Dagstuhl Seminar 15392). *Dagstuhl Reports*, 5(9):77–104, 2016.
- [8] Vasco Brattka, Stéphane Le Roux, Joseph Miller, and Arno Pauly. Connected choice and Brouwer's fixed point theorem. *J. Math. Log.*, 19(1), 2019.
- [9] Vasco Brattka and Arno Pauly. On the algebraic structure of Weihrauch degrees. *Log. Methods Comput. Sci.*, 14(4.4):1–36, 2018.
- [10] Vasco Brattka and Tahina Rakotoniaina. On the uniform computational content of Ramsey's theorem. *J. Symbolic Logic*, 82(4):1278–1316, 2017.
- [11] Peter A. Cholak, Carl G. Jockusch, Jr., and Theodore A. Slaman. On the strength of Ramsey's theorem for pairs. *J. Symbolic Logic*, 66(1):1–55, 2001.
- [12] Peter A. Cholak, Carl G. Jockusch, Jr., and Theodore A. Slaman. Corrigendum to: "On the strength of Ramsey's theorem for pairs". *J. Symbolic Logic*, 74(4):1438–1439, 2009.
- [13] François G. Dorais, Damir D. Dzhafarov, Jeffrey L. Hirst, Joseph R. Mileti, and Paul Shafer. On uniform relationships between combinatorial problems. *Trans. Amer. Math. Soc.*, 368(2):1321–1359, 2016.
- [14] Rodney G. Downey and Denis R. Hirschfeldt. *Algorithmic randomness and complexity. Theory and Applications of Computability*. Springer, New York, 2010.
- [15] Damir D. Dzhafarov. Strong reductions between combinatorial principles. *J. Symbolic Logic*, 81(4):1405–1431, 2016.
- [16] Damir D. Dzhafarov. Joins in the strong Weihrauch degrees. *Math. Res. Lett.*, 26:749–767, 2019.
- [17] Damir D. Dzhafarov and Carl G. Jockusch, Jr. Ramsey's theorem and cone avoidance. *J. Symbolic Logic*, 74(2):557–578, 2009.
- [18] Damir D. Dzhafarov, Ludovic Patey, Reed Solomon, and Linda Brown Westrick. Ramsey's theorem for singletons and strong computable reducibility. *Proc. Amer. Math. Soc.*, 145(3):1343–1355, 2017.
- [19] Kojiro Higuchi and Arno Pauly. The degree structure of Weihrauch reducibility. *Log. Methods Comput. Sci.*, 9(2), 2013.
- [20] Denis R. Hirschfeldt. *Slicing the Truth: On the Computable and Reverse Mathematics of Combinatorial Principles*. Lecture notes series / Institute for Mathematical Sciences, National University of Singapore. World Scientific Publishing Company Incorporated, 2014.
- [21] Denis R. Hirschfeldt, Jr. Carl G. Jockusch, Bjørn Kjos-Hanssen, Steffen Lempp, and Theodore A. Slaman. The strength of some combinatorial principles related to Ramsey's theorem for pairs. In Chitat Chong, Qi Feng, Theodore A Slaman, W Hugh Woodin, and Yue Yang, editors, *Computational Prospects of Infinity*, Lecture Notes Series, Institute for Mathematical Sciences, National University of Singapore, pages 143–161, 2008.
- [22] Denis R. Hirschfeldt and Carl G. Jockusch, Jr. On notions of computability-theoretic reduction between Π_2^1 principles. *J. Math. Log.*, 16(1):1650002, 59, 2016.
- [23] Denis R. Hirschfeldt, Richard A. Shore, and Theodore A. Slaman. The atomic model theorem and type omitting. *Trans. Amer. Math. Soc.*, 361(11):5805–5837, 2009.
- [24] Jeffrey L. Hirst and Carl Mummert. Reverse mathematics of matroids. In *Computability and Complexity*, volume 10010 of *Lecture Notes in Comput. Sci.*, pages 143–159. Springer, Cham, 2017.
- [25] Jeffrey L. Hirst and Carl Mummert. Using Ramsey's theorem once. *Arch. Math. Logic*, 58(7-8):857–866, 2019.
- [26] Carl G. Jockusch, Jr. Ramsey's theorem and recursion theory. *J. Symbolic Logic*, 37:268–280, 1972.
- [27] Jiayi Liu. RT_2^2 does not imply WKL_0 . *J. Symbolic Logic*, 77(2):609–620, 2012.
- [28] Benoit Monin and Ludovic Patey. Π_1^0 encodability and omniscient reductions. *Notre Dame J. Form. Log.*, 60:1–12, 2019.

- [29] Eike Neumann and Arno Pauly. A topological view on algebraic computation models. *J. Complexity*, 44:1–22, 2018.
- [30] David Nichols. Strong reductions between relatives of the stable Ramsey’s theorem. To appear, arXiv:1711.06532.
- [31] Ludovic Patey. Partial orders and immunity in reverse mathematics. In *Pursuit of the Universal*, volume 9709 of *Lecture Notes in Comput. Sci.*, pages 353–363. Springer, Cham, 2016.
- [32] Ludovic Patey. The weakness of being cohesive, thin or free in reverse mathematics. *Israel J. Math.*, 216(2):905–955, 2016.
- [33] Arno Pauly, Willem Fouché, and George Davie. Weihrauch-completeness for layerwise computability. *Log. Methods in Comput. Sci.*, 14, 2018.
- [34] Stephen G. Simpson. *Subsystems of Second Order Arithmetic*. Perspectives in Logic. Cambridge University Press, Cambridge, second edition, 2009.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CONNECTICUT, STORRS, CONNECTICUT, U.S.A.
Email address: `damir@math.uconn.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WISCONSIN,
U.S.A.
Email address: `junle.goh@wisc.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, ILLINOIS, U.S.A.
Email address: `drh@math.uchicago.edu`

INSTITUT CAMILLE JORDAN, UNIVERSITÉ CLAUDE BERNARD LYON 1, LYON, FRANCE
Email address: `ludovic.patey@computability.fr`

DEPARTMENT OF COMPUTER SCIENCE, SWANSEA UNIVERSITY, SWANSEA, U.K.
Email address: `a.m.pauly@swansea.ac.uk`