

Cardiovascular risk reduction with liraglutide: an exploratory mediation analysis of the LEADER trial

Running title: Exploratory mediation analysis of LEADER

John B Buse MD PhD¹, Stephen C Bain FRCP², Johannes FE Mann MD³, Michael A Nauck MD⁴, Steven E Nissen MD⁵, Stuart Pocock PhD⁶, Neil R Poulter F.Med.Sci.⁷, Richard E Pratley MD⁸, Martin Linder MSc⁹, Tea Monk Fries MD PhD⁹, David D Ørsted MD PhD⁹, Bernard Zinman MD¹⁰; on behalf of the LEADER Trial

Investigators

¹University of North Carolina School of Medicine, Chapel Hill, NC, USA;

²Swansea University Medical School, Swansea, UK;

³KfH Kidney Center, Munich, and Dept. of Nephrology, Friedrich Alexander University of Erlangen, Erlangen, Germany;

⁴Diabetes Center Bochum-Hattingen, St Josef Hospital (Ruhr-Universität Bochum), Bochum, Germany;

⁵Cleveland Clinic, Cleveland, OH, USA;

⁶Dept. of Medical Statistics, London School of Hygiene and Tropical Medicine, London, UK;

⁷School of Public Health, Imperial College London, London, UK;

⁸AdventHealth Translational Research Institute for Metabolism and Diabetes, Orlando, FL, USA;

⁹Novo Nordisk A/S, Søborg, Denmark;

¹⁰Lunenfeld–Tanenbaum Research Institute, Mt. Sinai Hospital, University of Toronto, Toronto, ON, Canada.

Corresponding author: John B Buse, MD, PhD. Address: University of North Carolina School of Medicine, 8025 Burnett Womack Building, Campus Box #7172, Chapel Hill, NC 27599-717271727172, USA. Tel: +1 919-966-0134. Email: jbuse@med.unc.edu

Article type: original article, cardiovascular and metabolic risk

Word count: 3603 (main text)

No. of tables/figures: 4

Abstract

Objective

The LEADER trial (NCT01179048) demonstrated a reduced risk of cardiovascular (CV) events for patients with type 2 diabetes who received the glucagon-like peptide-1 receptor agonist liraglutide versus placebo. The mechanisms behind this CV benefit remain unclear. We aimed to identify potential mediators for the CV benefit observed with liraglutide in the LEADER trial.

Research Design and Methods

We performed exploratory analyses to identify potential mediators of the effect of liraglutide on major adverse CV events (MACE: composite of CV death, non-fatal myocardial infarction or non-fatal stroke) from the following candidates: HbA_{1c}, body weight, urinary albumin-to-creatinine ratio (UACR), confirmed hypoglycemia, sulfonylurea use, insulin use, systolic blood pressure and LDL-cholesterol. These candidates were selected as CV risk factors on which liraglutide had an effect in LEADER such that a reduction in CV risk might result. We used two methods based on a Cox proportional hazards model, and the new Vansteelandt method designed to utilize all available information from the mediator and to control for confounding factors.

Results

Analyses using the Cox methods and Vansteelandt method indicated potential mediation by HbA_{1c} (up to 41% and 83% mediation, respectively) and UACR (up to 29% and 33% mediation, respectively) on the effect of liraglutide on MACE.

Mediation effects were small for other candidates.

Conclusions

These analyses identify HbA_{1c} and, to a lesser extent, UACR as potential mediators of the CV effects of liraglutide. Whether either is a marker of an unmeasured factor or a true mediator remains a key question that invites further investigation.

Liraglutide is a glucagon-like peptide-1 receptor agonist (GLP-1 RA) approved for the management of hyperglycemia in type 2 diabetes and for reduction of cardiovascular (CV) risk in patients with type 2 diabetes and clinical CV disease (CVD) (1,2). It is also approved at a higher dose for the treatment of obesity (3,4). The LEADER trial was initiated to assess the CV safety of liraglutide in patients with type 2 diabetes and showed that it reduced the risk of CV events, all-cause mortality and renal events compared with placebo (5,6).

Typical CV risk factors associated with type 2 diabetes (glycemic control, body weight, blood pressure, lipid profiles) are improved by GLP-1 RAs, including liraglutide (7). During the LEADER trial, liraglutide reduced HbA_{1c} and body weight, along with small but significant reductions in systolic blood pressure (6). Liraglutide has also shown multiple direct anti-inflammatory and anti-atherosclerotic effects in non-clinical studies (7,8). Given these numerous effects of GLP-1 RAs, it is a challenge to pinpoint the relevant mechanisms underlying the CV benefit of liraglutide (7,9). Mediation analyses allow investigation of the associations between known, measured variables, such as the aforementioned risk factors, and outcomes, but do not necessarily identify causality.

In the present exploratory analyses, we sought to identify potential mediators for the CV benefit observed with liraglutide, using data from the LEADER trial. We explored these with several mediation methods, including a new statistical methodology designed to integrate sequential confounders (a limitation of existing methods for mediation analysis) (10).

Research Design and Methods

Trial Design

The double-blind, randomized, placebo-controlled LEADER trial assessed the CV safety of liraglutide in the context of standard care, in patients with type 2 diabetes, HbA_{1c} ≥7% (≥53 mmol/mol) and a high risk for CVD (aged ≥50 years with established CVD or chronic kidney disease stage 3 or greater; or ≥60 years with at least one risk factor for CVD, defined fully in the protocol available as supplementary material to the primary publication (6)). Patients were randomized double-blind, 1:1 to once-daily injections of liraglutide (1.8 mg or maximum tolerated dose) or placebo, both in addition to standard of care treatment for type 2 diabetes and other CV risk factors, with a follow-up period of 3.5 to 5 years (6).

The primary endpoint was the time to first occurrence of a major adverse CV event (MACE) including CV death, non-fatal myocardial infarction or non-fatal stroke (6). Secondary endpoints included the individual components of the primary composite endpoint and all-cause death (6). Other variables assessed during the trial included: HbA_{1c}, body weight, waist circumference, fasting lipids, systolic blood pressure, diastolic blood pressure, pulse rate, biochemical and hematological parameters, calcitonin levels, anti-liraglutide antibodies, urinary albumin-to-creatinine ratio (UACR), the occurrence of hypoglycemia, adverse events and concomitant medication use.

Mediation analyses

Exploratory analyses were conducted to divide the total effect of liraglutide on time to first MACE, and other time-to-event outcomes, into an indirect effect (also known as mediated effect) and a direct effect (also known as remaining effect). The indirect effect may be interpreted as the difference in time to event between liraglutide and

placebo that could be explained by a difference in another variable such as body weight or HbA_{1c}, which we will refer to as the mediator. The direct effect is, in contrast, the effect that remained after controlling for the mediator.

Analyses were performed using methods based on a Cox proportional hazards model, which we refer to hereafter as Cox methods, and a new, more complex method, termed the Vansteelandt method (10).

Cox methods

Preliminary analyses were performed using two variants of a Cox proportional hazards model to estimate the direct effect.

1. Analysis including the change from baseline in the potential mediator over time as a time-dependent covariate. At any given time point, the most recent change from baseline in the mediator, before the occurrence of the outcome in question, was included as a covariate. Until the first mediator assessment after baseline, the change was 0. This was one of two approaches used by Inzucchi and colleagues to evaluate mediated effects on CV death in the EMPA-REG OUTCOME trial (11).
2. Analysis including the updated mean value of the potential mediator (before the occurrence of the outcome in question), calculated as the trapezoidal area under the curve divided by time, as a time-dependent covariate. This was the other approach used by Inzucchi and colleagues with the intention of capturing the cumulative effect of all prior values of the mediator on the outcome, in contrast to the acute effect in approach 1 above (11).

These two approaches were performed as separate Cox regression analyses for each potential mediator with randomized treatment as a factor and the baseline

mediator value as an additional covariate, to estimate the direct effect as a log hazard ratio. The total effect was estimated using a model with only treatment as a factor, and the indirect (mediated) effect was estimated as the difference between total and direct effects.

Vansteelandt method

The Cox analyses have shortcomings in that they may not capture the full complexity of the mediator and prohibit adequate control for confounding. Furthermore, they do not immediately lend themselves to a counterfactual interpretation (conceiving outcomes in conditions altered from those that were observed) as used in the modern theory of mediation analysis (12). As the main approach for our mediation analysis, we applied a new statistical method proposed by Vansteelandt and colleagues that addresses these shortcomings and is tailored for time-to-event endpoints with repeatedly measured mediators (10).

With this new method, the direct treatment effect was estimated as the difference between probabilities of event-free survival (i.e. proportions of patients without any event) in liraglutide and placebo groups, instead of a hazard ratio (HR). It involved estimation of the counterfactual probability that a patient in the liraglutide group would be event-free at a given time if their mediator values changed to the levels that would have been seen if the patient had been assigned to placebo. Alternatively, this may be interpreted as the survival curve for the liraglutide group after adjusting the mediator values to match those observed in the placebo group. Similar to the Cox method using the updated mean, the Vansteelandt method is based on a dynamic model that incorporates new values of the mediator as they become available over time. However, with the Vansteelandt method, mediator values are not summarized into a mean, but are added to the model as separate covariates. The method is

therefore more likely to detect a mediated effect if present. All analyses undertaken allowed adjustment for baseline covariates. However, the Vansteelandt method may also adjust for post-baseline confounders, such as concomitant medication, by including them as covariates in the models. Further details of this method have been published separately (10).

The direct effect was estimated as the difference between adjusted survival curve for the liraglutide group (mediator values adjusted to match those in the placebo group) and the actual survival curve for the placebo group. The actual curve was estimated by the same method but using each patient's actual mediator values as covariates.

To estimate the indirect (mediated) effect, the adjusted survival curve for the liraglutide group was compared with the actual curve for the same group. The total effect was estimated by comparing the actual curve for the liraglutide group with the actual curve for the placebo group.

The percentage mediation was calculated as the ratio between the indirect and total effects. Complete mediation would be indicated by a percentage mediation of 100%. The percentage mediation was evaluated and presented at 3 years, since this time point corresponds to the last scheduled visit which all patients should have attended. Thereafter, the number of patients under observation for MACE decreased over time, leading to an increased statistical uncertainty of the estimated survival curves. A confidence interval (CI) for the percentage mediation was estimated using a bootstrap resampling procedure. Since the calculation was highly computer-intensive, a CI was only calculated for the strongest candidate for mediation.

Identifying potential mediators

Potential mediators examined in all methods included candidate variables on which liraglutide had a clear treatment effect and that could be related to MACE – HbA_{1c}, body weight, UACR, confirmed hypoglycemic episodes, sulfonylurea use and insulin use – and candidate mediators on which liraglutide had a modest treatment effect and that could be related to MACE – systolic blood pressure and LDL-cholesterol (Table 1). These previously published treatment effects were evaluated at month 36 (6), but the mediation analyses undertaken herein are based on the mediator values indicated in the preceding sections, such as the value at 6 or 12 months, the most recent value or all values updated over time before the event in question. Each candidate was analyzed separately using the two Cox methods and the Vansteelandt method to explore if it was a potential mediator for MACE.

Additional analyses were performed using the Vansteelandt method for the candidate with the highest apparent degree of mediation, as the mediator:

1. Analysis of time to first MACE including the other candidate mediators as confounders;
2. On-treatment analysis of time to first MACE, i.e. an analysis excluding events occurring while the patient was off-treatment for >1 day. The intention of this analysis was to remove differences in treatment adherence (to either liraglutide or placebo) between patients, since treatment adherence could influence both the mediator and the risk of MACE and thereby act as a confounder. If such confounding existed, the estimated percentage mediation would be expected to be less in the on-treatment analysis than in the main analysis;

3. Analysis of time to first MACE with reversed adjustment for the mediator, i.e. where the survival curve for the placebo group was estimated after adjusting their mediator values to match those in the liraglutide group. The purpose of this analysis was to explore if the mediator might simply be a marker of exposure to liraglutide. If true, the adjusted survival curve for the placebo group would be expected to be close to the actual placebo curve, indicating no mediation.
4. Analyses of time to all-cause death and CV death.

Results

In the LEADER trial, 9340 patients were randomized to liraglutide (N=4668) or placebo (N=4672), in addition to standard of care therapy, with a median follow-up of 3.8 years and a median exposure to study drug of 3.5 years (6). To begin this mediation analysis, the published LEADER results were examined (6). Several risk factors were improved in patients in the liraglutide versus placebo groups (Table 1), and hence were analyzed as potential mediators of the previously reported benefit of liraglutide treatment for MACE (HR: 0.87, 95% CI: 0.78; 0.97) (6).

Cox methods to determine potential mediators of MACE

Mediation analyses using the Cox methods indicated a mediation effect of HbA_{1c} (Table 2). While there was only a very small mediation effect using the change from baseline as a time-dependent covariate, analysis with the updated mean indicated that HbA_{1c} could be a substantial mediator for the treatment effect of liraglutide on MACE (Table 2). The same analyses with the other potential mediators

demonstrated at most a small effect size, with the possible exception for UACR (Table 2).

Vansteelandt method to determine potential mediators of MACE

Mediation analysis using the Vansteelandt method (10) also indicated that HbA_{1c} was the most promising candidate among those studied as a potential mediator (Figure 1, Table 3). When adjusted for HbA_{1c}, the survival curve (proportion of patients without MACE) for patients randomized to liraglutide treatment shifted down toward the placebo curve. This indicated a reduced CV benefit in the liraglutide group when the difference in HbA_{1c} between the liraglutide and placebo groups was modeled as a mediator for the MACE results (Figure 1, Table 3). A potential small mediation effect was observed with UACR (Table 3), but the estimated mediation was relatively small for the other candidates (<20%) (Table 3).

The estimated contribution of HbA_{1c} as a mediator to the effect of liraglutide on MACE at 3 years was 82.0% (95% CI: 11.7; 449.3) with the direct effect of liraglutide accounting for the remaining 18% of the total observed CV benefit (Figure 1, Table 3). Since there was no clear mediation through any other candidate variable, none of them would be likely to confound the mediation through HbA_{1c}. Nevertheless, the analysis was repeated with body weight, UACR, systolic blood pressure, LDL-cholesterol, confirmed hypoglycemic episodes, sulfonylurea use and insulin use included as confounders in the model. The indirect effect of liraglutide on MACE mediated by HbA_{1c} was estimated to be 69.2%, which is consistent with the analysis without confounders considering the uncertainty in the estimates (Supplementary Figure S1).

On-treatment analysis (excluding MACE that occurred when a patient was off-treatment) showed a slightly stronger mediation effect for HbA_{1c} of 93% than the 82% observed in the first Vansteelandt mediation analysis (without confounders) (Supplementary Figure S2). The reversed adjustment analysis showed a lower but still substantial percentage mediation by HbA_{1c} of 47% (Supplementary Figure S3).

Vansteelandt method investigating all-cause death and CV death

The same types of analyses were applied to all-cause death and CV death. The effects of liraglutide on these endpoints in LEADER appeared to be mediated by HbA_{1c} to some extent (Supplementary Figure S4).

Conclusions

These mediation analyses of the LEADER trial data, with the Cox methods and the new Vansteelandt method, suggest that HbA_{1c} could be a significant mediator of the CV benefits observed with liraglutide. There was no evidence for a mediation effect with any other candidates that were assessed in the trial, with the possible exception of UACR. A mediation effect of HbA_{1c} on mortality was also observed. However, it is important to note that these analyses cannot distinguish between the possibilities that HbA_{1c} could be a mediator or a marker of unmeasured factors that are also affected by treatment with liraglutide, and which themselves affect CV outcomes. For example, a large change in HbA_{1c} could be indicative of adherence to treatment, including concomitant medications. Contrary to the hypothesis that HbA_{1c} could be acting as a marker of treatment adherence, an on-treatment analysis showed treatment adherence was unlikely to be a confounding factor, as the percentage mediation was not reduced compared with the main analysis. A related hypothesis is

that HbA_{1c} could be acting as a marker of exposure to liraglutide specifically, but this was not supported by the reversed adjustment analysis in which the association between HbA_{1c} and CV outcomes was also apparent in the placebo group.

Modern CV outcome trials with the GLP-1 RA lixisenatide, dipeptidyl peptidase (DPP)-4 inhibitors (alogliptin, saxagliptin and sitagliptin) and insulin (insulin glargine), on a background of standard of care therapy, showed CV safety but failed to show benefits on CV outcomes despite improvements in HbA_{1c} in the active- versus placebo-treatment groups (13-17). More recently, results from other CV outcome trials have shown CV benefits for the sodium-glucose linked transporter-2 (SGLT-2) inhibitors empagliflozin and canagliflozin and the GLP-1 RAs liraglutide, semaglutide, albiglutide and dulaglutide (6,18-22). In others, non-significant trends for reductions in CV events (MACE) were reported (for dapagliflozin – SGLT-2 inhibitor; exenatide [once-weekly] and semaglutide [oral] – GLP-1RAs) (23-25). Changes in HbA_{1c} from baseline with the study treatment versus placebo in these more recent trials ranged from -0.36% to -1.0% (6,18-25), which were marginally larger than changes in the previous trials that showed no CV benefits (range of changes in HbA_{1c}: -0.2% to -0.36% (13-17)). These modest changes in HbA_{1c} across all of these trials with varying CV outcome results (including variation within drug classes) suggest that mechanisms other than reductions in HbA_{1c} account for the CV benefits with liraglutide, semaglutide, albiglutide, dulaglutide, empagliflozin and canagliflozin (6,13-25). For example, within Harmony Outcomes, there was a relatively small (-0.5%) difference in HbA_{1c} between albiglutide and placebo at 16 months, but a relatively large (22%) risk reduction in MACE (22).

Epidemiological data support the association between HbA_{1c} and CV and mortality outcomes (26,27). Data from more than 250,000 people in the Swedish National

Diabetes Register found that HbA_{1c} was a strong predictor of all-cause death, myocardial infarction, stroke and heart failure (27). These epidemiological data appear to be supported by a meta-analysis using results from early CV outcome trials of intensive glucose control in patients with type 2 diabetes (UKPDS, PROactive, ADVANCE, VADT and ACCORD – trials with varying results for CV benefits) showing modest benefits for CV outcomes (28). (6,28). Taken as a whole, the data strongly suggest that the choice of therapies used to reduce HbA_{1c}, and not just HbA_{1c} reduction itself, has an important impact on CV outcomes.

For SGLT-2 inhibitors, it has been proposed that hemodynamic effects – reducing blood pressure and intravascular volume – and diuretic effects could be underlying mechanisms for the CV benefits (29,30). These hypotheses are supported by a mediation analysis of data from EMPA-REG OUTCOME, which identified hematocrit and hemoglobin – markers of effects on plasma volume – as potential key mediators for the effects of empagliflozin on CV death, with smaller mediation effects attributed to renal function and glycemic control (11). Though HbA_{1c} appears to have a smaller mediation effect relative to hematocrit and hemoglobin for empagliflozin, it was identified as a potential mediator in the EMPA-REG OUTCOME trial, based on simple Cox methods (11). The EMPA-REG OUTCOME analyses showed that change from baseline HbA_{1c} mediated 3% of the effect of empagliflozin on CV death while updated mean HbA_{1c} mediated 23% of the effect, suggesting a greater impact of chronic exposure to elevated HbA_{1c} during the trial (11). Similarly, our analyses of LEADER data using the same two Cox methods (but with the primary endpoint of time to first MACE) showed little effect of change as a time-dependent covariate, but a larger mediation effect of updated mean HbA_{1c}.

The percentage mediation of HbA_{1c} in LEADER is difficult to compare across different analysis methods that we used. However, the Cox analyses appeared to support the main Vansteelandt mediation analyses in identifying HbA_{1c} as the main candidate for a mediator, among those analyzed, of the effect of liraglutide on CV outcomes.

The occurrence of hypoglycemia was shown to be associated with an increased risk of CV events in LEADER (31). However, beyond HbA_{1c} and UACR, little to no mediation effect was identified for the other candidates analyzed, including hypoglycemia. We therefore speculated that factors such as reduction in hypoglycemia combined with a reduction in HbA_{1c} with liraglutide might explain the apparent mediation by HbA_{1c}. However, results were largely unchanged when hypoglycemia was included as a confounder (along with other candidates in the mediator model).

Our mediation analyses were *post hoc* and, in contrast to the primary analysis of the trial, they are not protected against confounding through patient randomization. The inability to distinguish true mediators from markers is the key limitation of this present and other published mediation analyses. Our analyses are limited by the trial data; only candidate mediators that were examined or measured during the trial could be evaluated. Future trials may help to allow examination of a wider range of potential mediators through collection of biobank samples. However, even in a large trial like LEADER, it was not possible to estimate the percentage mediation with precision. The estimates were subject to substantial uncertainty reflected by the wide CI for the percentage mediation through HbA_{1c}. It is important to note that there is uncertainty in the total treatment effect of liraglutide in the primary analysis that constitutes the starting point for this mediation analysis, and which, based on the 95% CI, could be

a 3% to 22% reduced risk of MACE compared with placebo (6). Furthermore, although the Vansteelandt method has the advantage that it improves the ability to detect a mediated effect, it does not allow us to distinguish what components of the mediator might contribute to the mediated effect, i.e. whether it is the early response, the average level over time within patients, the variability within patients or a combination of several such components. Also, this analysis was limited to MACE, which is comprised of several individual CV diseases, all of which have potentially different underlying pathologies and therefore mediators. These limitations emphasize that the results should be interpreted with caution and only be considered hypothesis-generating.

In summary, these mediation analyses have identified HbA_{1c} as a potential mediator of the CV effects of liraglutide. We did not identify any mediation effects for less well-studied but possible candidate mediators, which are also risk factors for CV events, including weight and hypoglycemia. Similar to all other mediation analyses, we cannot necessarily infer causality, and whether HbA_{1c} is a marker of an unmeasured factor or a true mediator remains a key question. Based on existing evidence, we consider it unlikely that HbA_{1c} is a true mediator of the CV benefit observed with liraglutide and this finding warrants further investigation.

Acknowledgments

The LEADER trial (ClinicalTrials.gov number: NCT01179048) and this *post hoc* analysis were funded by Novo Nordisk A/S.

The authors thank all trial personnel and participants; and Charlie Hunt and Izabel James, of Watermeadow Medical, an Ashfield company (funded by Novo Nordisk),

for medical writing and editorial assistance. A list of the LEADER Committee Members and Investigators can be found in the Supplementary Material.

Author contributions

JBB, JFEM, MAN, SEN, SP, NRP and BZ were members of the LEADER trial steering committee; JBB, JFEM, MAN, SEN, NRP and BZ are current members of the LEADER publication committee; all contributed significantly to the design and conduct of the study and acquisition of clinical data. SCB and REP are members of the LEADER publication committee, were trial investigators, co-chaired the global expert panel, and contributed to conduct of the study and acquisition of clinical data. ML performed the statistical analyses. All authors reviewed and interpreted the data, and were involved in drafting and critical revision of the manuscript. All authors approved the final version of the manuscript and take full responsibility for the content. JBB is the guarantor of this work and, as such, had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the analyses.

Declaration of interests

JBB's contracted consulting fees are paid to the University of North Carolina by Adocia, AstraZeneca, Dance Biopharm, Eli Lilly, MannKind, NovaTarg, Novo Nordisk, Senseonics, vTv Therapeutics, and Zafgen. He receives grant support for clinical trials from Novo Nordisk, Sanofi, and vTv Therapeutics. He is a consultant to Cirius Therapeutics Inc, CSL Behring, Neurimmune AG, and Pendulum Therapeutics. He holds stock options in Mellitus Health, PhaseBio, Stability Health, and Pendulum Therapeutics. He is supported by grants from the National Institutes

of Health (UL1TR002489, U01DK098246, UC4DK108612, U54DK118612), PCORI and ADA.

SCB has received research grants (includes principal investigator, collaborator or consultant and grants) from Healthcare and Research Wales (Welsh Government) and Novo Nordisk. He has received other research support from Healthcare and Research Wales (Welsh Government) and honoraria from Novo Nordisk, Sanofi, Lilly, Boehringer Ingelheim, and Merck. He has an ownership interest in Glycosmedia.

JFEM has received speaker honoraria from Astra, Boehringer Ingelheim, Eli Lilly & Co, Medice, Novo Nordisk, Roche, and Vifor Pharma; research support from the European Union, Canadian Institutes of Health Research, Astra, Boehringer Ingelheim, Novo Nordisk, Roche, and Sandoz; and consultation fees from Astra, Bayer, Novo Nordisk, and Vifor Pharma.

MAN has served on advisory boards or consulted for AstraZeneca, Boehringer Ingelheim, Eli Lilly & Co., Fractyl, GlaxoSmithKline, Hanmi, Intarcia/Servier, Menarini/Berlin Chemie, Merck, Sharp & Dohme, and Novo Nordisk. His institution has received grant support from AstraZeneca, Boehringer Ingelheim, Eli Lilly & Co., GlaxoSmithKline, Intarcia, Menarini/Berlin-Chemie, Merck, Sharp & Dohme, Novartis Pharma, Novo Nordisk A/S, and Sun Pharma. He has served on speakers' bureau of AstraZeneca, Boehringer Ingelheim, Eli Lilly & Co., GlaxoSmithKline, Menarini/Berlin Chemie, Merck, Sharp & Dohme, and Novo Nordisk A/S.

SEN has received reimbursement from Boehringer Ingelheim and grant support from Pfizer, Amgen, Esperion Therapeutics, Cerenis Therapeutics, The Medicines

Company, AstraZeneca, Takeda Pharmaceuticals, Orexigen Therapeutics, and Eli Lilly.

SP has no relevant conflict of interest to disclose.

NRP has received speaker fees from Servier, Takeda, and Novo Nordisk; has served on advisory boards for AstraZeneca and Novo Nordisk. He has received grants for his research group relating to type 2 diabetes from Diabetes UK, NIHR EME, Julius Clinical, and the British Heart Foundation, with a pending grant from Novo Nordisk.

REP has received research grants from Gilead Sciences, Lexicon Pharmaceuticals, Ligand Pharmaceuticals Inc., Lilly, Merck, Novo Nordisk, Sanofi-Aventis US LLC, and Takeda; has acted as a speaker for AstraZeneca, Novo Nordisk, and Takeda; and as a consultant for AstraZeneca, Boehringer Ingelheim, Eisai, Inc., GlaxoSmithKline, Janssen Scientific Affairs LLC, Ligand Pharmaceuticals Inc., Lilly, Merck, Novo Nordisk, Pfizer, and Takeda. All payments are made directly to his employer (AdventHealth).

TMF and DDØ are full time employees of, and hold stock in, Novo Nordisk.

ML is a full time employee of Novo Nordisk.

BZ has received consulting fees from Merck, Novo Nordisk, Sanofi-Aventis, Eli Lilly, AstraZeneca, Janssen, and Boehringer Ingelheim.

References

1. Novo Nordisk A/S. Victoza (liraglutide) Summary of Product Characteristics. 2017
2. Novo Nordisk A/S. Victoza (liraglutide) - Prescribing information highlights. In *FDA*, 2017
3. Novo Nordisk A/S. Saxenda (liraglutide) Summary of Product Characteristics. 2017
4. Novo Nordisk Inc. Saxenda (liraglutide) Highlights of Prescribing Information. 2017
5. Mann JFE, Ørsted DD, Brown-Frandsen K, et al. Liraglutide and Renal Outcomes in Type 2 Diabetes. *N Engl J Med* 2017;377:839-848
6. Marso SP, Daniels GH, Brown-Frandsen K, et al. Liraglutide and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med* 2016;375:311-322
7. Drucker DJ. The Cardiovascular Biology of Glucagon-like Peptide-1. *Cell Metab* 2016;24:15-30
8. Rakipovski G, Rolin B, Nohr J, et al. The GLP-1 Analogs Liraglutide and Semaglutide Reduce Atherosclerosis in ApoE(-/-) and LDLr(-/-) Mice by a Mechanism That Includes Inflammatory Pathways. *JACC Basic Transl Sci* 2018;3:844–857
9. Nauck MA, Meier JJ, Cavender MA, Abd El Aziz M, Drucker DJ. Cardiovascular Actions and Clinical Outcomes With Glucagon-Like Peptide-1 Receptor Agonists and Dipeptidyl Peptidase-4 Inhibitors. *Circulation* 2017;136:849-870
10. Vansteelandt S, Linder M, Vandenberghe S, Steen J, Madsen J. Mediation analysis of time-to-event endpoints accounting for repeatedly measured mediators subject to time-varying confounding. *Statistics in Medicine* 2019;38:4828-4840

11. Inzucchi SE, Zinman B, Fitchett D, et al. How Does Empagliflozin Reduce Cardiovascular Mortality? Insights From a Mediation Analysis of the EMPA-REG OUTCOME Trial. *Diabetes Care* 2018;41:356-363
12. Vanderweele T. *Explanation in causal inference: methods for mediation and interaction*. Oxford University Press, 2015
13. Green JB, Bethel MA, Armstrong PW, et al. Effect of Sitagliptin on Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med* 2015;373:232-242
14. Origin Trial Investigators, Gerstein HC, Bosch J, et al. Basal insulin and cardiovascular and other outcomes in dysglycemia. *N Engl J Med* 2012;367:319-328
15. Pfeffer MA, Claggett B, Diaz R, et al. Lixisenatide in Patients with Type 2 Diabetes and Acute Coronary Syndrome. *N Engl J Med* 2015;373:2247-2257
16. Scirica BM, Bhatt DL, Braunwald E, et al. Saxagliptin and cardiovascular outcomes in patients with type 2 diabetes mellitus. *N Engl J Med* 2013;369:1317-1326
17. White WB, Cannon CP, Heller SR, et al. Alogliptin after acute coronary syndrome in patients with type 2 diabetes. *N Engl J Med* 2013;369:1327-1335
18. Gerstein HC, Colhoun HM, Dagenais GR, et al. Dulaglutide and cardiovascular outcomes in type 2 diabetes (REWIND): a double-blind, randomised placebo-controlled trial. *Lancet* 2019;394:121-130
19. Marso SP, Bain SC, Consoli A, et al. Semaglutide and Cardiovascular Outcomes in Patients with Type 2 Diabetes. *N Engl J Med* 2016;375:1834-1844
20. Neal B, Perkovic V, Mahaffey KW, et al. Canagliflozin and Cardiovascular and Renal Events in Type 2 Diabetes. *N Engl J Med* 2017;377:644-657
21. Zinman B, Wanner C, Lachin JM, et al. Empagliflozin, Cardiovascular Outcomes, and Mortality in Type 2 Diabetes. *N Engl J Med* 2015;373:2117-2128

22. Hernandez AF, Green JB, Janmohamed S, et al. Albiglutide and cardiovascular outcomes in patients with type 2 diabetes and cardiovascular disease (Harmony Outcomes): a double-blind, randomised placebo-controlled trial. *Lancet* 2018;392:1519-1529
23. Holman RR, Bethel MA, Mentz RJ, et al. Effects of Once-Weekly Exenatide on Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med* 2017;377:1228-1239
24. Husain M, Birkenfeld AL, Donsmark M, et al. Oral Semaglutide and Cardiovascular Outcomes in Patients with Type 2 Diabetes. *N Engl J Med* 2019;381:841-851
25. Wiviott SD, Raz I, Bonaca MP, et al. Dapagliflozin and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med* 2019;380:347-357
26. Khaw KT, Wareham N, Bingham S, Luben R, Welch A, Day N. Association of hemoglobin A1c with cardiovascular disease and mortality in adults: the European prospective investigation into cancer in Norfolk. *Ann Intern Med* 2004;141:413-420
27. Rawshani A, Rawshani A, Franzen S, et al. Risk Factors, Mortality, and Cardiovascular Outcomes in Patients with Type 2 Diabetes. *N Engl J Med* 2018;379:633-644
28. Ray KK, Seshasai SR, Wijesuriya S, et al. Effect of intensive control of glucose on cardiovascular outcomes and death in patients with diabetes mellitus: a meta-analysis of randomised controlled trials. *Lancet* 2009;373:1765-1772
29. Sattar N, McLaren J, Kristensen SL, Preiss D, McMurray JJ. SGLT2 Inhibition and cardiovascular events: why did EMPA-REG Outcomes surprise and what were the likely mechanisms? *Diabetologia* 2016;59:1333-1339
30. Staels B. Cardiovascular Protection by Sodium Glucose Cotransporter 2 Inhibitors: Potential Mechanisms. *Am J Cardiol* 2017;120:S28-S36

31. Zinman B, Marso SP, Christiansen E, et al. Hypoglycemia, Cardiovascular Outcomes, and Death: The LEADER Experience. *Diabetes Care* 2018;41:1783-1791

Tables

Table 1. Potential mediators of the CV benefit with liraglutide identified from LEADER.

Variable	Difference between liraglutide and placebo groups (95% CI)	Difference between liraglutide and placebo groups: z-score
HbA _{1c} , %	ETD*: -0.40 (-0.45; -0.34)	-13.5
Body weight, kg	ETD*: -2.3 (-2.5; -2.0)	-16.2
UACR	ETR*: 0.83 (0.78; 0.88)	-6.1
Rate of confirmed hypoglycemia during the trial	Rate ratio: 0.81 (0.74; 0.88)	-4.8
Systolic blood pressure, mmHg	ETD*: -1.2 (-1.9; -0.5)	-3.3
LDL-cholesterol	ETR*: 0.98 (0.96; 0.99)	-3.0
Patients who initiated insulin use during the trial, n [%]	1336 [28.8%] vs 2019 [43.2%]	-
Patients who initiated SU use during the trial, n [%]	349 [7.5%] vs 505 [10.8%]	-

*Change from baseline to 36 months (ETR was used when the data were log-transformed, due to their known distributions)

ETD, estimated treatment difference; ETR, estimated treatment ratio; HbA_{1c}, glycated hemoglobin;

SU, sulfonylurea; UACR, urinary albumin-to-creatinine ratio

Table 2. Cox modelling for mediation of time to first MACE with liraglutide versus placebo by HbA_{1c}, body weight, UACR, systolic blood pressure, LDL-cholesterol, confirmed hypoglycemic episodes, sulfonylurea use and insulin use, assessed using two approaches.

Analysis	Risk of MACE with liraglutide vs placebo, HR (95% CI)	Percentage mediation, % (95% CI)[†]
Total treatment effect (primary LEADER analysis)	0.868 (0.778; 0.968)	–
Treatment effect adjusted for:		
HbA_{1c}		
Change as time-dependent covariate	0.880 (0.787; 0.984)	9.9 (-11.4; 59.0)
Updated mean* as time-dependent covariate	0.920 (0.820; 1.033)	41.1 (8.6; 161.0)
Body weight		
Change as time-dependent covariate	0.860 (0.770; 0.962)	-7.4 (-43.9; 13.3)
Updated mean* as time-dependent covariate	0.864 (0.772; 0.968)	-3.9 (-47.0; 26.2)
UACR		
Change as time-dependent covariate	0.899 (0.804; 1.004)	21.9 (7.5; 106.4)
Updated mean* as time-dependent covariate	0.907 (0.812; 1.013)	28.6 (12.3; 132.8)
Confirmed hypoglycemia		
Change as time-dependent covariate	0.868 (0.778; 0.967)	-0.3 (-4.5; 2.2)
Updated mean* as time-dependent covariate	0.868 (0.778; 0.968)	-0.1
SBP		
Change as time-dependent covariate	0.874 (0.784; 0.975)	5.2 (-0.6; 21.2)

Updated mean* as time-dependent covariate	0.877 (0.787; 0.979)	7.7 (0.2; 31.7)
LDL-cholesterol		
Change as time-dependent covariate	0.869 (0.778; 0.970)	0.6 (-4.2; 7.7)
Updated mean* as time-dependent covariate	0.869 (0.778; 0.970)	0.7 (-4.2; 8.8)
Insulin use		
Change as time-dependent covariate	0.889 (0.796; 0.993)	16.9 (3.1; 85.7)
Updated mean* as time-dependent covariate	0.894 (0.801; 0.998)	20.7
SU use		
Change as time-dependent covariate	0.861 (0.772; 0.960)	-5.6 (-25.2; -0.3)
Updated mean* as time-dependent covariate	0.865 (0.776; 0.965)	-2.3

*Calculated as the trapezoidal area under the curve divided by time. †CI for the percentage mediation was estimated using a bootstrap resampling procedure.

HbA1c, glycated hemoglobin; HR, hazard ratio; MACE, major adverse cardiovascular event; SBP, systolic blood pressure; SU, sulfonylurea; UACR, urinary albumin-to-creatinine ratio


Table 3. Vansteelandt modelling for mediation of time to first MACE with liraglutide versus placebo: by HbA_{1c}, body weight, UACR, systolic blood pressure, LDL-cholesterol, confirmed hypoglycemic episodes, sulfonylurea use and insulin use.

Variable	Estimated proportion of patients without MACE at 3 years			Percentage mediation, % (95% CI)
	Liraglutide	Liraglutide, adjusted for candidate mediator	Placebo	
HbA _{1c}	0.893	0.881	0.878	82.0 (11.7;449.3)
Body weight	0.893	0.891	0.879	14.3
UACR	0.892	0.888	0.880	33.3
Confirmed hypoglycemia	0.893	0.892	0.878	6.7
Systolic blood pressure	0.893	0.891	0.878	13.3
LDL-cholesterol	0.893	0.893	0.879	0.0
Insulin use	0.892	0.890	0.878	14.3
SU use	0.893	0.891	0.878	13.3

HbA_{1c}, glycated hemoglobin; SU, sulfonylurea; UACR, urinary albumin-to-creatinine ratio

Figure legend

Figure 1. Probability of no MACE occurring in patients treated with liraglutide and placebo, and with adjustment for HbA_{1c} as a candidate mediator using the Vansteelandt method.



HbA_{1c}, glycated hemoglobin; MACE, major adverse cardiovascular event