

## Biological Sciences, Ecology

**Title:** Climate drives the geography of marine consumption by changing predator communities

**Authors:** Matthew A. Whalen<sup>1,2,3\*</sup>, Ross D.B. Whippo<sup>3,4</sup>, John J. Stachowicz<sup>5</sup>, Paul H. York<sup>6</sup>, Erin Aiello<sup>7</sup>, Teresa Alcoverro<sup>8,9</sup>, Andrew H. Altieri<sup>10,11</sup>, Lisandro Benedetti-Cecchi<sup>12</sup>, Camilla Bertolini<sup>13</sup>, Midoli Bresch<sup>1</sup>, Fabio Bulleri<sup>12</sup>, Paul E. Carnell<sup>14</sup>, Stéphanie Cimon<sup>15</sup>, Rod M. Connolly<sup>16</sup>, Mathieu Cusson<sup>15</sup>, Meredith S. Diskin<sup>17</sup>, Elrika D'Souza<sup>9</sup>, Augusto A.V. Flores<sup>18</sup>, F. Joel Fodrie<sup>19</sup>, Aaron W.E. Galloway<sup>4</sup>, Leo C. Gaskins<sup>20</sup>, Olivia J. Graham<sup>21</sup>, Torrance C. Hanley<sup>22</sup>, Christopher J. Henderson<sup>23</sup>, Clara M. Hereu<sup>24</sup>, Margot Hessing-Lewis<sup>1,25</sup>, Kevin A. Hovel<sup>26</sup>, Brent B. Hughes<sup>27</sup>, A. Randall Hughes<sup>22</sup>, Kristin M. Hultgren<sup>28</sup>, Holger Jänes<sup>14,29</sup>, Dean S. Janiak<sup>30</sup>, Lane N. Johnston<sup>31</sup>, Pablo Jorgensen<sup>32</sup>, Brendan P. Kelaher<sup>33</sup>, Claudia Kruschel<sup>34</sup>, Brendan S. Lanham<sup>35</sup>, Kun-Seop Lee<sup>36</sup>, Jonathan S. Lefcheck<sup>3</sup>, Enrique Lozano-Álvarez<sup>37</sup>, Peter I. Macreadie<sup>14</sup>, Zachary L. Monteith<sup>1</sup>, Nessa E. O'Connor<sup>13</sup>, Andrew D. Olds<sup>23</sup>, Jennifer K. O'Leary<sup>7,38</sup>, Christopher J. Patrick<sup>39</sup>, Oscar Pino<sup>40</sup>, Alistair G.B. Poore<sup>41</sup>, Michael A. Rasheed<sup>6</sup>, Wendel W. Raymond<sup>42</sup>, Katrin Reiss<sup>43</sup>, O. Kennedy Rhoades<sup>5,30</sup>, Max T. Robinson<sup>44</sup>, Paige G. Ross<sup>45</sup>, Francesca Rossi<sup>46</sup>, Thomas A Schlacher<sup>23</sup>, Janina Seemann<sup>11</sup>, Brian R. Silliman<sup>20</sup>, Delbert L. Smee<sup>17</sup>, Martin Thiel<sup>40,47,48</sup>, Richard K.F. Unsworth<sup>44</sup>, Brigitta I. van Tussenbroek<sup>37</sup>, Adriana Vergés<sup>41</sup>, Mallarie E. Yeager<sup>26</sup>, Bree K. Yednock<sup>49</sup>, Shelby L. Ziegler<sup>19</sup>, J. Emmett Duffy<sup>3</sup>

**Affiliations:** <sup>1</sup>Hakai Institute, PO Box 309, Heriot Bay, BC V0P 1H0, Canada. <sup>2</sup>Biodiversity Research Centre, University of British Columbia, 2212 Main Mall Vancouver, BC V6T 1Z4, Canada. <sup>3</sup>Tennenbaum Marine Observatories Network, Smithsonian Institution, Edgewater, Maryland 21037 USA. <sup>4</sup>University of Oregon, Oregon Institute of Marine Biology, 63466 Boat Basin Rd., Charleston, OR 97420, USA. <sup>5</sup>Bodega Marine Laboratory, University of California, Davis, 2099 Westside Road, Bodega Bay, CA 94923, USA. <sup>6</sup>James Cook University, Centre for Tropical Water and Aquatic Ecosystem Research, PO Box 6811, Cairns, Qld, Australia. <sup>7</sup>Biology Department, California Polytechnic State University, 1 Grand Avenue, San Luis Obispo, CA 93422 USA. <sup>8</sup>Centre d'Estudis Avançats de Blanes (Spanish National Research Council), Blanes, Girona, Spain. <sup>9</sup>Nature Conservation Foundation, 1311, "Amritha", 12th Main, Opp Kodava Samaj, Vijayanagar 1st Stage, Mysore 570 017, India. <sup>10</sup>Department of Environmental Engineering Sciences, University of Florida, Gainesville, FL 32611, USA. <sup>11</sup>Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancon, Republic of Panama. <sup>12</sup>Department of Biology and CoNISMa, Via Derna 1 I-56126, University of Pisa, Pisa, Italy. <sup>13</sup>Trinity College Dublin, The University of Dublin, College Green, Dublin 2, Ireland. <sup>14</sup>School of Life and Environmental Sciences, Centre for Integrative Ecology, Deakin University, Geelong, Victoria 3220, Australia. <sup>15</sup>Département des sciences fondamentales & Québec-Océan, Université du Québec à Chicoutimi, 555, Boulevard de l'Université, Chicoutimi, QC G7H 2B1, Canada. <sup>16</sup>Griffith University, Australian Rivers Institute - Coast & Estuaries, School of Environment and Science. Gold Coast, Queensland 4222, Australia. <sup>17</sup>Dauphin Island Sea Lab, 101 Bienville Blvd., Dauphin Island, AL 36528, USA. <sup>18</sup>Centre for Marine Biology, University of São Paulo, <sup>19</sup>Institute of Marine Sciences, University of North Carolina at Chapel Hill, 3431 Arendell St., Morehead City, NC 28557 USA. <sup>20</sup>Nicholas School for

the Environment, Duke University Marine Lab, 135 Duke Marine Lab Road, Beaufort, NC 28516, USA.  
<sup>21</sup>Cornell University, Dept. of Ecology and Evolutionary Biology, 215 Tower Rd, Ithaca, NY 14853.  
<sup>22</sup>Northeastern University, Marine Science Center, 430 Nahant Road, Nahant, MA 01907, USA. <sup>23</sup>School  
of Science and Engineering, University of the Sunshine Coast, Maroochydore QLD 4558, Australia.  
<sup>24</sup>Universidad Autónoma de Baja California, Carretera Ensenada, Tijuana 3917, C.P. 22860 Ensenada,  
Baja California, Mexico, <sup>25</sup>Institute of the Oceans & Fisheries, 2202 Main Mall, Vancouver, BC V6T  
1Z4, Canada. <sup>26</sup>Department of Biology, San Diego State University, 5500 Campanile Drive, San Diego,  
CA 92182, USA. <sup>27</sup>Department of Biology, Sonoma State University, 1801 E. Cotati Ave, Rohnert Park,  
CA 94954, USA. <sup>28</sup>Department of Biology, Seattle University, 901 12th Ave., Seattle, WA 98122, USA.  
<sup>29</sup>Estonian Marine Institute, University of Tartu, Mäealuse 14, 12618 Tallinn, Estonia, <sup>30</sup>Smithsonian  
Marine Station, 701 Seaway Dr, Fort Pierce, FL 34949, USA. <sup>31</sup>School of Marine Sciences and Policy,  
University of Delaware, 700 Pilottown Rd., Lewes DE 19958, USA. <sup>32</sup>Geomare, A. C., Paseo del  
Pedregal 82, C.P. 22860 Ensenada, Baja California, Mexico. <sup>33</sup>National Marine Science Centre, Southern  
Cross University, 2 Bay Drive, Coffs Harbour, NSW 2450, Australia. <sup>34</sup>Department of Ecology,  
University of Zadar, Agronomy and Aquaculture M. Pavlinovica 1, 23000 Zadar, Croatia. <sup>35</sup>School of  
Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, NSW 2052,  
Australia. <sup>36</sup>Department of Biological Sciences, Pusan National University, Busan 46241, Korea.  
<sup>37</sup>Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Prolongación  
Niños Heroes S/N, Puerto Morelos, Quintana Roo 77580, Mexico. <sup>38</sup>California Sea Grant, 9500 Gilman  
Dr., La Jolla, CA 92093-0232, USA <sup>39</sup>Texas A&M University Corpus Christi, HRI 121 Unit 5892, 6300  
Ocean Drive, Corpus Christi TX 78412, USA. <sup>40</sup>Facultad Ciencias del Mar, Universidad Católica del  
Norte, Coquimbo, Chile. <sup>41</sup>School of Biological, Earth and Environmental Sciences, University of New  
South Wales, Sydney, NSW 2052, Australia. <sup>42</sup>University of Alaska Fairbanks Fisheries Division, 17101  
Point Lena Loop Rd., Juneau 99801, USA. <sup>43</sup>Faculty for Biosciences and Aquaculture, Nord University,  
8049 Bodø, Norway. <sup>44</sup>Seagrass Ecosystem Research Group, College of Science, Swansea University,  
Wallace Building, Swansea SA2 8PP, UK. <sup>45</sup>Virginia Institute of Marine Science, PO Box 350,  
Wachapreague, VA 23480, USA. <sup>46</sup>CNRS, National Centre for Scientific Research, Université Côte  
d'Azur, CNRS, FRE 3729 ECOMERS, Nice 6100, France. <sup>47</sup>Millenium Nucleus Ecology and Sustainable  
Management of Oceanic Islands ESMOI, Coquimbo, Chile. <sup>48</sup>Center for Advanced Studies in Arid  
Systems, CEAZA, Coquimbo, Chile. <sup>49</sup>South Slough National Estuarine Research Reserve, P.O. Box  
5417, Charleston, OR 97420, USA.

**Competing interests:** authors declare no competing interests

**Corresponding Author:**

Matthew Adam Whalen

Postdoctoral Scholar in Marine Biodiversity

Hakai Institute, Smithsonian MarineGEO, University of British Columbia

Department of Botany

3200-6270 University Blvd.

Vancouver, BC, V6T 1Z4, CANADA

1-703-909-5732

[mawhal@gmail.com](mailto:mawhal@gmail.com)

**Abstract:** The global distribution of primary production and consumption by humans (fisheries) are well-documented, but we have no map linking the central ecological process of consumption within food webs to temperature and other ecological drivers. Using standardized assays that span 105 degrees of latitude on four continents, we show that rates of bait consumption by generalist predators in shallow marine ecosystems are tightly linked to both temperature and the composition of consumer assemblages. Unexpectedly, rates of consumption peaked at mid-latitudes (25-35°) in both northern and southern hemispheres across both seagrass and unvegetated sediment habitats. This pattern contrasts with terrestrial systems, where biotic interactions reportedly weaken away from the equator, but it parallels an emerging pattern of a subtropical peak in marine biodiversity. The higher consumption at mid-latitudes was closely related to the type of consumers present, which explained rates of consumption better than consumer density, biomass, species diversity, or habitat. Indeed, the apparent effect of temperature on consumption was mostly driven by temperature-associated turnover in consumer community composition. Our findings reinforce the key influence of climate warming on altered species composition and highlight its implications for the functioning of Earth's ecosystems.

**Keywords:** latitudinal diversity gradient, macroecology, biogeography, trophic processes, climate, seagrass

**Significance Statement (max 120 words):**

Consumption transfers energy and materials through food chains and fundamentally influences ecosystem productivity. Therefore, mapping the distribution of consumer feeding intensity is key to understanding how environmental changes influences biodiversity, with consequent effects on trophic transfer and top-down impacts through food webs. Our global comparison of standardized bait consumption in shallow coastal habitats finds a previously unrecognized peak in feeding intensity away from the equator that is better explained by the presence of particular consumer families than by latitude or temperature. This study complements recent demonstrations that changes in biodiversity can have similar or larger impacts on ecological processes than those of climate.

## Text

Latitudinal diversity gradients have stimulated decades of research, much of it invoking a decline from tropics to poles in rates of key biological processes and species interactions (1–3). General explanations for ecological patterns across latitude, however, remain elusive in part because so many environmental and biological variables change in parallel with latitude. As a result, the connections between ecological patterns and processes at global scales remain controversial (4–6). This uncertainty has recently been amplified by demonstrations that diversity of many modern and ancient lineages peaks at mid-latitudes rather than at the equator, particularly in the ocean (7, 8).

Temperature is among the environmental factors that vary most consistently with latitude, and is a fundamental driver of biological processes. Metabolic theory mechanistically links environmental temperatures to a suite of biological processes, including metabolism and trophic transfer (9–12). For example, metabolic theory predicts that per capita consumption rates of ectothermic consumers should follow increased metabolic needs and activity, and increase with rising temperature (13). But the traits of consumers, their abundance, and the resources available to them also change with temperature (14, 15), so total consumption rates may be poorly predicted by temperature alone. Separating these effects requires data on geographic variation in consumption.

Our understanding of global variation in top-down processes in marine systems is largely indirect, based on inferences from distributions of organismal traits such as body size and morphological defenses (e.g., 2, 16) and on comparisons of primary production, prey abundance, and predator abundance (e.g., 17, 18). The distribution and abundance of species respond to ecosystem productivity, reproductive rate, migration, mortality, and evolutionary history, all of which are modified by temperature (e.g., 19, 20). But while the spatial distribution of primary production and human predation (fishing) are well-documented (21, 22), we have only sparse empirical measurements of geographic variation in consumption by natural predators, which is needed to predict trophic transfer and prey abundance. Thus, we lack a global map linking the central ecological process of consumption to temperature and other drivers.

We approached this problem by measuring feeding intensity of generalist marine consumers across 42 sites around the globe representing two widespread coastal habitats: seagrass and unvegetated sediments. These critical habitats provide shelter and fuel primary and secondary production (23–25), and seagrass persistence is in turn linked to trophic processes, as mid-level carnivores consume herbivores that can facilitate or damage seagrass (26–30). Therefore, understanding consumption by mid-level predators is key to seagrass conservation and restoration efforts (30).

To compare consumption rigorously around the world's coasts, we used a simple, standardized feeding assay, offering small (~1 cm) discs of dried squid mantle as bait. Squid is attractive to many generalist marine predators, including mid-sized fishes and crustaceans, which we surveyed in both seagrass and unvegetated habitats. Importantly, standardizing bait allowed us to estimate consumption rates on a comparable basis while avoiding confounding influences of geographic variation in prey type, prey behavior, and prey defenses. Previous studies (e.g., 31, 32) have shown changes in feeding on standardized prey across latitude, but ours covers nearly the entire latitudinal range of seagrasses on four continents, and multiple ocean basins in both northern and southern hemispheres (our study, 38°S-67°N; seagrass, 45°S-70°N). This allowed us to test the consistency of latitudinal gradients in consumption in two widespread habitats, and to begin disentangling the role of correlated drivers. Based on previous studies documenting broad-scale patterns in biodiversity (3, 33, 34), prey defense (2), trophic interactions (5, 35, 36), and metabolism (12), we hypothesized that rates of bait consumption would increase with temperature toward the equator.

## Results and Discussion

Contrary to our expectations, bait consumption peaked away from the equator in both hemispheres (25-35° North and South), and was consistent between seagrass and sediment habitats (Fig. 1A), despite slightly higher density, richness, and biomass of consumers in seagrass (SI Appendix, Fig. S1). Although our sampling near the equator was relatively sparse, the decline in measured bait consumption at the lowest latitudes was robust in two independent regions (SI Appendix, Fig. S2). This mid-latitude peak was supported by a hump-shaped relationship between absolute latitude and consumption (comparison of models with and without second-order polynomial; quadratic model Akaike weight  $w_{quad} = 0.86$ ). In contrast, satellite-derived mean annual sea surface temperature decreased monotonically with latitude (SST; Fig. 1C), and the hump-shaped relation of SST to consumption was much stronger than that for latitude ( $w_{quad} > 0.99$ ). This nonlinear relationship between SST and consumption was also supported in two independent, well-sampled transects along the Northwest Atlantic ( $w_{quad} = 0.97$ ) and Southwest Pacific ( $w_{quad} = 0.99$ ; SI Appendix, Fig. S2), strengthening the inference that the equatorial dip in consumption reflects a response to temperature, rather than some other correlate of latitude.

Fish and crabs were the main consumers of bait at all sites, and their taxonomic composition varied greatly across sites and with temperature (Fig. 2). Because species and genera of coastal animals differ markedly across ocean basins and hemispheres, we analyzed consumer composition at the level of taxonomic families, which allowed us to describe gradual shifts in global consumer biogeography across broad environmental gradients, while maintaining functional distinction among taxa. We used presence vs. absence rather than abundance data because we wanted to explore consumer composition and abundance separately. Mean annual sea surface temperature alone explained 16% of the dissimilarity in fish and crustacean

assemblages across sites (canonical analysis coefficient = 0.13,  $P = 0.001$ ). However, a single unconstrained index of compositional dissimilarity (Principal Components Axis 1 in Fig. 2A) explained 19.5% of the total variation in consumer composition, and separated cool-water assemblages from warm assemblages (37). Indeed, this index of consumer composition was a stronger predictor of bait consumption than water temperature (either measured during the assays or using mean annual sea surface temperatures; SI Appendix, Fig. S3), latitude, consumer density and biomass, or estimates of ocean productivity, fishing pressure, or human population density (Table 1). Consumer density and biomass only became important predictors of consumption once we reduced the dataset to include only those consumer families whose presence was associated with increased consumption rates (see Methods, SI Appendix, Table S1). When viewed as a simple network of causal relationships, the effect of thermal environment on consumption rate was largely indirect, being mediated by consumer community composition, and this remained true even when allowing for a nonlinear relationship between consumption rate and mean annual sea surface temperature (Fig. 3). Roughly three-quarters of the total effect of thermal environment on bait consumption flowed indirectly through differences in consumer taxa in different climates.

Locations with high consumption rates had consumer assemblages composed largely of invertivores and omnivores that actively forage on or just above the seafloor (SI Appendix, Fig. S3-4). Actively swimming foragers should consume bait faster due to increased encounter rates, all else being equal, and arguably consumption by these foragers might rise more rapidly with temperature than for more sedentary taxa. Video evidence confirmed the association of key families with high consumption. Porgies (family Sparidae), for example, removed bait at every site where they were observed in video footage (SI Appendix, Table S2) and the presence of this family showed the strongest association with consumption rate in our analysis of community composition (Fig. 2B, SI Appendix, Table S1).

The equatorial decline in bait consumption appears to be related to consumer community composition, as many of the actively foraging taxa associated with high consumption rates, including porgies, half-beaks (Hemiramphidae), and grunts (Haemulidae), were rare or absent at the sites closest to the equator (SI Appendix, Fig. S5). Some of these consumer families (e.g., porgies) are known from low latitude waters but were not recorded in our surveys (38). It is possible that larger enemies reduce mesopredator abundance or restrict their foraging (39, 40) to a greater extent at low latitudes, but we do not have any direct evidence to support this hypothesis. Similarly, human harvest or other activities could have restricted the abundance of these key consumers, but we know of no reason to expect this to be more intense at low latitudes, as many of the middle and high latitude sites in this study are heavily influenced human activities, including overfishing (41). Alternatively, environmental tolerances could limit consumer access or abundance in shallow seagrass habitats at low latitude (42). There is currently little evidence to evaluate these explanations.

Our finding that feeding intensity peaked at mid-latitudes differs strongly from most previous studies on latitudinal gradients in species interactions (5, 12, 31, 36). Non-linear ecological transitions between warm-temperate and subtropical locations might help explain this result. These regions feature rapid transition between thermal guilds of consumers with cool- vs. warm-water affinities (37) and these biogeographic transitions are correlated with shifts in the relative strength of bottom-up vs. top-down processes that are directly and indirectly related to temperature (17). We find it interesting that such transitions co-occur in similar climatic regions: transitions in consumption from this study ( $\sim 19\text{--}22^\circ\text{C}$  SST), transitions in dominant fish guilds ( $\sim 21\text{--}25^\circ\text{C}$  SST; 37), and transitions in top-down vs. bottom-up control ( $\sim 17\text{--}20^\circ\text{C}$  temperature 0-200m; 17). These comparisons suggest that zones of biogeographic and trophic transitions associated with climate are also areas of transition for consumptive pressure by small mesopredators.

The weak and inconsistent differences we found in consumption rates between seagrass vs. unvegetated sediment habitats (SI Appendix, Fig. S6-7) was surprising given decades of research showing that the structure provided by seagrasses and other foundation species can strongly influence predator-prey relationships (43–46). While we found no overall difference in consumer composition between seagrass and unvegetated habitats (permutation test,  $P = 0.75$ ), consumer densities were generally higher inside than outside of seagrass habitat (SI Appendix, Fig. S1). Thus, any protection provided by seagrass structure may have been offset by consumer aggregation in seagrass. Yet the consistency of latitudinal patterns in consumption between the two habitats suggests that broad-scale environment and consumer biogeography had stronger influences on consumption than local differences in habitat structure.

Our feeding assay used identical bait at all sites to isolate the effect of consumer activity from the behavioral and morphological traits of prey, which vary widely across space. No single bait will attract all predators equally; ours targeted the small to medium-sized generalists that dominate many shallow marine habitats. Thus, the consumption rates that we describe are relative measures of one-half of a predator-prey interaction (i.e., consumption in the absence of prey behavior and other trait variation). Whereas this design cannot completely characterize species interactions, standardization more rigorously estimates how potential consumption varies across the globe. Our assays did not measure top-down control per se, but the kind of information we gathered is critical to understanding trophic interactions, including cascading effects in seagrass ecosystems (30, 47, 48), because it measures the willingness of consumers to eat prey of a certain size. The consistency of our results across ocean basins and hemispheres, along with similar recent findings for pelagic top predators (49), suggests that the mid-latitude peak in marine consumption is indeed general. The importance of particular predator taxa and traits in the geography of consumption we found parallels the outsized role of endothermy in the effectiveness of marine predators (e.g., 50, 51), including in some seagrass meadows (47, 48).

We focused on the smaller ectothermic consumers that consume herbivorous invertebrates that can be critical to seagrass persistence (32), but these are potential prey of larger endotherms like fishing birds and small marine mammals, so endothermy might well influence the broader food webs we studied. However, given that many endothermic predators are most abundant and diverse in cooler regions of the world ocean (51), we would expect the distribution of their collective feeding intensity to differ from the pattern we observed.

Changing climate, overfishing, and global species introductions are altering the biogeography of marine life and the composition of communities (52, 53), with wide-ranging effects on ecosystems (54), including in seagrass habitats (55). Shifting biogeography of consumers can alter community and ecosystem structure and processes (56, 57) independent of temperature, as we show here. Simultaneously, warming can directly influence physiology of ectothermic consumers (e.g., metabolic demand, activity; 58). We show that variation in water temperature influences marine trophic process mainly indirectly by changing consumer community composition. The hump-shaped relationship between temperature and consumption we found suggests that predation and trophic transfer may intensify at middle to high latitudes and decline near the equator as the world ocean warms and species continue to shift their ranges. Such shifts in species ranges and biomass distributions could lead to large changes in consumption, with repercussions for community structure and trophic flows through marine food webs. It is already clear that many ectotherms are expanding or contracting their ranges with climate change (59, 60). Our findings suggest that such distributional shifts may affect ecological processes as much or more than those predicted based only on temperature effects on metabolism.

## **Materials and Methods**

We assessed rates of consumption using a simple, standardized field assay (61). We tethered a 1-1.3 cm diameter piece of dried squid mantle with monofilament to a fiberglass garden stake (hereafter, ‘squidpop’), that we inseted into the sediment such that the bait dangled 20-30 cm above the sediment surface in or just above the seagrass canopy. At most sites, we deployed 20-30 squidpops within a seagrass meadow and 20-30 squidpops in nearby unvegetated sediments (SI Appendix, Table S3). We checked the squidpops for presence (1) or absence (0) of bait after one hour and again after 24 hours. Most sites repeated this assay for a total of three deployments in each of the two habitat types, and measured water temperature during each deployment.

To characterize variation in environments across the range of the study, we drew upon several publicly-available datasets with global-scale variables of interest. We accessed sea surface temperature and chlorophyll records using Bio-ORACLE (62), which packages data collected by the Aqua-MODIS satellite. We used mean annual sea surface temperature (SST) because it showed stronger relationships with consumption rate than maximum or minimum annual SST and it summarizes well the differences between thermal conditions across the globe (Fig. 1). Most assays were conducted during the summer, but differences in timing of assays generated



variation in *in situ* temperature that altered the rank-order of our estimates of the thermal environments compared to sea surface temperature, making nearby sites appear less similar environmentally (Fig. 3). We used mean annual chlorophyll *a* as a proxy for surface ocean primary productivity across sites. We also accessed data on human population densities from the Gridded Population of the World (63), which we used as a proxy for local human disturbance. Finally, we accessed fishing pressure data from the Sea Around Us project (64) using the R package ‘searoundus’ (65).

At most sites (30 of 42 sites) we also conducted consumer surveys in the areas adjacent to feeding assays. These surveys used hand-pulled seines in seagrass and unvegetated sediment habitats to sample epibenthic consumers (mainly fishes, but also large crustaceans) adjacent to feeding assays. All consumers were identified, counted, and released. The total lengths of the first 25 individual fish of each species were also measured. We used these data to estimate consumer density, size distribution, biomass, and diversity, as well as to generate a species list for each site. Species lists from five additional sites were added using data from video footage and diver transects (FL, India, Italy, Yuca1, Yuca2; SI Appendix Movie S1). Biomass estimates were calculated using length-weight regressions available in Fishbase (66).

For each of the squidpop assays we independently fitted an exponential decay model and estimated consumption rate (bait loss through time) using the slope parameter. We then used the resulting rate estimates as data points in subsequent analysis.

We predicted individual consumption rates in generalized linear mixed effects models (logit link, random intercepts for sites) using a variety of potential abiotic and biotic drivers, and compared models using Akaike Information Criterion corrected for small sample size calculated using the R package ‘bbmle’ (67). We also explored a variety of polynomial terms and LOESS curves to investigate possibilities of non-linear relationships between temperature and consumption, although for model comparison we only included linear terms. We restricted the data used in model comparison to the 27 sites for which we had the full complement of explanatory variables. For simplicity, and because our analysis was largely exploratory using a large set of candidate explanatory variables, we compared models with individual predictor variables only. All mixed models were fitted using maximum likelihood in the package ‘lme4’ in R (68).

When estimating consumer species (alpha) diversity, we used both species richness and Hurlbert’s probability of interspecific encounter as effective numbers of species (69). We also wanted to investigate changes in consumer community composition across sites (beta diversity), but given the scale of our analysis and the large biogeographic gradients we captured, comparing composition in terms of species identity was not possible. Species level overlap was low among sites, especially across ocean basins and hemispheres, so we chose to compare composition (presence-absence) at the level of families across sites using Raup-Crick dissimilarities. While

this metric has been used to investigate small spatial scale differences in species composition within regions (70), we use it here to investigate global among-site turnover of consumers at higher taxonomic levels. In order to visualize and quantify major axes of community variation, we used Principal Coordinates Analysis (PCoA) to ordinate consumer communities based on their dissimilarities, and then assess how these dissimilarities related to the thermal environment and consumption rate. We used a combination of unconstrained (PCoA) and constrained (Canonical Analysis of Principal Coordinates) techniques in this analysis. Unconstrained ordination reduces dimensionality of the dataset by finding orthogonal axes of decreasing variation in the dataset, while constrained ordination uses a regression-based approach to define a set of axes of interest a priori based on explanatory variables (71). We used the resultant axes from unconstrained ordination (PCoA) as explanatory variables in the models described above because the unconstrained ordination does not require a priori assumptions about which factors are important. We also assessed relationships between consumer community composition and thermal environment by constraining the first ordination axis to SST or *in situ* temperature. Multivariate analyses were performed using the R package ‘vegan’ (72).

In order to identify which consumer families were positively and negatively associated with consumption intensity across sites, we constrained the first axis of the ordination to align with our estimates of consumption rate. Then we selected families that mapped onto the positive side of this axis as candidate taxa driving spatial variation in consumption rate, and calculated the density and biomass of these consumers at sites with seining data (27 of 42 sites). Finally, we compared the results from multivariate analysis to direct observations of squidpop attacks and bait removal from video footage captured at 14 sites (SI Appendix Table S2).

To explore which predator traits might explain feeding intensity in our assays, we scored six traits for each taxon in our dataset (416 morphospecies in 103 taxonomic families). Four traits were derived from Fishbase (feeding habit, lateral body shape; 73) and Reef Life Survey (trophic group, water column usage; 74). A fifth binomial trait scored whether each taxon is an actively swimming forager or tends towards ambush or sit-and-wait behavior, either on the benthos or in the water column. We applied the most common value of this trait to all taxa in each family, but we acknowledge that variation in foraging activity can occur within families. Traits missing in these databases were filled using expert opinion of co-authors and available trait information from related taxa. A sixth continuous trait describing body size as the average total length of each taxon (carapace width for crabs) was calculated from our seining data. Whereas published total length estimates are available for many taxa, we opted to use length estimates from our own dataset because many taxa only utilize seagrass and other nearshore habitats for part of their development, when they may differ greatly from the species’ maximum size. Using the R package ‘FD’ (75) we calculated community-level weighted means of trait values to derive estimates of average conditions for each of the six individual traits in each site and habitat combination in the dataset, and we calculated a variety of functional diversity metrics (functional

richness, functional dispersion, functional evenness, functional diversity, and Rao's Q) following published methods (75–77). For all consumer functional diversity metrics and all community-level weighted means except body size we used presence-absence data instead of weighting by relative abundance so that we could include sites with seining and video data. We did weight mean consumer body size estimates by relative abundance because we only had size estimates from seine sampling. Weighting by abundance did not qualitatively change the results. We regressed each functional diversity metric and each community-level weighted mean trait against consumption rate individually using the linear mixed effects models described above.

We tested whether consumer composition mediated the influence of mean annual SST on consumption rates using the package 'mediation' in R (78). Because we found support for a hump-shaped relationship between SST and consumption rate, we tested whether consumer composition mediated the non-linear relationship between temperature and consumption rate (using 33 of 42 sites with all three variables). We modeled the relationships using 1) smooth terms for SST on consumption rate and a linear term for composition (PCoA1) on consumption rate in a generalized additive model (GAM; logit link function; R package 'mgcv' 79) and 2) a general linear model for SST on composition. We report the standardized linear regression coefficients, estimated degrees of freedom for smoothed GAM terms (and associated chi-square statistic), estimates of the mediation effect and direct effect, and the proportion of the direct effect of SST mediated by composition for the second mediation analysis, along with 95% confidence intervals around estimates of the direct effect, mediation effect, and the proportion mediated. All models in mediation analysis used data that were averaged at the level of habitats within sites, which is the lowest level of pairwise comparisons we can make between squidpop assays and consumer composition.

All analyses were performed in R version 3.5.3 (80). Data and analyses for this project are available at <https://doi.org/10.5281/zenodo.3998836>.

## References

1. T. Dobzhansky, Evolution in the tropics. *Am. Sci.* **38**, 208–221 (1950).
2. G. J. Vermeij, *Evolution and Escalation: An Ecological History of Life* (Princeton University Press, 1993).
3. A. Z. Krug, D. Jablonski, J. W. Valentine, K. Roy, Generation of Earth's first-order biodiversity pattern. *Astrobiology* **9**, 113–124 (2009).
4. S. A. Levin, The problem of pattern and scale in ecology: the Robert H. MacArthur Award Lecture. *Ecology* **73**, 1943–1967 (1992).
5. D. W. Schemske, G. G. Mittelbach, H. V. Cornell, J. M. Sobel, K. Roy, Is there a latitudinal gradient in the importance of biotic interactions? *Annu. Rev. Ecol. Evol. Syst.* **40**, 245–269 (2009).

- 434 6. A. T. Moles, S. P. Bonser, A. G. B. Poore, I. R. Wallis, W. J. Foley, Assessing the evidence  
435 for latitudinal gradients in plant defence and herbivory. *Funct. Ecol.* **25**, 380–388 (2011).
- 436 7. M. G. Powell, V. P. Beresford, B. A. Colaianne, The latitudinal position of peak marine  
437 diversity in living and fossil biotas: latitudinal position of peak marine diversity. *J.*  
438 *Biogeogr.* **39**, 1687–1694 (2012).
- 439 8. C. Chaudhary, H. Saeedi, M. J. Costello, Bimodality of latitudinal gradients in marine  
440 species richness. *Trends Ecol. Evol.* **31**, 670–676 (2016).
- 441 9. J. F. Gillooly, J. H. Brown, West, Geoffrey B., V. M. Savage, Charnov, Eric L., Effects of  
442 size and temperature on metabolic rate. *Science* **293**, 2248–2251 (2001).
- 443 10. J. H. Brown, J. F. Gillooly, A. P. Allen, V. M. Savage, G. B. West, Toward a metabolic  
444 theory of ecology. *Ecology* **85**, 1771–1789 (2004).
- 445 11. G. G. Mittelbach, *et al.*, Evolution and the latitudinal diversity gradient: speciation,  
446 extinction and biogeography. *Ecol. Lett.* **10**, 315–331 (2007).
- 447 12. J. R. Schramski, A. I. Dell, J. M. Grady, R. M. Sibly, J. H. Brown, Metabolic theory  
448 predicts whole-ecosystem properties. *Proc. Natl. Acad. Sci.* **112**, 2617–2622 (2015).
- 449 13. M. I. O’Connor, B. Gilbert, C. J. Brown, Theoretical predictions for how temperature  
450 affects the dynamics of interacting herbivores and plants. *Am. Nat.* **178**, 626–638 (2011).
- 451 14. M. I. O’Connor, M. F. Piehler, D. M. Leech, A. Anton, J. F. Bruno, Warming and resource  
452 availability shift food web structure and metabolism. *PLoS Biol.* **7**, e1000178 (2009).
- 453 15. A. I. Dell, S. Pawar, V. M. Savage, Temperature dependence of trophic interactions are  
454 driven by asymmetry of species responses and foraging strategy. *J. Anim. Ecol.* **83**, 70–84  
455 (2014).
- 456 16. M. B. J. Harfoot, *et al.*, Emergent global patterns of ecosystem structure and function from  
457 a mechanistic general ecosystem model. *PLOS Biol.* **12**, e1001841 (2014).
- 458 17. D. G. Boyce, K. T. Frank, B. Worm, W. C. Leggett, Spatial patterns and predictors of  
459 trophic control in marine ecosystems. *Ecol. Lett.* **18**, 1001–1011 (2015).
- 460 18. P. D. van Denderen, M. Lindegren, B. R. MacKenzie, R. A. Watson, K. H. Andersen,  
461 Global patterns in marine predatory fish. *Nat. Ecol. Evol.* **2**, 65–70 (2018).
- 462 19. J. R. Bernhardt, J. M. Sunday, P. L. Thompson, M. I. O’Connor, Nonlinear averaging of  
463 thermal experience predicts population growth rates in a thermally variable environment.  
464 *Proc. R. Soc. B Biol. Sci.* **285**, 10 (2018).
- 465 20. D. Pauly, On the interrelationships between natural mortality, growth parameters, and mean  
466 environmental temperature in 175 fish stocks. *ICES J. Mar. Sci.* **39**, 175–192 (1980).

- 467 21. C. B. Field, Primary production of the biosphere: integrating terrestrial and oceanic  
468 components. *Science* **281**, 237–240 (1998).
- 469 22. R. A. Watson, *et al.*, Global marine yield halved as fishing intensity redoubles. *Fish Fish.*  
470 **14**, 493–503 (2013).
- 471 23. J. Cebrian, Patterns in the fate of production in plant communities. *Am. Nat.* **154**, 449  
472 (1999).
- 473 24. S. L. Williams, K. L. Heck, Jr., “Seagrass Community Ecology” in *Marine Community*  
474 *Ecology*, M. D. Bertness, S. D. Gaines, M. E. Hay, Eds. (Sinauer, 2001), pp. 317–337.
- 475 25. M. Sievers, *et al.*, The role of vegetated coastal wetlands for marine megafauna  
476 conservation. *Trends Ecol. Evol.* **34**, 807–817 (2019).
- 477 26. J. F. Valentine, J. E. Duffy, “The Central Role of Grazing in Seagrass Ecology” in  
478 *Seagrasses: Biology, Ecology and Conservation*, A. W. D. Larkum, R. J. Orth, C. M.  
479 Duarte, Eds. (Springer Netherlands, 2006), pp. 463–501.
- 480 27. J. E. Duffy, J. P. Richardson, K. E. France, Ecosystem consequences of diversity depend on  
481 food chain length in estuarine vegetation. *Ecol. Lett.* **8**, 301–309 (2005).
- 482 28. S. P. Baden, A. Emanuelsson, L. Pihl, C.-J. Svensson, P. Åberg, Shift in seagrass food web  
483 structure over decades is linked to overfishing. *Mar. Ecol. Prog. Ser.* **451**, 61–73 (2012).
- 484 29. P. L. Reynolds, J. P. Richardson, J. E. Duffy, Field experimental evidence that grazers  
485 mediate transition between microalgal and seagrass dominance. *Limnol. Oceanogr.* **59**,  
486 1053–1064 (2014).
- 487 30. J. E. Duffy, A. R. Hughes, P.-O. Moksnes, “Ecology of Seagrass Communities” in *Marine*  
488 *Community Ecology and Conservation*, 2nd Edition, M. D. Bertness, J. F. Bruno, B. R.  
489 Silliman, J. J. Stachowicz, Eds. (Sinauer Associates Inc., 2013), pp. 271–297.
- 490 31. P. L. Reynolds, *et al.*, Latitude, temperature, and habitat complexity predict predation  
491 pressure in eelgrass beds across the Northern Hemisphere. *Ecology* **99**, 29–35 (2018).
- 492 32. C. A. Musrri, *et al.*, Variation in consumer pressure along 2500 km in a major upwelling  
493 system: crab predators are more important at higher latitudes. *Mar. Biol.* **166** (2019).
- 494 33. P. D. Mannion, P. Upchurch, R. B. J. Benson, A. Goswami, The latitudinal biodiversity  
495 gradient through deep time. *Trends Ecol. Evol.* **29**, 42–50 (2014).
- 496 34. B. Worm, D. P. Tittensor, *A Theory of Global Biodiversity* (Princeton University Press,  
497 2018) <https://doi.org/10.2307/j.ctt1zkjz6q> (September 19, 2019).
- 498 35. A. L. Freestone, R. W. Osman, G. M. Ruiz, M. E. Torchin, Stronger predation in the tropics  
499 shapes species richness patterns in marine communities. *Ecology* **92**, 983–993 (2011).

- 500 36. T. Roslin, *et al.*, Higher predation risk for insect prey at low latitudes and elevations.  
501 *Science* **356**, 742–744 (2017).
- 502 37. R. D. Stuart-Smith, G. J. Edgar, N. S. Barrett, S. J. Kininmonth, A. E. Bates, Thermal  
503 biases and vulnerability to warming in the world’s marine fauna. *Nature* (2015)  
504 <https://doi.org/10.1038/nature16144> (May 10, 2018).
- 505 38. K. E. Carpenter, Ed., “The Living Marine Resources of the Western Central Atlantic.  
506 Volume 3: Bony fishes part 2 (Opistognathidae to Molidae), sea turtles and marine  
507 mammals” in *FAO Species Identification Guide for Fishery Purposes and American Society*  
508 *of Ichthyologists and Herpetologists Special Publication No.5*, (FAO, 2002), pp. 1375–  
509 2127.
- 510 39. F. Micheli, C. H. Peterson, Estuarine Vegetated Habitats as Corridors for Predator  
511 Movements. *Conserv. Biol.* **13**, 869–881 (1999).
- 512 40. E. M. P. Madin, *et al.*, Multi-Trophic Species Interactions Shape Seascape-Scale Coral Reef  
513 Vegetation Patterns. *Front. Ecol. Evol.* **7** (2019).
- 514 41. B. K. Eriksson, *et al.*, Effects of altered offshore food webs on coastal ecosystems  
515 emphasize the need for cross-ecosystem management. *AMBIO* **40**, 786 (2011).
- 516 42. J. M. Sunday, A. E. Bates, N. K. Dulvy, Global analysis of thermal tolerance and latitude in  
517 ectotherms. *Proc. R. Soc. Lond. B Biol. Sci.*, 1823–1830 (2011).
- 518 43. K. A. Hovel, R. N. Lipcius, Habitat fragmentation in a seagrass landscape: patch size and  
519 complexity control blue crab survival. *Ecology* **82**, 1814–1829 (2001).
- 520 44. R. J. Orth, K. L. Heck, Jr., J. van Montfrans, Faunal communities in seagrass beds: a review  
521 of the influence of plant structure and prey characteristics on predator-prey relationships.  
522 *Estuaries* **7**, 339–350 (1984).
- 523 45. K. L. Main, Predator avoidance in seagrass meadows: prey behavior, microhabitat  
524 selection, and cryptic coloration. *Ecology* **68**, 170–180 (1987).
- 525 46. J. Mattila, *et al.*, Increased habitat structure does not always provide increased refuge from  
526 predation. *Mar. Ecol. Prog. Ser.* **361**, 15–20 (2008).
- 527 47. B. B. Hughes, *et al.*, Recovery of a top predator mediates negative eutrophic effects on  
528 seagrass. *Proc. Natl. Acad. Sci.* **110**, 15313–15318 (2013).
- 529 48. A. C. Huang, M. Essak, M. I. O’Connor, Top–down control by great blue herons *Ardea*  
530 *herodias* regulates seagrass-associated epifauna. *Oikos* **124**, 1492–1501 (2015).
- 531 49. M. Roesti, *et al.*, Pelagic fish predation is stronger at temperate latitudes than near the  
532 equator. *Nat. Commun.* **11**, 1–7 (2020).

50. Y. Y. Watanabe, K. J. Goldman, J. E. Caselle, D. D. Chapman, Y. P. Papastamatiou, Comparative analyses of animal-tracking data reveal ecological significance of endothermy in fishes. *Proc. Natl. Acad. Sci.* **112**, 6104–6109 (2015).
51. J. M. Grady, *et al.*, Metabolic asymmetry and the global diversity of marine predators. *Science* **363**, eaat4220 (2019).
52. M. L. Pinsky, B. Worm, M. J. Fogarty, J. L. Sarmiento, S. A. Levin, Marine taxa track local climate velocities. *Science* **341**, 1239–1242 (2013).
53. D. J. McCauley, *et al.*, Marine defaunation: Animal loss in the global ocean. *Science* **347**, 1255641 (2015).
54. J. A. Estes, *et al.*, Trophic downgrading of planet earth. *Science* **333**, 301–306 (2011).
55. G. A. Hyndes, *et al.*, Accelerating tropicalization and the transformation of temperate seagrass meadows. *BioScience* **66**, 938–948 (2016).
56. L. D. Zeidberg, B. H. Robison, Invasive range expansion by the Humboldt squid, *Dosidicus gigas*, in the eastern North Pacific. *Proc. Natl. Acad. Sci.* **104**, 12948–12950 (2007).
57. I. M. Côté, S. J. Green, M. A. Hixon, Predatory fish invaders: Insights from Indo-Pacific lionfish in the western Atlantic and Caribbean. *Biol. Conserv.* **164**, 50–61 (2013).
58. B. J. Sinclair, *et al.*, Can we predict ectotherm responses to climate change using thermal performance curves and body temperatures? *Ecol. Lett.* **19**, 1372–1385 (2016).
59. R. D. Sagarin, J. P. Barry, S. E. Gilman, C. H. Baxter, Climate-related change in an intertidal community over short and long time scales. *Ecol. Monogr.* **69**, 465–490 (1999).
60. E. Sanford, J. L. Sones, M. García-Reyes, J. H. R. Goddard, J. L. Largier, Widespread shifts in the coastal biota of northern California during the 2014–2016 marine heatwaves. *Sci. Rep.* **9**, 4216 (2019).
61. J. E. Duffy, S. L. Ziegler, J. E. Campbell, P. M. Bippus, J. S. Lefcheck, Squidpops: a simple tool to crowdsource a global map of marine predation intensity. *PLOS ONE* **10**, e0142994 (2015).
62. L. Tyberghein, *et al.*, Bio-ORACLE: a global environmental dataset for marine species distribution modelling: Bio-ORACLE marine environmental data rasters. *Glob. Ecol. Biogeogr.* **21**, 272–281 (2012).
63. Center for International Earth Science Information Network - CIESIN - Columbia University, Gridded Population of the World, Version 4 (GPWv4): Population Density, Revision 10 (2017).
64. D. Pauly, D. Zeller, Eds., Sea Around Us Concepts, Design and Data (searoundus.org) (2015).

65. S. Chamberlain, R. S. Reis, *searounds: Sea Around Us API Wrapper* (2017).
66. R. Froese, D. Pauly, Fishbase. *World Wide Web Electron. Publ.* (2019).
67. B. M. Bolker, R. D. C. Team, *bbmle: Tools for General Maximum Likelihood Estimation* (2017).
68. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).
69. J. M. Chase, T. M. Knight, Scale-dependent effect sizes of ecological drivers on biodiversity: why standardised sampling is not enough. *Ecol. Lett.* **16**, 17–26 (2013).
70. J. M. Chase, N. J. B. Kraft, K. G. Smith, M. Vellend, B. D. Inouye, Using null models to disentangle variation in community dissimilarity from variation in  $\alpha$ -diversity. *Ecosphere* **2**, art24 (2011).
71. M. J. Anderson, T. J. Willis, Canonical Analysis of Principal Coordinates: a useful method of constrained ordination for ecology. *Ecology* **84**, 511–525 (2003).
72. J. Oksanen, *et al.*, *vegan: Community Ecology Package* (2018).
73. C. Boettiger, D. T. Lang, P. Wainwright, rfishbase: exploring, manipulating and visualizing FishBase data from R. *J. Fish Biol.* (2012).
74. R. D. Stuart-Smith, *et al.*, Integrating abundance and functional traits reveals new global hotspots of fish diversity. *Nature* **501**, 539–542 (2013).
75. E. Laliberté, P. Legendre, A distance-based framework for measuring functional diversity from multiple traits. *Ecology* **91**, 299–305 (2010).
76. S. Villéger, N. W. H. Mason, D. Mouillot, New multidimensional functional diversity indices for a multifaceted framework in functional ecology. *Ecology* **89**, 2290–2301 (2008).
77. Z. Botta-Dukát, Rao’s quadratic entropy as a measure of functional diversity based on multiple traits. *J. Veg. Sci.* **16**, 533–540 (2005).
78. D. Tingley, T. Yamamoto, K. Hirose, L. Keele, K. Imai, mediation: R Package for Causal Mediation Analysis. *J. Stat. Softw.* **59**, 1–38 (2014).
79. S. N. Wood, Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models. *J. R. Stat. Soc. B* **73**, 3–36 (2011).
80. R Core Team, *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, 2019).

## Acknowledgments



We would like to thank M.I. O'Connor, J.R. Bernhardt, and P.L. Thompson for helpful reviews of early versions of the manuscript. We are grateful for three anonymous reviews that greatly improved the paper and for G.J. Vermeij, who provided several rounds of thoughtful review and suggested that high-level consumers might depress mesopredator consumption at low latitudes. Funding: we acknowledge funding from the Smithsonian Institution and the Tula Foundation; Author Contributions: JED conceived of the Bitemap project, JED and RDBW generated the methodology for field research and data collection and supervised the investigations across sites, RDBW, PHY, and MAW curated the data, MAW performed the formal analyses and visualization, MAW and JED wrote the original draft, MAW, JJS, and JED wrote the final draft, all authors contributed to field investigations and to reviewing and editing the manuscript. This is contribution XXX from the Smithsonian's MarineGEO and Tennenbaum Marine Observatories Network.

## Figures

Figure 1. Distributions of bait consumption by generalist marine predators and temperature across the 42 sites in this study. A) Consumption rate of tethered dried squid bait peaks at mid-latitudes in both hemispheres. Point color represents habitat, and lines show independent quadratic GLMs fitted for each habitat type in each hemisphere. B) Map of study sites. C) Latitudinal pattern of mean annual sea surface temperature (SST).


Figure 2. The composition of consumer assemblages reflects global gradients in environmental temperatures and consumption rate. A) Principal coordinates analysis, where locations of symbols reflect compositional differences among sites and habitats based on family-level presence-absence data. Symbol color represents mean annual sea surface temperature (°C), and symbol size corresponds to bait consumption rate. B) The same ordination showing scores for consumer families driving differences in composition and consumption rate among sites. Symbol color represents average in situ temperature at sites where the predator family was observed label color represents positive (red), negative (purple), or non-significant (black) correlations with consumption rate, and body length (width for crabs) is proportional to the magnitude of the correlation. Asterisks denote families that were seen feeding on bait in video footage.



Figure 3. Predator composition mediates the effect of thermal environment on consumption rates. (A-C) Bivariate relationships between consumer composition (PCoA1, Fig. 2A), thermal environment (SST) and consumption rate. Lines show predictions from models used in mediation analysis (A, linear regression; B, logistic regression; C, generalized additive modeling). D) Paths represent causal hypotheses about relationships. Numbers next to paths leading to and from consumer composition are standardized regression coefficients and standard errors. Numbers above and below the path from thermal environment to consumption rate are estimated degrees of freedom and chi-square values for the smooth term in the presence and absence of mediation, respectively. Numbers above the path diagram are estimates of the direct and indirect (mediation) effects with 95% bootstrapped confidence intervals.




642   **Tables**

643

644   Table 1. Comparison of generalized linear mixed effects models predicting bait consumption by  
645   generalist consumers in two shallow marine habitats







**D**

Mediation Effect (AB)	0.24 ( 0.14,0.34)
Direct Effect (C)	0.09 (-0.06,0.22)
Proportion Mediated	0.73 ( 0.46,1.46)




Table 1. Comparison of generalized linear mixed effects models predicting bait consumption by generalist consumers in two shallow marine habitats

Candidate models	$k$	AICc	$\Delta$ AICc	$w_i$	$R^2$
Taxonomic Composition	3	70.3	0	0.932	0.51
Selected Abundance	3	77.0	6.7	0.033	0.39
(Sea Surface Temperature) <sup>2</sup>	4	79.2	8.9	0.011	0.66
Selected Biomass	3	79.6	9.3	0.009	0.38
Functional Richness	3	79.9	9.5	0.008	0.26
Sea Surface Temperature	3	80.5	10.2	0.006	0.39
<i>in situ</i> Temperature	3	82.8	12.5	0.002	0.31
Productivity	3	90.5	20.2	<0.001	0.12
Proportion of Active Foragers	3	90.5	20.2	<0.001	0.10
Body size	3	90.6	20.3	<0.001	0.10
Consumer Species Richness	3	90.8	20.5	<0.001	0.05
Functional Evenness	3	91.4	21.1	<0.001	0.04
Intercept-Only	2	92.0	21.7	<0.001	0
Trophic Group	6	92.1	21.8	<0.001	0.27
Functional Group Richness	3	92.2	21.9	<0.001	0.04
Feeding Type	3	92.4	22.1	<0.001	0.15
Habitat	3	93.1	22.8	<0.001	0.01
Lateral Body Shape	6	93.1	22.8	<0.001	0.23
Total Biomass	3	93.2	22.9	<0.001	0.02
Effective Number of Species	3	93.2	22.9	<0.001	0.01
Total Abundance	3	93.3	23.0	<0.001	0.02
Fishing Pressure	3	93.6	23.3	<0.001	0.02
Water Column Use	6	93.7	23.4	<0.001	0.20
Human Population Density	3	94.0	23.7	<0.001	<0.01
Rao Q	3	94.0	23.7	<0.001	<0.01
Functional Dispersion	3	94.0	23.7	<0.001	<0.01

Taxonomic Composition refers to the first axis from the PCoA of consumer assemblages (Fig. 2). Selected Abundance and Biomass refer to density or biomass of fish and decapod families selected through constrained ordination (Fig. 2B, SI Appendix, Table S1). Productivity refers to remotely-sensed mean annual chlorophyll *a*. Habitat categorically relates seagrass and unvegetated habitats. Abundance, biomass, and human population density were log<sub>10</sub>-transformed. Body size, Trophic Group, Lateral Body Shape, and Water Column Use are community-weighted mean trait values by site and habitat. We provide marginal pseudo- $R^2$  values for comparison of fixed effects. For model comparison we only included data from sites with the full complement of predictors (27 of 42 sites).