

Title: Alternate leg bounding acutely improves change of direction performance in women's team sports players irrespective of ground type

Running head: Bounding and change of direction

Eleanor Dann ^a, Samuel Quinn ^a, Mark Russell ^b, Liam P. Kilduff ^{c,d}, Anthony N. Turner ^e, Samuel P. Hills ^{a*}

^a Faculty of Health and Social Sciences, Bournemouth University, Bournemouth, United Kingdom

^b Faculty of Health, Sport and Life Sciences, Leeds Trinity University, Leeds, United Kingdom

^c Applied Sports Technology, Exercise Medicine Research Centre (A-STEM), Swansea University, Swansea, United Kingdom.

^d Welsh Institute of Performance Science, Faculty of Science and Engineering, Swansea University, Swansea,

United Kingdom.

^e London Sports Institute, Middlesex University, Faculty of Science and Technology, London, United Kingdom

*Corresponding author: Dr Samuel P. Hills

Figures: 1

Tables: 1

Title: Alternate leg bounding acutely improves change of direction performance in women's team sports players irrespective of ground type

Running head: Bounding and change of direction

1 **ABSTRACT**

2 This study aimed to assess whether post-warm-up body mass only alternate leg bounding performed on
3 grass or a hard surface acutely improves pre-planned change of direction performance in women’s team
4 sports players relative to a control condition and, if so, profile the time-course of such changes. On three
5 occasions, 14 amateur women’s team sports players performed 20 m pre-planned change of direction
6 (‘Pro-Agility’) tests at 4 min, 8 min, and 12 min following interventions. Interventions were
7 implemented immediately after a standardized warm-up and consisted of three sets of 10 repetitions of
8 alternate leg bounding (five ground contacts per limb) on a hard indoor surface (HARD) or natural grass
9 (GRASS), or a control condition involving ~75 s of continuous walking with no bounding (CON).
10 Performance was similar between conditions at 4 min post-intervention. Performance at 8 min was
11 greater in HARD (2.9%, $p = 0.015$), and GRASS (3.8%, $p = 0.029$) relative to CON, whilst GRASS
12 also exceeded CON at 12 min post-bounding (5.2%, $p = 0.004$). All effects were large. No differences
13 existed between HARD and GRASS at any timepoint. Alternate leg bounding performed with body
14 mass only can acutely improve indices of change of direction performance in women’s team sports
15 players irrespective of the ground surface when an appropriate post-stimulus recovery period is
16 provided. Bounding on grass or a hard surface represents a feasible match-day practice that enhances
17 subsequent change of direction performance and could therefore be used as part of practically applicable
18 pre-match, half-time, and/or pitch-side (re)warm-up activities.

19

20 **KEY WORDS:** Running, plyometric, football, power, agility, warm-up

21

22

23

24

25

26 **INTRODUCTION**

27 The capacity for muscular force production is influenced by the recent contractile history of a given
28 muscle group (15). If potentiating effects exceed any co-existing fatigue, performing certain high-
29 intensity muscle actions as a conditioning stimulus can acutely enhance subsequent exercise outcomes
30 during explosive tasks such as jumps and sprints (27, 28, 31). This acute and temporary performance
31 improvement, postactivation performance enhancement (PAPE), has been attributed to mechanisms
32 including increased actin-myosin myofilament sensitivity to Ca^{2+} , enhanced motor neuron recruitment,
33 increased body temperature, and/or more favourable central input to the motor neuron (2, 27, 31).

34

35 Many team sports such as the football codes are characterized by intermittent bouts of high-speed
36 activity such as sprinting, jumping, and changing direction (25). Acknowledging that myriad factors
37 contribute to overall team success, explosive physical actions are involved in many of the most decisive
38 passages of play (6, 8). A player's ability to combine sprinting with rapid changes of direction, either
39 as a pre-planned manoeuvre or in response to a stimulus, is an important indicator of physical
40 performance (9, 22, 23) and can discriminate between playing levels in men's and women's soccer (17).
41 The potential to acutely enhance high-intensity exercise performance means that targeting PAPE could
42 be a worthwhile pre- or during-competition strategy to improve elements of sport-specific physical
43 performance for team sports players.

44

45 Studies have often used moderate to heavy resistance exercise to elicit PAPE (27, 31). The greatest
46 benefits have typically been reported in trained individuals using multi-set routines when ~7-10 min of
47 rest separates the conditioning stimulus and subsequent exercise (15, 31). However, heavy resistance
48 exercise may not always be feasible or desirable to implement within the practical, logistical, and
49 regulatory constraints associated with pre- and within-competition practices of many team sports
50 players. Identifying alternative methods of inducing PAPE that require less equipment and/or might be

51 better tolerated by players and coaches on the day of competition may allow this strategy to be more
52 widely implemented in practice.

53

54 Maximal isometric contractions may activate more motor units than dynamic movements and can
55 therefore elicit PAPE in certain contexts (7, 10, 27), whilst ballistic and plyometric activities such as
56 weighted jumps and throws may also be used as a conditioning stimulus due to their preferential
57 recruitment of type II motor units (5, 11, 30). Turner et al. (28) observed that three sets of 10 repetitions
58 of alternate-leg bounding on a hard indoor surface whilst wearing a weighted vest (equivalent to 10%
59 of body mass) improved 10 m and 20 m sprint performance by 2-3% at 4 min and 8 min post-bounding,
60 compared with a walking control condition. It would be valuable to ascertain whether the benefits of
61 bounding can transfer to other important indices of team sports specific physical performance (e.g.,
62 movement sequences incorporating changes of direction), especially given that an isometric squat
63 protocol has failed to enhance pre-planned change of direction outcomes in men's academy rugby
64 players (16).

65

66 As plyometric exercises are characterized by rapid transfer from eccentric to concentric muscle actions
67 and involve high ground reaction forces (4), the ground surface on which these movements are
68 performed can affect the kinematics and physiological responses to such activities (18, 21). Whilst
69 existing findings (28) suggest a potential benefit to incorporating bounding exercise within a pre-match,
70 half-time, or pre-pitch-entry active warm-up for improving acceleration performance in plyometrically
71 trained men's team sports players, it remains to be determined whether favourable responses also occur
72 on softer surfaces more relevant to outdoor match-play (e.g., natural grass). Moreover, as very limited
73 research has assessed PAPE in female athletes, exploring the efficacy of similar feasible conditioning
74 strategies in women's team sports players would help to determine the value of such strategies for this
75 population. This is especially relevant given that individual characteristics such as strength, speed,
76 training experience, and proportion of type II muscle fibres may each influence the magnitude of the

77 PAPE response (10, 15, 31). Therefore, the aim of this study was to assess whether plyometric bounding
78 exercise performed on natural grass or a hard surface with body mass only loading acutely improves
79 subsequent change of direction performance in women's team sports players relative to a control
80 condition and to determine the timeframe over which any PAPE may occur.

81

82 **METHODS**

83 *Experimental approach to the problem*

84 In a randomized, counterbalanced, cross-over fashion, participants completed three trials with
85 approximately seven days between trials. On arrival at the testing venue on the day of each trial,
86 participants completed a ~15 min standardized active warm-up which involved jogging (~5 min,
87 moderate intensity) and lower-body dynamic stretching (~7 min, focusing on the musculature involved
88 in the subsequent bounding and change of direction activities), before concluding with sprinting and
89 changing direction at increasing intensities (~3 min, involving 10 m sprints including 180° turns at near
90 maximal intensities). The intervention stimulus followed immediately thereafter, which consisted of
91 either a: a) walking control condition (CON), b) bounding on a hard surface condition (HARD), or c)
92 bounding on grass condition (GRASS). A 20 m 'Pro-Agility test' was completed at 4 min, 8 min, and
93 12 min post-intervention.

94

95 *Subjects*

96 Following Bournemouth University ethics approval, 14 amateur standard women's team sports players
97 (age: 20 ± 1 years, mass: 62.9 ± 7.6 kg, stature: 1.66 ± 0.06 m) volunteered to participate. *A priori*
98 sample size calculation was completed using commercially available software (G*Power; Version
99 3.9.1.2, Germany). With an anticipated large effect size (28) and alpha set at 0.05, a sample of 12 was
100 deemed sufficient for at least 80% power to detect significant effects. Participants were informed of the
101 risks and benefits of participation and provided written consent before data collection. Eligibility

102 required that participants had at least one year of plyometric training experience (i.e., performed specific
103 plyometric exercises on average at least once per week over this period) and were active team sports
104 players.

105

106 *Procedures*

107 Participants attended two familiarization sessions before the first trial to ensure familiarity with all
108 exercise and testing procedures, which included performing bounding and multiple repetitions of the
109 ‘Pro-Agility test’ with maximal effort. Trials were completed at the same time of day on each occasion
110 to avoid the influence of diurnal variation in performance (26). For all trials, participants were asked to
111 avoid alcohol, caffeine, or strenuous exercise in the preceding 24 hours and maintain consistent
112 nutrition, hydration, and footwear on each occasion.

113

114 For HARD and GRASS, the standardized warm-up was followed immediately by three sets of 10
115 repetitions (i.e., five ground contacts per leg per set) of alternate leg bounding with no additional loading
116 applied other than body mass. Participants were instructed to perform the bounding as per Turner et al.
117 (28). After a three-step run-up, participants pushed off their preferred foot before flexing the hip to
118 bring the opposite limb through so the thigh was approximately parallel to the ground with the knee
119 flexed to $\sim 90^\circ$. Hip and knee extension followed to forcefully contact the ground with the foot and push
120 off, before participants repeated this sequence until 5 contacts were completed on each leg. A 15 s active
121 recovery separated each set. Participants were instructed to maximize distance per bound whilst
122 minimizing ground contact time. The only difference between HARD and GRASS was that bounding
123 in HARD was performed on a hard indoor sports hall surface, whereas bounding in GRASS was
124 performed on a flat natural grass surface which had not been exposed to precipitation within the
125 preceding 24 h. The warm-up in CON was followed by continuous walking for the equivalent duration
126 of the bounding intervention in HARD and GRASS (~ 75 s). Walking was included rather than passive
127 rest to minimize losses of warm-up induced body temperature in CON relative to HARD and GRASS.

128

129 At 4 min, 8 min, and 12 min after the respective intervention, change of direction ability was assessed
130 using a 'Pro-Agility test' (Figure 1). This test was selected because it was anticipated to elicit minimal
131 fatigue per repetition, combined acceleration, deceleration, and changes of direction, was similar to
132 activities performed by many of the participants in sport-specific training, and has demonstrated good
133 reliability (coefficient of variation; CV% = 1.8) in recreational standard women's team sports athletes
134 (24). The current sample demonstrated test-retest CV% = 1.3 following familiarization. Participants
135 began each repetition stationary in a neutral stance 0.3 m behind the start line. On hearing a verbal start
136 command, participants were required to turn 90° to sprint to touch with their foot a line 5 m to their
137 right. Having reached the line, participants changed direction (180° change) and sprinted to a line 10 m
138 in the opposite direction, before a further 180° change of direction and 5 m sprint back to the start line
139 (24). A single repetition was performed at each timepoint and time taken to complete the 20 m course
140 was recorded using electronic timing gates (Brower Timing Systems, USA) at a height of approximately
141 0.8 m. All procedures for CON and HARD were performed in a temperature-controlled indoor sports
142 hall (air temperature: 18.7 ± 0.6 °C, relative humidity: $51.3 \pm 0.9\%$), whereas the bounding in GRASS
143 was performed outdoors on an area of natural grass immediately adjacent to the sports hall entrance.
144 The standardized warm-up, recovery periods, and testing in GRASS were completed in the indoor sports
145 hall.

146

147 ****INSERT FIGURE 1 HERE****

148

149 *Statistical analyses*

150 Statistical analyses were conducted using SPSS software (Version 28; SPSS Inc, USA) and $p < 0.05$ was
151 used as the threshold for statistical significance. Following checks for normality of distribution, two-
152 way analysis of variance with repeated measures was used, with condition (CON, HARD, GRASS) and
153 time (4 min, 8 min, 12 min) representing within-participant factors alongside their interaction.

154 Mauchly's test was consulted and the Greenhouse-Geisser correction was applied if the assumption of
155 sphericity was violated. Significant main effects were explored using Bonferroni-adjusted pairwise
156 comparisons, whilst significant condition x time interactions were broken down via simple effects
157 analysis. Hedge's g effect sizes (ES) were calculated for significant comparisons and were interpreted
158 as trivial (0.00–0.19), small (0.20– 0.49), moderate (0.50–0.79), or large (>0.80) (3).

159

160 **RESULTS**

161 Table 1 shows change of direction performance in each condition. There were significant effects of
162 condition ($F_{(2, 26)} = 9.907$, $p < 0.001$, partial eta-squared = 0.432), time ($F_{(1.36, 17.72)} = 10.496$, $p = 0.002$,
163 partial eta-squared = 0.447), and a significant condition x time interaction ($F_{(4, 52)} = 3.958$, $p = 0.007$
164 partial eta-squared = 0.233). No difference between conditions existed at 4 min post-intervention but at
165 8 min performances in HARD (2.9%, $p = 0.015$, ES:1.17, *large*) and GRASS (3.8%, $p = 0.029$, ES:
166 1.18, *large*) were superior to CON. At 12 min, times in GRASS, but not HARD, remained faster than
167 CON (5.2%, $p = 0.004$, ES: 1.64, *large*). Results were similar between HARD and GRASS throughout.

168

169 In CON, performance at 12 min was worse than that recorded at 4 min post-walking ($p = 0.018$, ES:
170 1.09, *large*). In HARD, times were faster at 8 min relative to both 4 min ($p = 0.029$, ES: 0.46, *small*)
171 and 12 min ($p = 0.002$, ES: 0.85, *large*) post-bounding.

172

173 ****INSERT TABLE 1 HERE****

174

175 **DISCUSSION**

176 This study assessed whether alternate leg bounding on grass or a hard surface acutely improved change
177 of direction performance in women's team sports players. On both surfaces, completing three sets of 10

178 repetitions of body mass only alternate leg bounding improved pre-planned change of direction times
179 at 8 min post-intervention compared with a walking control condition. Bounding on natural grass also
180 elevated performance relative to the control after 12 min. These findings indicate that alternate leg
181 bounding can acutely improve indices of change of direction performance in women's team sports
182 players when ~8-12 min recovery is provided. Such data highlight the potential for this strategy to be
183 incorporated where feasible into the match-day practices of this athletic population. Depending on the
184 length of any post-warm-up transition period, bounding may be implemented as part of a pre-match
185 warm-up, half-time rewarm-up, and/or pitch-side preparations for substitutes awaiting pitch-entry.
186 Moreover, this strategy may be implemented at the beginning of training sessions that are targeted at
187 improving physical capabilities such as speed and change of direction ability.

188

189 At 8 min post-intervention, change of direction performance was elevated by 2.9% and 3.8% in HARD
190 and GRASS, respectively, compared with the same timepoint in CON. Whilst improvements in
191 explosive physical performance following an appropriate conditioning stimulus have been well
192 established (16, 27, 31), conditioning protocols have often involved using external equipment to
193 facilitate appropriate loading. Given the practical and regulatory constraints that often exist on match-
194 day, these strategies are unlikely to be feasible immediately before or during a match in many team
195 sports. Plyometric or ballistic activities may offer more a practical alternative conditioning stimulus
196 when compared with traditional methods of inducing PAPE (5, 11, 30). Improvements in change of
197 direction performance after 8 min in the current study are of similar magnitude to previous observations
198 of augmented 10 m and 20 m straight line sprint performance after trained men completed alternate leg
199 bounding whilst wearing a weighted vest (with an additional 10% of body mass of loading) (28). This
200 study therefore extends previous research to demonstrate a bounding-induced PAP effect on another
201 crucial aspect of team sports match-play (i.e., change of direction) and without the need for any
202 additional equipment.

203

204 Soft surfaces such as sand can dissipate ground reaction forces and reduce stretch-shortening cycle
205 efficiency compared with the same activities performed on hard ground (18). Moreover, increased
206 shock absorption and vertical deformation have been observed when running on natural grass relative
207 to a hard asphalted running track (21). As exercises such as alternate leg bounding are characterized by
208 rapid transfer from eccentric to concentric muscle actions and involve high ground reaction forces (4),
209 the surface on which these activities are performed could plausibly influence the length of the
210 amortization phase and thus the extent to which they have can elicit PAPE. However, no statistically
211 significant differences in change of direction performance existed between HARD and GRASS at any
212 timepoint. These findings suggest that alternate leg bounding performed on natural grass may be at least
213 as effective for improving this performance outcome as the same intervention performed on a hard
214 indoor surface. Moreover, the benefit relative to CON in GRASS extended to 12 min post-intervention
215 whereas improvements in HARD were restricted to the 8 min timepoint. Although the reasons for this
216 finding remain unclear, greater muscle activation has been reported during movements on softer
217 compared with harder surfaces (18). Therefore, it is possible that the likely slightly softer natural surface
218 may have led to greater muscle activation and thus a more sustained performance enhancement in
219 GRASS than in HARD. Alternatively, or in conjunction, because exercises on grass have a greater
220 metabolic cost than those performed on a hard surface (21), the bounding in GRASS could have led to
221 a greater and/or more sustained elevation of muscle temperature than in both HARD and CON. Given
222 the positive relationship between body temperature and explosive physical performance capacity (12,
223 19, 20), any such elevations could have contributed to significant performance elevations relative to
224 CON at 12 min in GRASS but not in HARD.

225

226 Movement specificity alongside the intensity of the prior contraction may be an important factor
227 influencing the PAPE response following a conditioning stimulus, whilst greater familiarity with a task
228 could reduce 'warm-up decrement' (29). Turner et al. (28) reported greater improvements in 10 m and
229 20 m sprint performance at 4 min and 8 min following bounding with a weighted vest compared with
230 the same volume of bounding performed without additional external loading. The authors speculated

231 that increased ground contact time during each bound in the weighted condition had greater
232 biomechanical specificity to the acceleration phase of sprinting. Similar considerations may have
233 contributed to the performance improvements (i.e., relative to CON) persisting at 12 min post-bounding
234 in GRASS if greater shock absorption on the natural grass surface led to longer ground contact time in
235 this condition than in HARD (21). Indeed, short-duration acceleration and deceleration as required in
236 the change of direction test involve maximizing horizontal orientation of forces (1, 13). Further research
237 is needed to elucidate whether greater improvements in change of direction performance could be
238 elicited by adding external loading (e.g., via a weighted vest or horizontally applied resistance) or
239 incorporating conditioning movements in the frontal plane, and whether this bounding stimulus can
240 simultaneously improve other valuable physical performance indicators (e.g., jumping activities with a
241 more vertical orientation of force).

242

243 As well as the intervention itself (e.g., the type, duration, and intensity of exercise performed, alongside
244 the subsequent recovery duration), characteristics of the individual participant can influence the PAPE
245 response. Indeed, possessing superior strength, speed, training experience, and proportion of type II
246 muscle fibres may allow an athlete to benefit from PAPE to a greater extent than those who are weaker
247 or less well-trained (10, 15, 31). Whilst differences in PAPE between males and females have not been
248 confirmed in the literature, current evidence is limited and many relevant characteristics typically vary
249 between sexes (14). The fact that this study observed PAPE of change of direction performance relative
250 to a control condition in recreational standard female athletes with typically ~1-2 years of plyometric
251 training experience is therefore an important and novel observation. It is possible that greater benefits
252 may have been experienced if participants had been more highly trained athletes. Notably, the more
253 well-trained an individual, the greater volume and/or intensity of conditioning stimulus may be required
254 to maximize PAPE (31).

255

256 Whilst this study demonstrated improved 'Pro-Agility test' performance in HARD and GRASS relative
257 to CON, it is not possible to determine whether the bounding interventions elicited improvements in
258 the specific change of direction component or the acceleration/deceleration component of the test. In
259 addition, because physiological and electromyographical measurements were not taken, the precise
260 mechanism(s) underpinning the acute performance enhancement observed cannot be conclusively
261 determined. Nonetheless, this study has shown that a ~75 s long alternate leg bounding intervention
262 completed with body mass only on either natural grass or an indoor hard surface can acutely enhance
263 change of direction test performance in women's team sports players compared with a walking control
264 condition.

265

266 This study involved between-condition comparisons, without including a pre-intervention baseline
267 measurement. The findings must be interpreted as such. The lack of baseline measurement means that
268 it is not possible to determine within-condition performance changes relative to pre-intervention.
269 However, as the change of direction test was consistent across all three trials, the design of the current
270 study avoids any potential confounding influence from the test itself producing a fatiguing and/or
271 potentiation effect at subsequent timepoints (i.e., performances at 8 min and 12 min had been preceded
272 in all trials by one and two repeats of the change of direction test, respectively). Performance in HARD
273 and GRASS could thus be compared with responses produced in the absence of any bounding
274 intervention (i.e., CON).

275

276 **PRACTICAL APPLICATIONS**

277 In women's team sports players, completing three sets of 10 repetitions of body mass only alternate leg
278 bounding on either a hard indoor surface or natural grass elicited improvements in change of direction
279 performance 8 min post-intervention when compared with a walking control condition. Improvements
280 relative to the control were also seen at 12 min post-bounding when bounding was performed on grass.
281 These findings suggest that players and coaches may consider implementing alternate leg bounding at

282 specific timepoints during training or on match-day (e.g., at the end of the pre-match warm-up, at half-
283 time, or for substitutes awaiting pitch-entry) as a means of potentially enhancing indices of sport-
284 specific physical performance even if a suitable hard surface is not immediately available.

285

286 ACKNOWLEDGMENTS

287 No financial assistance was received for this study. The authors report that there are no conflicts of
288 interest to declare. The results of the present study do not constitute endorsement of the product by the
289 authors or the National Strength and Conditioning Association.

290

291 REFERENCES

- 292 1. Baena-Raya A, Soriano-Maldonado A, Conceição F, Jiménez-Reyes P, and Rodríguez-Pérez
293 MA. Association of the vertical and horizontal force-velocity profile and acceleration with
294 change of direction ability in various sports. *Eur J Sport Sci* 21: 1659-1667, 2021.
- 295 2. Blazevich AJ and Babault N. Post-activation potentiation versus post-activation performance
296 enhancement in humans: historical perspective, underlying mechanisms, and current issues.
297 *Front Physiol* 10: 1359, 2019.
- 298 3. Cohen J. Statistical power analysis. *Curr Dir Psychol Sci* 1: 98-101, 1992.
- 299 4. Davies G, Riemann BL, and Manske R. Current concepts of plyometric exercise. *Int J Sports*
300 *Physiol Perform* 10: 760, 2015.
- 301 5. Desmedt JE and Godaux E. Ballistic contractions in man: characteristic recruitment pattern of
302 single motor units of the tibialis anterior muscle. *J Physiol* 264: 673-693, 1977.
- 303 6. Dos' Santos T, McBurnie A, Thomas C, Jones PA, and Harper D. Attacking agility actions:
304 Match play contextual applications with coaching and technique guidelines. *Strength Cond J*:
305 epub ahead of print, 2022.
- 306 7. Duchateau J and Hainaut K. Isometric or dynamic training: differential effects on mechanical
307 properties of a human muscle. *J Appl Physiol* 56: 296-301, 1984.
- 308 8. Faude O, Koch T, and Meyer T. Straight sprinting is the most frequent action in goal
309 situations in professional football. *J Sports Sci* 30: 625-631, 2012.
- 310 9. Gabbett TJ, Kelly JN, and Sheppard JM. Speed, change of direction speed, and reactive
311 agility of rugby league players. *J Strength Cond Res* 22: 174-181, 2008.
- 312 10. Hamada T, Sale DG, MacDougall JD, and Tarnopolsky MA. Interaction of fibre type,
313 potentiation and fatigue in human knee extensor muscles. *Acta Physiol Scand* 178: 165-173,
314 2003.
- 315 11. Hilfiker R, Hubner K, Lorenz T, and Marti B. Effects of drop jumps added to the warm-up of
316 elite sport athletes with a high capacity for explosive force development. *J Strength Cond Res*
317 21: 550-555, 2007.
- 318 12. Hills SP, Aben HGJ, Starr DP, et al. Body temperature and physical performance responses
319 are not maintained at the time of pitch-entry when typical substitute-specific match-day
320 practices are adopted before simulated soccer match-play. *J Sci Med Sport* 24: 511-516, 2020.
- 321 13. Hunter JP, Marshall RN, and McNair PJ. Interaction of step length and step rate during sprint
322 running. *Med Sci Sports Exerc* 36: 261-271, 2004.

- 323 14. Hunter SK. The relevance of sex differences in performance fatigability. *Med Sci Sports*
324 *Exerc* 48: 2247-2256, 2016.
- 325 15. Kilduff LP, Owen N, Bevan H, et al. Influence of recovery time on post-activation
326 potentiation in professional rugby players. *J Sports Sci* 26: 795-802, 2008.
- 327 16. Marshall J, Turner AN, Jarvis PT, et al. Postactivation potentiation and change of direction
328 speed in elite academy rugby players. *J Strength Cond Res* 33: 1551-1556, 2019.
- 329 17. Mujika I, Santisteban J, Impellizzeri FM, and Castagna C. Fitness determinants of success in
330 men's and women's football. *J Sports Sci* 27: 107-114, 2009.
- 331 18. Pereira LA, Freitas TT, Marín-Cascales E, et al. Effects of training on sand or hard surfaces
332 on sprint and jump performance of team-sport players: A systematic review with meta-
333 analysis. *Strength Cond J* 43: 56-66, 2021.
- 334 19. Russell M, Tucker R, Cook CJ, Giroud T, and Kilduff LP. A comparison of different heat
335 maintenance methods implemented during a simulated half-time period in professional rugby
336 union players. *J Sci Med Sport* 21: 327-332, 2018.
- 337 20. Sargeant AJ. Effect of muscle temperature on leg extension force and short-term power
338 output in humans. *Eur J Appl Physiol Occup Physiol* 56: 693-698, 1987.
- 339 21. Sassi A, Stefanescu A, Bosio A, Riggio M, and Rampinini E. The cost of running on natural
340 grass and artificial turf surfaces. *J Strength Cond Res* 25: 606-611, 2011.
- 341 22. Sekulic D, Pehar M, Krolo A, et al. Evaluation of basketball-specific agility: applicability of
342 preplanned and nonplanned agility performances for differentiating playing positions and
343 playing levels. *J Strength Cond Res* 31: 2278-2288, 2017.
- 344 23. Sheppard J, Young WB, Doyle T, Sheppard T, and Newton RU. An evaluation of a new test
345 of reactive agility and its relationship to sprint speed and change of direction speed. *J Sci Med*
346 *Sport* 9: 342-349, 2006.
- 347 24. Stewart PF, Turner AN, and Miller SC. Reliability, factorial validity, and interrelationships of
348 five commonly used change of direction speed tests. *Scand J Med Sci Sports* 24: 500-506,
349 2014.
- 350 25. Taylor JB, Wright AA, Dischiavi SL, Townsend MA, and Marmon AR. Activity demands
351 during multi-directional team sports: a systematic review. *Sports Med* 47: 2533-2551, 2017.
- 352 26. Teo W, McGuigan MR, and Newton MJ. The effects of circadian rhythmicity of salivary
353 cortisol and testosterone on maximal isometric force, maximal dynamic force, and power
354 output. *J Strength Cond Res* 25: 1538-1545, 2011.
- 355 27. Tillin NA and Bishop D. Factors modulating post-activation potentiation and its effect on
356 performance of subsequent explosive activities. *Sports Med* 39: 147-166, 2009.
- 357 28. Turner AP, Bellhouse S, Kilduff LP, and Russell M. Postactivation potentiation of sprint
358 acceleration performance using plyometric exercise. *J Strength Cond Res* 29: 343-350, 2015.
- 359 29. Verhoeven FM and Newell KM. Unifying practice schedules in the timescales of motor
360 learning and performance. *Hum Mov Sci* 59: 153-169, 2018.
- 361 30. West DJ, Dietzig BM, Bracken RM, et al. Influence of post-warm-up recovery time on swim
362 performance in international swimmers. *J Sci Med Sport* 16: 172-176, 2013.
- 363 31. Wilson JM, Duncan NM, Marin PJ, et al. Meta-analysis of postactivation potentiation and
364 power: effects of conditioning activity, volume, gender, rest periods, and training status. *J*
365 *Strength Cond Res* 27: 854-859, 2013.

366

367

368

369

370

LEGENDS:

Figure 1. Overview of Pro-Agility test procedures

Table 1. Change of direction performance times (s) by condition and timepoint following intervention

Table 1. Change of direction performance times (s) by condition and timepoint following intervention

	4 min	8 min	12 min
CON	5.72 ± 0.10	5.76 ± 0.15	5.91 ± 0.22
HARD	5.66 ± 0.13	5.60 ± 0.12 *	5.71 ± 0.14
GRASS	5.58 ± 0.15	5.54 ± 0.21 *	5.60 ± 0.14 *

CON: Control condition (walking), GRASS: Bounding intervention on a natural grass surface, HARD: Bounding intervention on a hard indoor surface
Data are presented as mean ± standard deviation. *: Statistically significantly different from the equivalent timepoint in CON (p <0.05, large effect size).